

Diagonalization in proof complexity

Jan Krajíček*

Mathematical Institute[†]
Academy of Sciences, Prague

Abstract

We study the diagonalization in the context of implicit proofs of [10]. We prove that at least one of the following three conjectures is true:

- There is a function $f : \{0, 1\}^* \rightarrow \{0, 1\}$ computable in \mathcal{E} that has circuit complexity $2^{\Omega(n)}$.
- $\mathcal{NP} \neq \text{co}\mathcal{NP}$.
- There is no p -optimal propositional proof system.

We note that a variant of the statement (either $\mathcal{NP} \neq \text{co}\mathcal{NP}$ or $\mathcal{NE} \cap \text{co}\mathcal{NE}$ contains a function $2^{\Omega(n)}$ hard on average) seems to have a bearing on the existence of good proof complexity generators. In particular, we prove that if a minor variant of a recent conjecture of Razborov [17, Conjecture 2] is true (stating conditional lower bounds for the Extended Frege proof system EF) then actually unconditional lower bounds would follow for EF .

The only method for demonstrating unprovability of a Π_1^0 -sentence in a theory (containing some amount of arithmetic) is the diagonalization. One would like to adapt this to a non-uniform setting in which theories are replaced by propositional proof systems and Π_1^0 -sentences by tautologies. This fails, at least in the straightforward adaptation. In particular, many strong proof systems do prove their own consistency by polynomial size

*Partially supported by grant # A 101 94 01 of the Academy of Sciences of the Czech Republic and by project LN00A056 of The Ministry of Education of the Czech Republic.

[†]Also member of the Institute for Theoretical Computer Science of the Charles University. A part of this work was done while visiting the Mathematical Institute at Oxford.

proofs (cf.[5, Chpt.14]). However, there is a non-uniform setting in first-order theories where the diagonalization gives non-trivial lower bounds.

To explain this we need to fix some notation first. By a theory we shall mean a set of axioms, not the set of its consequences. We shall assume (w.l.o.g.) that the language of all theories in question is the language of bounded arithmetic theory S_2^1 . For a natural number N the symbol \underline{N} is the dyadic numeral inductively defined by: $\underline{0} := 0$, $\underline{1} := 1$, $\underline{2} := (1 + 1)$, $\underline{2k} := (\underline{2} \cdot \underline{k})$, and $\underline{2k+1} := (\underline{2k} + 1)$.

Given a theory T , let $Prf_T(y, x)$ be a formula in the language of S_2^1 expressing that y is a T -proof of formula x . Assuming that $T \supseteq S_2^1$, as we shall do, there is a canonical formalization of syntax of logic (terms, formulas, proofs, etc.) and of the notion of provability (see [13] or [5] for details), and its usual properties are provable in S_2^1 .

Define a diagonal formula $A(x)$ satisfying:

$$A(x) \equiv \forall y, |y| \leq s(x) \longrightarrow \neg Prf_T(y, \lceil A(\dot{x}) \rceil).$$

where $s(x)$ is any term and $\lceil A \rceil$ is a (canonical) number encoding A such that its length is proportional to the size of A , and \dot{x} is the formalization of dyadic numerals. Then, by the standard argument, we have the following theorem. (The condition that $T \in \mathcal{NP}$ is technical and implies, in particular, that T is definable and that the binary relation $Prf_T(y, x)$ is in \mathcal{NP} and definable by a formula, and that all its true instances have polynomial size proofs in S_2^1 .)

Theorem 0.1 *Let $T \supseteq S_2^1$ be a consistent theory such that $T \in \mathcal{NP}$. Then for any $N \geq 1$ the sentence $A(\underline{N})$ is true and provable in T but any T -proof of the sentence must have the size at least $s(N)$.*

Note that the length of the formula $A(\underline{N})$ is $O(\log N)$ and hence the lower bound is non-trivial, as long as $s(N) \gg \log(N)$. A foremost example of this reasoning is the Finitistic Gödel's theorem of Friedman [3] and Pudlák [13, 14] (the theorem says that any T -proof of a formula $Con_T(\underline{N}) := \forall y, |y| \leq \underline{N} \rightarrow \neg Prf(y, \lceil 0 = 1 \rceil)$ expressing the consistency of T w.r.t. proofs of length at most N , must have the size at least $N^{\Omega(1)}$). Other, quantitatively more subtle applications, can be found in bounded arithmetic, cf.[5, Chpt.10].

One would like to adapt this to propositional proof complexity. The problem with this is not that we deal with first-order theories (as we know how to pass between them and propositional proof systems) but with the fact

that $A(x)$ is not a bounded formula and the instances $A(\underline{N})$ translate into propositional formulas of length at least $s(N)$, which is super-polynomial in order to get non-trivial lower bound. An idea suggests itself at this point: Try to use the concept of implicit proofs from [10]. This is what we investigate in this paper. A variant of the idea of implicit proofs is recalled in Section 1. In Section 2 we derive the theorem mentioned in the abstract. Its variant is derived in Section 3 and linked to proof complexity generators in Section 4.

The paper is self-contained, assuming that the reader has a general background in bounded arithmetic and proof complexity (only basic things are assumed), see [5]. Let us just recall the terminology and some more notation. $Time(t(n))$ is the class of languages computable in deterministic time $O(t(n))$, $\Sigma_i Time(t(n))$ is the Σ_i -level of the time $O(t(n))$ hierarchy, $NTime(t(n)) = \Sigma_1 Time(t(n))$, and $\mathcal{E} = Time(2^{O(n)})$. For a function $f : \{0,1\}^* \rightarrow \{0,1\}$, $C_f(n)$ is the circuit complexity of computing f on $\{0,1\}^n$, while $H_f(n)$ is its hardness on average in the sense of [12]. A “proof system” tacitly means a “propositional proof system”: It is a non-deterministic acceptor of the set of propositional tautologies in the DeMorgan language. A proof of a tautology in a proof system is any particular computation of the proof system accepting the formula. A proof system is p -bounded iff the proof system accepts all tautologies in a fixed polynomial time, and it is p -optimal iff it polynomially simulates other proof systems, cf.[2]. The length of a string w is denoted $|w|$, and a number is identified with the string of its bits (i.e. $|m| \sim \log(m)$).

Finally let us recall a well-known translation of formulas into propositional formulas, just for the case of $A(x)$. Given $N \geq 1$, there is a propositional 3DNF formula denoted $\|A(x)\|_N$ of size $s(N)^{O(1)}$ that expresses (by the fact of being a tautology) that $A(N)$ is true. It is constructed as in the proof of the \mathcal{NP} -completeness of the satisfiability. The formula has $s(N)$ atoms for bits of a potential y and auxiliary atoms ($s(N)^{O(1)}$ of them) used for values of subcircuits used in the canonical computation of the truth value of $Prf_T(y, \lceil A(\underline{N}) \rceil)$, and says that if all local conditions in the computation are satisfied then $Prf_T(y, \lceil A(\underline{N}) \rceil)$ fails. See [5, Chpt.9] for details of the translation.

1 Implicit proofs of implicit formulas

Let w be a 0-1 string of length 2^k . Identify $i < 2^k$ with vectors $i = (i_1, \dots, i_k) \in \{0, 1\}^k$ ordered lexicographically. We say that a circuit $C(x_1, \dots, x_k)$ represents w iff $C(i_1, \dots, i_k) = w_i$ (the i th bit of w), for all $i < 2^k$. Similarly, if W is a 0-1 $2^k \times 2^k$ matrix then we say that a circuit $D(x_1, \dots, x_k, y_1, \dots, y_k)$ represents W iff $D(i, j) = W_{i,j}$, for all $i, j < 2^k$.

Let M be a non-deterministic polynomial time machine. For any input w of length 2^ℓ and represented by a circuit $C(z_1, \dots, z_\ell)$, for suitable $k = O(\ell)$ (given by the time of M) and any $2^k \times 2^k$ matrix W represented by a circuit $D(x_1, \dots, x_k, y_1, \dots, y_k)$ we can write down a propositional formula $\sigma_{C,D}^M$ expressing that W is an accepting computation of M (given by W listing in its rows all instantaneous descriptions of the computation) on input w . The formula has atoms $x_1, \dots, x_k, y_1, \dots, y_k, z_1, \dots, z_\ell$ and auxiliary atoms for values of subcircuits of C and D , and formalizes that all local conditions posed on W by M are met and that the input as given in W is indeed w . The meaning of the verb “expressing” is that $\sigma_{C,D}^M$ is a tautology iff W is indeed an accepting computation of M on w .

We will need one additional formula, this time first-order. Assume that a circuit $C(x_1, \dots, x_k)$ represents a 3DNF formula φ_C (of size at most 2^k). Let $[C]$ be its number code. Then there is a formula $BigTaut(x)$ such that $BigTaut([C])$ formalizes that φ_C is a tautology.

2 A theorem

Theorem 2.1 *At least of one the following three statements is true:*

- (i) *There is a function $f : \{0, 1\}^* \rightarrow \{0, 1\}$ computable in \mathcal{E} that has circuit complexity $2^{\Omega(n)}$.*
- (ii) *$\mathcal{NP} \neq \text{co}\mathcal{NP}$.*
- (iii) *There is no p -optimal propositional proof system.*

Proof :

We shall assume that all three statement are false and derive a contradiction via Theorem 0.1.

(1) Let P be a proof system witnessing that both (ii) and (iii) fail: P is p -bounded and also p -optimal. Assume w.l.o.g. that P contains EF . Let $Prf_P(u, v)$ be a Σ_1^b -formula formalizing that “ u is a P -proof of formula v ”.

(2) Define theory T to be S_2^1 augmented by an extra axiom, a form of reflection principle:

$$\forall x, y, u, \text{Prf}_P(u, [\sigma_{x,y}^P]) \longrightarrow \text{BigTaut}(y) .$$

(3) Let $A(x)$ be the diagonal formula from the introduction, with the term $s(x)$ being simply x . Consider the propositional formula $\|A(x)\|_N$. The formula is a tautology as $A(N)$ is true. The size of the formula is $N^{O(1)}$ but there is clearly a circuit C_N of size $n^{O(1)}$, $n := \log(N)$, representing the formula (in the sense of Section 1).

(4) The set of all formulas $\|A(x)\|_N$, $N \geq 1$, is polynomial time decidable and hence we can use it as axioms in some proof system. By the hypothesis that P is p -optimal there is a (deterministic) polynomial time algorithm M computing from the string $\|A(x)\|_N$ a P -proof of $\|A(x)\|_N$.

(5) The output of M is a particular accepting computation of P , i.e. an $2^{O(n)} \times 2^{O(n)}$ matrix W^N encoding the computation. As M runs in deterministic polynomial time, $W_{i,j}^N$ as a function of $i, j \in \{0, 1\}^{O(n)}$ is in \mathcal{E} .

(6) Assuming that also statement (i) fails, there exists a circuit $D(i, j)$ in 2-times $O(n)$ variables and of size $2^{\delta \cdot n}$ that represents W^N , for arbitrary small $\delta > 0$. We shall choose a particular δ in the step (12).

(7) Take an instance of the reflection principle by substituting for x and y the codes of D and C_N respectively:

$$\forall u, \text{Prf}_P(u, [\sigma_{[C_N], [D]}^P]) \longrightarrow \text{BigTaut}([C_N]) .$$

(8) By Section 1 the size of $\sigma_{C_N, D}^P$ is polynomial in the sizes of C_N and D , i.e. it is $2^{O(\delta \cdot n)}$. Now we use the hypothesis that P is also p -bounded. Hence there is a P -proof e of $\sigma_{C_N, D}^P$ of size $2^{O(\delta \cdot n)}$. Note that the constants implicit in the O -notation are fixed and independent of δ . Substituting $[e]$ for u in the formula in (7) we get:

$$\text{Prf}_P([e], [\sigma_{[C_N], [D]}^P]) \longrightarrow \text{BigTaut}([C_N]) .$$

(9) The antecedent of the formula in (8) is a true Σ_1^b -sentence of size $2^{O(\delta \cdot n)}$ and has a proof in S_2^1 (and hence in T) of polynomial size, i.e. of size $2^{O(\delta \cdot n)}$.

(10) Applying modus ponens to the formulas in (8) and (9) we get a proof of size $2^{O(\delta \cdot n)}$ the sentence:

$$\text{BigTaut}([C_N]) .$$

(11) We claim that the implication:

$$BigTaut(\lceil C_N \rceil) \rightarrow A(\underline{N})$$

has an S_2^1 -proof of size $n^{O(1)}$. This is because it is an instance of a universal, S_2^1 -provable, statement saying that the translation of Π_1^b -sentences into propositional formulas is sound. Cf. [5, Chpt.9] for an analogous statement.

(12) Putting (10) and (11) together we get a size $2^{O(\delta \cdot n)}$ proof in T of $A(\underline{N})$. Taking $\delta > 0$ small enough so that $2^{O(\delta \cdot n)} < N$ (this we can do as the O -constant is independent of δ) we get a contradiction with Theorem 0.1.

q.e.d.

Let us remark that instead of using \mathcal{E} and circuit size $2^{\Omega(n)}$ in (i) we could have used $Time(t(n))$ and circuit size $t(n)^{\Omega(1)}$, as long as $t(n) \geq n^{\omega(1)}$. This follows by a padding argument or by a simple change to the proof above: Use the diagonal formula for term $s(x) := t(|x|)$ instead of $s(x) := x$ in the step (3).

3 A variant of the theorem

It is not difficult to see that the property of a string to be the truth table of a function on $\{0,1\}^n$ with $2^{\delta \cdot n}$ circuit complexity, or even $2^{\delta \cdot n}$ hard on average, is definable in the polynomial time hierarchy \mathcal{PH} . Taking the lexicographically first such strings (at least one exists of each length 2^n , $n \gg 0$, by a simple counting) we see that there is such an f computable in $\mathcal{E}^{\mathcal{PH}}$.

If $\mathcal{NP} = co\mathcal{NP}$ then such an f is in $\mathcal{E}^{\mathcal{NP} \cap co\mathcal{NP}} = \mathcal{NE} \cap co\mathcal{NE}$. If, in addition, $\mathcal{E} = \mathcal{NE}$ then such an f is in \mathcal{E} .

This simple (apparently folklore) argument yields the following theorem.¹ We shall use only Part 2 in Section 4 but we state also Part 1 as it is a weaker version (if hardness on average is replaced by circuit size) of Theorem 2.1 actually: It is known, by [11], that the non-existence of a p -optimal proof

¹This argument has been pointed out to me by E. Jeřábek, and has been also noted by V. Kabanets, and replaces my original proof: A simple modification of the proof of Theorem 2.1 shows that $\mathcal{NP} = co\mathcal{NP}$ implies that $\mathcal{NE} \cap co\mathcal{NE}$ contain a function with exponential circuit complexity which was then turned into a function with exponential hardness on average by the construction from [4].

system implies that $Time(t(n)) \neq NTime(t(n))$ (as long as $t(n) \leq 2^{n^{O(1)}}$) and hence also $\mathcal{E} \neq \mathcal{NE}$ (the opposite implication is unknown but Verbitsky [18] constructed a relativized world where it does not hold).

Theorem 3.1 *The following two proposition hold.*

1. *At least of one the following three statements is true:*

- (a) *There is a function $f : \{0,1\}^* \rightarrow \{0,1\}$ computable in \mathcal{E} that has $2^{\Omega(n)}$ hardness on average.*
- (b) $\mathcal{NP} \neq \text{coNP}$.
- (c) $\mathcal{E} \neq \mathcal{NE}$.

2. *At least of one the following two statements is true:*

- (a) *There is a function $f : \{0,1\}^* \rightarrow \{0,1\}$ computable in $\mathcal{NE} \cap \text{coNE}$ that has $2^{\Omega(n)}$ hardness on average.*
- (b) $\mathcal{NP} \neq \text{coNP}$.

4 Proof complexity generators

By a proof complexity generator we mean a map $g : \{0,1\}^n \rightarrow \{0,1\}^m$, $m = m(n)$ and $m > n$, whose bits can be computed in $NTime(m(n)^{O(1)}) \cap \text{coNTime}(m(n)^{O(1)})$. This assumption about the computability of the bits is the weakest one allowing to write down, for any $b \in \{0,1\}^m$, a size $m^{O(1)}$ propositional formula $\tau_b(g)$ that is a tautology iff $b \notin \text{Rng}(g)$, see below. (The notation $\tau_b(g)$ is somewhat misleading as the formula depends on a particular definition of g and not only on g , but we will ignore this here: The lower bounds conjectured later should hold for all $NTime(m(n)^{O(1)}) \cap \text{coNTime}(m(n)^{O(1)})$ -definitions.)

A generator is good if for a strong proof system P and with high probability in choosing a random $b \in \{0,1\}^m$, the formula $\tau_b(g)$ requires very long (in particular, super-polynomial) P -proofs. The quality of the generator is measured by the strength of P and by the probability that b yields a hard τ -formula. At present it is not ruled out that some generator g works for all P and all b . Following [17] we shall say that generator g is *hard for P* if all $\tau_b(g)$, for all b 's, require super-polynomial size P -proofs (cf.[17]).

The τ -formulas have been defined in [6] and independently in [1], and their theory has made first steps in [7, 16, 8, 17]. I shall not describe the

development of the ideas and known lower bound results; this can be found in the introductions to [8] or [17]. Instead I shall briefly describe one motivation and why we speak about "generators".

Proving lower bounds for strong propositional proof system appears hard. In fact, we do not know any such lower bounds. A factor contributing to this is that it is actually not easy to come up with sensible tautologies that would be good candidates for requiring long proofs even in strong systems (cf. [8] for a detailed discussion). The τ -formulas seem to be candidates worth studying in this context.

The word generator is used because some of the usual pseudo-random number generators seem to be good candidates. In particular, a good proof complexity generator must behave as a hitting set generator w.r.t. $\mathcal{NP}/poly$ -test (cf.[8]).

Similarly as the existence of good pseudo-random generators can be proved under some computational hardness assumptions (cf.[12, 4]) we may try to reduce the existence of good proof complexity generators to a suitable computational hardness assumption too. This is discussed in [9] with a broader perspective and in the introduction to [17].

The most studied map in this context is the classic Nisan-Wigderson generator, cf. [12]. This has been proposed as possibly a good proof complexity generator in [1] and taken up in [8], although the motivations (and, more importantly, the choice of parameters in the construction and the formalization of the notion of *hardness* of the generators²) are different. Let us first recall the definition of the NW-generators (and fix the notation in the process).

Let A be an $m \times n$ 0-1 matrix with ℓ ones per row. $J_i(A) := \{j \leq n \mid A_{ij} = 1\}$. Let $f : \{0,1\}^\ell \rightarrow \{0,1\}$ be a boolean function. $NW_{A,f} : \{0,1\}^n \rightarrow \{0,1\}^m$ is the NW-generator based on A and f : The i -th bit of output is computed by f from the bits of the input that belong to $J_i(A)$.

Assume that $f \in NTime(t(n)) \cap coNTime(t(n))$. Given particular $NTime(t(n))$ -definitions $\alpha: \exists v, \alpha_\epsilon(u, v)$ ($|u| = \ell$, $|v| \leq t(n)$ and α_ϵ p-time) of $f(u) = \epsilon$, for $\epsilon = 0, 1$, the τ -formulas are defined by:

$$\tau_b^\alpha := \bigvee_{i \leq m} \neg \alpha_{b_i}(x \downarrow J_i(A), v^i)$$

where $b = (b_1, \dots, b_m) \in \{0,1\}^m$, x is an n -tuple of variables and v^i are disjoint $t(n)$ -tuples of variables. Clearly $\tau_b \in TAUT$ iff $b \notin Rng(NW_{A,f})$.

²I shall describe neither the set-up from [8] nor the conjectured hardness in terms of pseudo-surjectivity here as it is not relevant to the topic of this paper.

Note that the size of τ_b^α is $t(n)^{O(1)} \cdot m(n)$, and hence we get a size $m^{O(1)}$ formula as long as $t(n) \leq m(n)^{O(1)}$.

An idea, formulated in [1] in general terms and then quite specifically in [17], is that $NW_{A,f}$ forms a good proof complexity generator, as long as A has suitable combinatorial properties (being an (ℓ, d) combinatorial design in the sense of [12]: $J_i(A)$'s have size ℓ and the intersection of any two different rows has size $\leq d$) and as long as f is computationally hard. Specifically, Razborov [17] has made the following conjecture.

Conjecture 4.1 (A. A. Razborov[17, Conjecture 2])

Any NW-generator based on a matrix A which is a combinatorial design with the same parameters as in [12] and on any function f in $\mathcal{NP} \cap \text{coNP}$ that is hard on average for \mathcal{P}/poly , is hard for EF .

Let us interpret the specifications. The parameters in the main construction of combinatorial designs in [12, L.2.5] satisfy: $d = \log(m)$, $\log(m) \leq \ell \leq m$ and $n = O(\ell^2)$.³ Writing this dually:

$$(1) \quad m = 2^{\epsilon \cdot n^{1/2}} \quad \text{and} \quad \ell = \epsilon \cdot n^{1/2}$$

where $\epsilon > 0$ is a constant (determined by the O -constant in the expression $n = O(\ell^2)$). We are taking the maximal value allowed for m by [12, L.2.5] as that is the chief case in [12], and it also links with [16, 17] studying “function” generators.

The phrase “hard on average for \mathcal{P}/poly ” presumably means that the hardness of f in the sense of [12] is exponential, similarly as it is needed in [12]:

$$(2) \quad H_f(\ell) \geq 2^{\Omega(\ell)} .$$

The conjecture requires that $f \in \mathcal{NP} \cap \text{coNP}$. We shall relax this condition to:

$$(3) \quad f \in \text{NTime}(2^{O(\ell)}) \cap \text{coNTime}(2^{O(\ell)}) .$$

By (1) we have $2^{O(\ell)} = m^{O(1)}$ so (3) means that f is in $\text{NTime}(m^{O(1)}) \cap \text{coNTime}(m^{O(1)})$. By the discussion earlier this means that the size of the τ -formula will be still $m^{O(1)}$. Thus this modification of the original formulation

³There is also a construction specific for the application to \mathcal{BPP} with parameters satisfying: $\ell = c \cdot \log(m)$ and $n = O(c^2 \cdot \log(m))$, where c is a constant, cf. [12, L.2.6]. That is not good in proof complexity as a trivial proof of the τ -formulas going through all possible seeds would have a polynomial size.

of the conjecture seems quite harmless and, moreover, very much in the spirit of [12] allowing to compute the output bits of the generator in time $m^{O(1)}$ rather than just $n^{O(1)}$.

Nevertheless, (3) is a modification of the original specifications and so we do not want to talk about Conjecture 4.1 in the next theorem. For this reason let us formulate the following **Statement R**:

R Let g be an NW-generator based on an $m \times n$ matrix A that is an $(\ell, \log(m))$ combinatorial design and on any function f such that the constraints in (1), (2) and (3) are satisfied. Then g is hard for EF .

Theorem 4.2 *Assume that Statement R is true. Then EF is not p -bounded.*

Proof :

Assume that EF is p -bounded. We shall arrive at a contradiction with Statement R. If EF is p -bounded then, in particular, $\mathcal{NP} = co\mathcal{NP}$.

By Theorem 3.1 (Part 2) there is a function f in $\mathcal{NE} \cap co\mathcal{NE}$ that has exponential hardness on average. Having such f we may apply Statement R (suitable matrices A are constructed in [12]) to conclude that EF is not p -bounded. That is a contradiction.

q.e.d.

Note that if we postulate smaller m then although the size of the formula might not be polynomial in m anymore it will still be superpolynomially smaller than $2^{O(n)}$, i.e. than a lower bound to the size of the trivial proof going through all seeds.

An optimist may conclude that it only takes to prove a conditional statement in order to prove that EF is not p -bounded. A pessimist may conclude that if the conclusion of R holds even without it's hypothesis then the hypothesis is irrelevant. I think an interesting modification of R may be obtained by stating it for one particular function f , e.g. the hard bit of the discrete logarithm, and maybe claiming the lower bound for $\tau_b(g)$ for a random b with high probability, rather than for all b 's.

Acknowledgements: I thank E. Jeřábek and P. Pudlák for discussions and suggestions. In particular, P. Pudlák suggested that some variant of Theorem 2.1 should be relevant to proof complexity generators.

References

- [1] M. ALEKHNovich, E. BEN-SASSON, A. A. RAZBOROV, and A. WIGDERSON, Pseudorandom generators in propositional proof complexity, *Electronic Colloquium on Computational Complexity*, Rep. No.**23**, (2000). Ext. abstract in: *Proc. of the 41st Annual Symp. on Foundation of Computer Science*, (2000), pp.43-53.
- [2] S. A. COOK and A. R. RECKHOW, The relative efficiency of propositional proof systems, *J. Symbolic Logic*,**44(1)**, (1979), pp.36-50.
- [3] H. FRIEDMAN, On the consistency, completeness, and correctness problems, (unpublished preprint, 1979).
- [4] R. IMPAGLIAZZO and A. WIGDERSON, $P = BPP$ unless E has sub-exponential circuits: derandomizing the XOR lemma, in: *Proc. of the 29th Annual ACM Symposium on Theory of Computing*, (1997), pp. 220-229.
- [5] J. KRAJÍČEK, *Bounded arithmetic, propositional logic, and complexity theory*, Encyclopedia of Mathematics and Its Applications, Vol. **60**, Cambridge University Press, (1995).
- [6] J. KRAJÍČEK, On the weak pigeonhole principle, *Fundamenta Mathematicae*, Vol.**170(1-3)**, (2001), pp.123-140.
- [7] J. KRAJÍČEK, Tautologies from pseudo-random generators, *Bulletin of Symbolic Logic*, **7(2)**, (2001), pp.197-212.
- [8] J. KRAJÍČEK, Dual weak pigeonhole principle, pseudo-surjective functions, and provability of circuit lower bounds, *Journal of Symbolic Logic*, to app. (preprint Nov.'02).
- [9] J. KRAJÍČEK, Hardness assumptions in the foundations of theoretical computer science, (preprint in ITI series: <http://iti.mff.cuni.cz/series/index.html>, Jan.'03).
- [10] J. KRAJÍČEK, Implicit proofs, *J. of Symbolic Logic*, to app., (preprint July 2003).
- [11] J. KRAJÍČEK, and P. PUDLÁK, Propositional proof systems, the consistency of first order theories and the complexity of computations, *J. Symbolic Logic*, **54(3)**, (1989), pp.1063-1079

- [12] N. NISAN, and A. WIGDERSON, Hardness vs. randomness, *J. Comput. System Sci.*, Vol.**49**, (1994), pp.149–167.
- [13] P. PUDLÁK, On the length of finitistic consistency statements in first order theories, in: *Logic Colloquim 84*, (1986), pp.165-196, North-Holland.
- [14] P. PUDLÁK, Improved bounds to the length of proofs of finitistic consistency statements, *Contemporary Mathematics*, **65**, (1987), pp. 309-331.
- [15] A. A. RAZBOROV, Unprovability of lower bounds on the circuit size in certain fragments of bounded arithmetic, *Izv. Ross. Akad. Nauk Ser. Mat.*, **59(1)**, (1995), pp. 201-224.
- [16] A. A. RAZBOROV, Resolution lower bounds for perfect matching principles, in: *Proc. of the 17th IEEE Conf. on Computational Complexity*, (2002), pp.29-38.
- [17] A. A. RAZBOROV, Pseudorandom generators hard for k -DNF resolution and polynomial calculus resolution, preprint, (May'03).
- [18] O. V. VERBITSKY, Optimal algorithms for co-NP sets and the problem $EXP \stackrel{?}{=} NEXP$. (Russian) *Mat. Zametki*, Vol.**50(2)**, (1991), pp.37-46.

Mailing address:

Mathematical Institute
 Academy of Sciences
 Žitná 25, Prague 1, CZ - 115 67
 The Czech Republic
 krajicek@math.cas.cz