



On k -term DNF with the largest number of prime implicants*

Robert H. Sloan

University of Illinois at Chicago, Chicago, IL 60607, USA,
sloan@uic.edu,

WWW home page: <http://www.cs.uic.edu/~sloan>

Balázs Szörényi

Hungarian Academy of Sciences and University of Szeged
Research Group on Artificial Intelligence, Szeged, Hungary,
szorenyi@rgai.hu

György Turán

University of Illinois at Chicago, Chicago, IL 60607, USA,
Hungarian Academy of Sciences and University of Szeged
Research Group on Artificial Intelligence, Szeged, Hungary,
gyt@uic.edu

February 8, 2005

Abstract

It is known that a k -term DNF can have at most $2^k - 1$ prime implicants and this bound is sharp. We determine all k -term DNF having the maximal number of prime implicants. It is shown that a DNF is maximal if and only if it corresponds to a non-repeating decision tree with literals assigned to the leaves in a certain way. We also mention some related results and open problems.

1. Introduction

Prime implicants of a Boolean function, or, in other words, maximal subcubes of a subset of the n -dimensional hypercube, form a basic concept for the theory of Boolean functions and their applications. Concerning the maximal number of prime implicants, it is known that an n -variable Boolean function can have at most $O(\frac{3^n}{\sqrt{n}})$ prime implicants, and there are n -variable Boolean functions with $\Omega(\frac{3^n}{n})$ prime implicants (see, e.g., [4]).

*This material is based upon work supported by the National Science Foundation under Grant Nos. CCR-0100036 and CCF-0431059.




Figure 1: A non-repeating, unate-leaf decision tree (NUD)

Another case considered is the maximal number of prime implicants of Boolean functions represented by disjunctive normal forms (DNF) with a bounded number of terms. The result that a k -term DNF can have at most $2^k - 1$ prime implicants was discovered independently by Chandra and Markowsky [4], Levin [15] and McMullen and Shearer [17]. For a recent application in computational learning theory, see Hellerstein and Raghavan [9]. It was shown by Laborde [14], Levin [15] and McMullen and Shearer [17] that the bound is sharp, i.e., there are k -term DNF with $2^k - 1$ prime implicants (Chandra and Markowsky gave an example with more than $2^{k/2}$ prime implicants). In view of these results, we call a DNF *maximal* if it has k terms and $2^k - 1$ prime implicants for some k .

In this paper we complete the results of [4, 14, 15, 17] by determining all the maximal disjunctive normal forms. In order to formulate the description, let us introduce the following definition.

By a tree we mean a rooted binary tree such that for every inner node, the edge leading to its left (resp., right) child is labeled 0 (resp., 1). For a given $k \geq 2$ and $r \geq 0$, let us consider the variables x_1, \dots, x_{k-1} , and the literals y_1, \dots, y_k and z_1, \dots, z_r (all variables in the list are assumed to be different). A *non-repeating, unate-leaf decision tree (NUD)* T over these variables and literals is constructed by taking a tree with $k - 1$ inner nodes (and thus with k leaves), assigning to each inner node a distinct variable x_i , assigning to each leaf a distinct literal y_j , and, in addition, assigning to each leaf an arbitrary subset of the z literals. The set of leaves of T is denoted by L . If we want to mention the number of x variables and y literals used in the construction, then we refer to T as a k -NUD (the value r is irrelevant). Figure 1 gives an example of a 5-NUD (the labelling of the edges is omitted for simplicity).

A k -NUD represents a k -term DNF, determined as follows. For a leaf $\ell \in L$, let the term t_ℓ be the conjunction of the x literals along the path leading to ℓ (where traversing an edge labeled 1 corresponds to an unnegated literal, and traversing an edge labeled 0 corresponds to a negated literal) and of the y and z literals assigned to ℓ . The k -term DNF represented by the k -NUD T is

$$\varphi_T = \bigvee_{\ell \in L} t_\ell.$$

For example, the 5-term DNF represented by the 5-NUD of Figure 1 is

$$\overline{x_1} \overline{x_2} \overline{x_4} y_1 z_1 \vee \overline{x_1} \overline{x_2} x_4 y_2 z_2 z_3 \vee \overline{x_1} x_2 y_3 z_1 \vee x_1 \overline{x_3} y_4 z_1 z_4 \vee x_1 x_3 y_5 z_2.$$

The Boolean function represented by φ_T can also be thought of in the following way: given a truth assignment a to all the variables, use the values of the x variables to determine a path from the root to a leaf. The function value is 1 if a makes all the y and z literals assigned to this leaf true, and it is 0 otherwise. It is clear from the definition that the input vectors accepted at a leaf ℓ are precisely those vectors which satisfy the term t_ℓ . The function φ_T is a generalized addressing function or multiplexer [18, 23]. If a DNF φ comes from a NUD T , then T can be reconstructed from φ . The y and z literals are those which are unate in φ , i.e., their negation does not occur in φ , while the x variables are those which occur both negated and unnegated. Among the x variables, the one labeling the root is the only one which occurs in every term (either unnegated or negated). The left child is the only x variable which occurs in every term containing the negation of the root variable, etc. In view of this correspondence, with some abuse of terminology, we can talk about a DNF being a NUD, rather than being equivalent to a NUD. The maximal DNF of [14, 17] (resp., [15]) corresponds to a tree which is a single path (resp., a complete binary tree), without any z literals. A NUD generalizes these examples by allowing for an arbitrary tree and for the additional z literals. Now we can formulate the description of maximal DNF.

Theorem 1. *A DNF is maximal if and only if it is a NUD.*

A closely related class of DNF *tautologies* is obtained if we consider trees with the same kind of inner nodes, but without any literals assigned to the leaves. In the case of the example of Figure 1, the corresponding DNF tautology is

$$\overline{x_1} \overline{x_2} \overline{x_4} \vee \overline{x_1} \overline{x_2} x_4 \vee \overline{x_1} x_2 \vee x_1 \overline{x_3} \vee x_1 x_3.$$

Let us refer to this class of tautologies as *nonrepeating decision tree* tautologies, or *ND*'s. The main step in the proof of Theorem 1 is to show that for every DNF tautology the following two properties are equivalent: *a*) any two of its terms have exactly one conflicting pair of literals (in other words, the terms are pairwise neighboring), *b*) it is an ND. This result was proven recently, independently from our work, by Kullmann [12, 13]. Kullmann's proof uses the concept of Hermitian defect and other concepts from linear algebra. It also uses the characterization of ND's as strongly minimal tautologies with the additional property that the number of terms is one more than the number of variables (Aharoni and Linial [1], Davydov *et al.* [5] Kullmann [11]), proved using Hall's theorem or resolution techniques. (A tautology is strongly minimal if deleting any term, or adding any literal to a term results in a non-tautology.) Our proof is an elementary combinatorial argument.

We note that ND's come up in other contexts as well, e.g., in connection with the complexity of analytic tableaux (Urquhart [22], referring to earlier unpublished work of Cook, and Arai *et al.* [2]). Another related topic is the decision tree complexity of tautologies (Lovász *et al.* [16]), which is discussed further in [21].

The characterization of ND's as pairwise neighboring DNF tautologies is a direct consequence of the following splitting lemma: if the n -dimensional hypercube is partitioned into subcubes of pairwise

distance one, then there is a split of the whole cube into two half cubes such that every cube of the partition is contained in one of the two halves. We also consider the question of what can be said about cube partitions without the distance assumption. The goodness of a split into two half cubes can be measured by the fraction of the total volume of subcubes contained in one of the two halves (thus in the distance 1 case one always has a split of measure 1). It corresponds to a notion of influence of the variable determining the split on the partition (for other notions of influence, see, e.g., Hammer *et al.* [8] and Kahn *et al.* [10]). We give general lower and upper bounds for the best achievable split. The upper bound uses a result of Savicky and Sgall [19] on DNF tautologies with bounded occurrences of the variables.

Recent related work on the combinatorial aspects of the satisfiability problem (see Kullmann [13] for a recent survey) makes use of the connection with partitioning complete graphs into complete bipartite graphs (bicliques). This connection, and in particular, the Graham - Pollak theorem [7] is used by Laborde [14] to show that a maximal k -term DNF contains at least $2k - 1$ variables. (This result, in turn, follows immediately from Theorem 1 above without using the Graham - Pollak theorem.)

The paper is organized as follows. After some preliminaries in Section 2, the results of [4, 14, 15, 17] are presented in Section 3. The proof of Theorem 1 is given in Section 4. The Splitting Lemma is proved in Section 5. Section 6 contains the bounds for the general splitting problem. The connection to the Graham - Pollak theorem is discussed briefly in Section 7. Section 8 contains some further open problems on the number of prime implicants.

2. Preliminaries

A literal is a variable or a negated variable, a term is a conjunction (or a set) of literals, and a disjunctive normal form (DNF) is a disjunction of terms. The empty conjunction (resp. disjunction) is identically true (resp. false). It is assumed that terms do not contain both a variable and its negation. The size of a term t , denoted by $|t|$, is the number of its literals. The number of *conflicts* between two terms is the number of variables occurring unnegated in one term and negated in the other. A DNF is *disjoint* if any two of its terms have at least one conflict. We write $\psi \leq \varphi$ if every truth assignment satisfying ψ also satisfies φ , and $\psi < \varphi$ if, in addition, there is a truth assignment a with $\psi(a) = 0$ and $\varphi(a) = 1$. The set of vectors in $\{0, 1\}^n$ satisfying φ are denoted by $T(\varphi)$. If t is a term then $T(t)$ is a subcube (or simply cube) in $\{0, 1\}^n$, with $|T(t)| = 2^{n-|t|}$. With an abuse of notation, we usually write cube t instead of cube $T(t)$. For a literal z , the z half cube of $\{0, 1\}^n$ is the $(n - 1)$ -dimensional subcube formed by the vectors for which z is true.

A term t is an *implicant* of a DNF $\varphi = t_1 \vee \dots \vee t_k$ if $t \leq \varphi$. In this case we also say that φ is a *cover* of t , as the union of the cubes $T(t_i)$ covers the cube $T(t)$. Note that the variables occurring in t and φ may differ. It may be assumed *w.l.o.g.* that by a truth assignment we mean an assignment

of truth values to every variable occurring in t or φ . The term t is a *prime implicant* of φ , if t is an implicant of φ , but every term obtained by deleting a literal from t is not an implicant of φ . The DNF φ is a *minimal cover* of the term t , if φ is a cover of t (i.e., t is an implicant of φ), but every DNF obtained from φ by deleting a term is not a cover of t .

Let t be a term, and $\varphi = t_1 \vee \dots \vee t_k$ be a DNF. Every term t_i of φ can be uniquely written in the form

$$t_i = t'_i \wedge t''_i, \quad (1)$$

where t'_i contains all the literals from t_i which also occur in t , and t''_i contains the remaining literals of t_i .

Given a DNF φ , let $Var(\varphi)$ (resp., $Lit(\varphi)$) denote the set of variables (resp., literals) occurring in any term of φ , and let

$$UL(\varphi) = \{z \in Lit(\varphi) : \bar{z} \notin Lit(\varphi)\} \quad (2)$$

be the set of *unate* literals in φ , i.e. the set of those literals occurring in φ , for which their negation does not occur in φ .

For $a \in \{0, 1\}^n$, the vector $a^{(z)}$ is the vector obtained from a by flipping its component corresponding to the literal z . Given $x, y \in \{0, 1\}^n$, the smallest subcube containing both x and y is denoted by $Cube(x, y)$. It is obtained by including every literal corresponding to components where x and y agree. The Hamming distance $d(x, y)$ of $x, y \in \{0, 1\}^n$ is the number of components where x and y differ. The graph of the n -dimensional cube has $\{0, 1\}^n$ as vertices, and edges (x, y) for every x, y of Hamming distance 1. The distance of two subcubes C_1 and C_2 is $\min\{d(x, y) : x \in C_1, y \in C_2\}$. Note that the distance of $T(t_1)$ and $T(t_2)$ is equal to the number of conflicts between the terms t_1 and t_2 . A partition of the cube into subcubes can also be viewed as a disjoint DNF tautology. A partition of a cube into subcubes is *pairwise neighboring*, if any two subcubes in the partition have distance 1. A set of terms forms a pairwise neighboring partition, if the corresponding set of cubes forms a pairwise neighboring partition.

3. Prime implicants and k -term DNF

In this section we describe the results of [4, 14, 15, 17] on prime implicants of k -term DNF. We give a complete presentation in order to make the paper self-contained, to clarify what are the consequences of the separate assumptions of being an implicant, a prime implicant, resp. a minimal cover, and to give an explicit formulation of results implicit in [14]. We use the notation introduced above in (1) and (2).

Proposition 2. *A term t is an implicant of a DNF φ if and only if $\bigvee_{i=1}^k t''_i = 1$.*

Proof For the \Leftarrow direction, let a be a truth assignment such that $t(a) = 1$. Then $t'_i(a) = 1$ for every i and $t''_i(a) = 1$ for some i , so $t_i(a) = 1$ for some i , and thus $\varphi(a) = 1$.

For the \Rightarrow direction assume $\bigvee_{i=1}^k t_i'' < 1$, i.e., $\left(\bigvee_{i=1}^k t_i''\right)(a) = 0$ for some a . The literals occurring in $\bigvee_{i=1}^k t_i''$ do not occur in t , but it may be the case that the negation of such a literal occurs in t . Let b be the truth assignment obtained from a by setting all the literals of t to 1. Then every literal in $\bigvee_{i=1}^k t_i''$ is either unchanged, or is changed to 0, thus $\left(\bigvee_{i=1}^k t_i''\right)(b) = 0$, and so $\varphi(b) = 0$. But $t(b) = 1$, contradicting the fact that t is an implicant of φ . \square

Proposition 3. *If t is a prime implicant of φ then*

- a) $t = \bigwedge_{i=1}^k t_i'$,
- b) $Lit(t) \subseteq Lit(\varphi)$.

Proof For a), it follows from the definition that $t \leq \bigwedge_{i=1}^k t_i'$. Assume that a variable x in t does not occur in any t_i . Then x does not occur in φ at all, though \bar{x} may occur in some t_i'' . But then t is an implicant of the disjunction of those terms in φ which do not contain \bar{x} , and so by deleting x from t we still get an implicant of φ . Part b) follows trivially from a). \square

Proposition 4. *If φ is a minimal cover of t then*

- a) $Lit(t) \cap Lit(\varphi) = UL(\varphi)$,
- b) $\bigvee_{i=1}^k t_i''$ is a minimal cover of 1.

Proof For the \subseteq part of a) note that if t contains a non-unate literal z of φ , then terms containing \bar{z} can be deleted from φ and we still get a cover of t , contradicting the minimality of φ . For the \supseteq part of a), assume that a unate literal z is not contained in t . Then $\bar{z}t$ is also an implicant of φ , which is covered by the terms of φ not containing z . As these terms do not contain \bar{z} either, their disjunction covers t as well, again contradicting the minimality of φ . Part b) follows from Proposition 2. \square

Putting together Propositions 2, 3 and 4, we get the following.

Theorem 5. *If t is a prime implicant of φ and φ is a minimal cover of t then*

- a) $Lit(t) = UL(\varphi)$,
- b) $\bigvee_{i=1}^k t_i''$ is a minimal cover of 1. \square

Theorem 6. ([4, 15, 17])

Every k -term DNF has at most $2^k - 1$ prime implicants.

Proof Let φ be a k -term DNF and t be a prime implicant of φ . Consider a minimal set of terms of φ covering t . Then, by Theorem 5 a), t is uniquely determined this set of terms. \square

The next result gives important structural information on maximal DNF's.

Theorem 7. ([14])

Let $\varphi = t_1 \vee \dots \vee t_k$ be a k -term DNF with $2^k - 1$ prime implicants. Then

a) $\bigvee_{i=1}^k t_i''$ is a minimal cover of 1,

b) t_i'' and t_j'' conflict in exactly one variable, for every $1 \leq i < j \leq k$.

Proof By Theorems 5 and 6, every nonempty subset of the terms of φ is a minimal covering of some prime implicant of φ . Part a) follows by applying Theorem 5 b) to all the terms.

Let us consider now $\psi_{i,j} = t_i \vee t_j$. Again, this is a minimal cover of a prime implicant of φ . If t_i and t_j do not conflict in any variable, then, by Theorem 5 a), the corresponding prime implicant is the term formed by all the literals in t_i and t_j . But that term is not a prime implicant. Indeed, it must be the case that $t_i \neq t_j$, and so $t_i \wedge t_j < t_i$ or $t_i \wedge t_j < t_j$. If t_i and t_j conflict in more than one variable, then we get a contradiction to Theorem 5 b), as a the disjunction of two terms with at least two conflicts cannot be 1. \square

4. Proof of Theorem 1

In this section we prove Theorem 1. First we consider the \Leftarrow direction.

Lemma 8. *Every NUD is maximal.*

Proof Let T be a k -NUD, and let H be a nonempty subset of its leaves. Define the term

$$t_H = UL(\{t_\ell : \ell \in H\}).$$

Let a be a truth assignment satisfying t_H . It follows by induction of the number of inner nodes evaluated, that on input a we arrive to a leaf belonging to H , and it follows from the definition of t_H that a satisfies every literal assigned to that leaf. Thus t_H is an implicant of φ_T .

Assume that we delete an x literal, say x_i^ϵ from t_H , to get the term t' . As $x_i^\epsilon \in UL(\{t_\ell : \ell \in H\})$, there is a leaf ℓ_1 belonging to H below the ϵ -child of the inner node x_i , but no leaf below the $(1 - \epsilon)$ -child of x_i is in H . Let a be the vector satisfying all the literals in t_{ℓ_1} and t_H , with every variable not occurring in these terms set to 0. Let $b = a^{(x_i)}$. On the input b we arrive to a leaf ℓ_2 below the $(1 - \epsilon)$ -child of x_i . But the y literal assigned to ℓ_2 is set to 0 in b , and hence $\varphi_T(b) = 0$. On the other hand, b still satisfies t' . Thus t' is not an implicant.

Assume now that we delete a y literal, say y_j , from t_H , to get the term t' . Let ℓ be the leaf containing y_j . It follows from the definition of t_H that $\ell \in H$. Let a be a vector satisfying t_ℓ and t_H , and let $b = a^{(y_j)}$. Then the input b leads to ℓ , but as its y_j component is 0, we get $\varphi_T(b) = 0$. On the other hand, b still satisfies t' . Thus t' is not an implicant. The case when we delete a z literal, say z_j , from t_H is the same, except now there may be several leaves in H containing z_j . We

can choose any such leaf, and repeat the same argument as for y_j . It again follows that the term obtained after deleting the literal is not an implicant.

Thus the term t_H is a prime implicant of φ_T . Terms corresponding to different subsets of L are different, as each leaf has its unique y literal. Hence φ_T has at least $2^k - 1$ prime implicants, and so it is maximal by Theorem 6. \square

The rest of this section contains the proof of the converse.

Lemma 9. *Every maximal DNF is a NUD.*

Proof Let $\varphi = t_1 \vee \dots \vee t_k$ be a k -term DNF with $2^k - 1$ prime implicants. Consider the term $t = UL(\varphi)$, and the decomposition $t_i = t'_i \wedge t''_i$ of the terms of φ w.r.to t , as in (1). According to Theorem 7, the terms t''_1, \dots, t''_k form a pairwise neighboring partition over the non-unate variables occurring in φ , i.e., over $\{0, 1\}^s$, where $s = |Var(\varphi) \setminus UL(\varphi)|$.

The proof of the following lemma is given in Section 5.

Lemma 10. (*Splitting Lemma*) *If a set of $k \geq 2$ terms form a pairwise neighboring partition, then there is a variable that occurs (unnegated or negated) in every term.*

This lemma implies the characterization of nonrepeating decision tree tautologies mentioned in the introduction.

Lemma 11. (*ND Lemma*) [12] *A set of $k \geq 2$ terms form a pairwise neighboring partition if and only if it is an ND.*

Proof Apply Lemma 10 to the pairwise neighboring partition to get a variable x_1 occurring in every term. It must be the case that x_1 occurs both unnegated and negated, as otherwise the cubes would not cover the whole cube. If the x_1^c half cube contains just one cube then we stop at that branch, otherwise we use the lemma again to get a variable which occurs in every subcube of the partition, belonging to the x_1^c half cube, etc. In this way we get a tree, where the inner nodes are labeled with variables and there are k leaves ℓ_1, \dots, ℓ_k corresponding to the cubes in the partition. (The tree constructed is (the dual of) a special *search tree* in the sense of [16] for the partition.) The labels of the inner nodes are *different*, as the same label appearing twice would mean that some pair of cubes have distance at least 2. Indeed, if variable x_i occurs twice then let x_j be the variable labeling the least common ancestor of the two occurrences in the tree. By construction, there are terms containing $\bar{x}_i \bar{x}_j$, resp. $x_i x_j$. Thus the partition is an ND. \square

Now we can complete the proof of Lemma 9. Lemma 11 gives a nonrepeating decision tree for the pairwise neighboring terms t''_1, \dots, t''_k . We claim that by adding the literals in t'_i to the leaf ℓ_i , we get a k -NUD for φ . Consider any truth assignment a to the variables in φ . Evaluating the tree on a , we arrive to a leaf corresponding to a term t''_i . As $\varphi(a) = 1$ iff $t'_i(a) = 1$, the tree computes

φ correctly. By construction, all the literals in the leaves are unate. Thus, in order to verify the NUD-ity of the tree, it only remains to show that for every leaf there is a literal which occurs only in that leaf (that literal will be its y literal). Assume that this is not the case, and every (unate) literal assigned to leaf ℓ_i occurs in some other leaf. Let x_j^ϵ be the last literal on the path leading to ℓ_i . Then $x_j^{1-\epsilon} \in UL(\varphi \setminus t_i)$. We claim that $UL(\varphi \setminus t_i) \setminus \{x_j^{1-\epsilon}\}$ is an implicant of φ . Let a be a truth assignment satisfying every literal in $UL(\varphi \setminus t_i) \setminus \{x_j^{1-\epsilon}\}$, and let us evaluate the tree on a . If we arrive to a leaf other than ℓ_i , then $\varphi(a) = 1$ by construction. But $\varphi(a) = 1$ if we arrive to ℓ_i as well, as all unate literals in ℓ_i occur in other leaves, and thus they must be set to 1 in a . Thus $UL(\varphi \setminus t_i)$ is not a prime implicant of φ , contradicting Theorems 5 and 6. \square

5. Proof of the Splitting Lemma (Lemma 10)

Let u_1, \dots, u_k be terms forming a pairwise neighboring partition of $\{0, 1\}^s$. For a literal z consider the union of cubes $T(u_i)$ contained in the z half cube, i.e., put

$$S_z = \bigcup_{\{i: z \in u_i\}} T(u_i).$$

We show that S_z is always a cube, and that the largest S_z is the entire z half cube. As then $S_{\bar{z}}$ is the entire \bar{z} half cube, this implies the lemma.

Note that if neither z nor \bar{z} occur in a term u , then for every vector a it holds that $a \in T(u)$ iff $a^{(z)} \in T(u)$. If a vector a in the z half cube is not in S_z , then it is covered by a cube not containing z or \bar{z} , and so $a^{(z)}$ is covered by the same cube. Thus $a^{(z)} \notin S_{\bar{z}}$. Therefore, for every vector a in the z half cube it holds that

$$a \in S_z \Leftrightarrow a^{(z)} \in S_{\bar{z}}. \quad (3)$$

Lemma 12. *For every literal z it holds that S_z is a cube.*

Proof Suppose that S_z is not a cube. We show below that there is a path (a, b, c) in the graph of the z -half cube such that $a, c \in S_z$ and $b \notin S_z$. Then $b = a^{(x)}, c = b^{(y)}$, for some variables $x \neq y$. Consider the cubes $T(u)$ (resp., $T(u')$) containing a (resp., c). These cubes must be different, as otherwise b would be in the same cube, and thus in S_z as well. By the definition of S_z , both u and u' contain z . We know that u and u' have a conflict. As $a \in T(u)$ and $a^{(x,y)} \in T(u')$, the conflicting variable must be x or y . Assume *w.l.o.g.* that $x \in u$ and $\bar{x} \in u'$. Using (3) we get $a^{(z)} \in S_{\bar{z}}$ and $b^{(z)} \notin S_{\bar{z}}$. Let $a^{(z)}$ be covered by the cube $T(u'')$. By the definition of $S_{\bar{z}}$, the term u'' contains \bar{z} , and furthermore, $b^{(z)} \notin T(u'')$. As $a^{(z)}$ and $b^{(z)}$ only differ in their x component, it must be the case that $x \in u''$. Thus $\bar{z}, x \in u''$ and $z, \bar{x} \in u'$, so u'' and u' conflict in at least two variables, a contradiction.

We still need to show that, as claimed above, if S_z is not a cube then there is a path (a, b, c) in the z -half cube such that $a, c \in S_z$ and $b \notin S_z$. First we note that if a set is not a cube then this fact

can be certified by three points (see, e.g., [9] for a precise definition of a certificate and applications of this notion, and [20] for related results).

Proposition 13. *If a set A is not a cube, then there are $a_0, c_0 \in A$ and $b_0 \notin A$ such that $b_0 \in \text{Cube}(a_0, c_0)$.*

Proof Let \wedge (resp. \vee) of a set of vectors denote their componentwise \wedge (resp. \vee). For any two vectors a_0, c_0 it holds that $a_0 \wedge c_0 \in \text{Cube}(a_0, c_0)$ and $a_0 \vee c_0 \in \text{Cube}(a_0, c_0)$. Thus if the proposition is false then $A = \text{Cube}(\bigwedge_{a \in A} a, \bigvee_{a \in A} a)$, a contradiction. \square

Therefore, if S_z is not a cube then there are three vectors a_0, b_0, c_0 in the z half cube such that $a_0, c_0 \in S_z, b_0 \notin S_z$ and $b_0 \in \text{Cube}(a_0, c_0)$. Let $T(u)$ (resp., $T(u')$) be the cube containing a_0 (resp., c_0). The two cubes must be different, as otherwise b_0 would be in the same cube, and thus also in S_z . The terms u and u' have exactly one conflict. We now observe that there is a shortest path between a_0 and c_0 , which is contained in the union of the two cubes. This is a special case of a more general result of Ekin *et al.* [6].

Proposition 14. [6] *Let u and u' be terms conflicting in exactly one variable. If $a_0 \in T(u)$ and $c_0 \in T(u')$, then $T(u \vee u')$ contains a shortest path connecting a_0 and c_0 .*

Proof Write *w.l.o.g.* $a_0 = 0a^1a^2a^3a^4$ and $c_0 = 1c^1c^2c^3c^4$, where the first component corresponds to the variable where the two terms conflict, the second subvector corresponds to literals common in the two terms, the third subvector to literals only occurring in u , the fourth to literals only occurring in u' , and the fifth to literals that do not occur in either of the two terms. Then a required shortest path can be built by completing the sequence $a_0, 0a^1a^2c^3c^4 \in T(u), 1c^1a^2c^3c^4 \in T(u')$ and c_0 , noting that $a^1 = c^1$. \square

Thus so far we know that $a_0, c_0 \in S_z, b_0 \notin S_z$ and $b_0 \in \text{Cube}(a_0, c_0)$ and there is a shortest path in $T(u \vee u')$ connecting a_0 and c_0 . The shortest path, therefore, is in $S_z \cap \text{Cube}(a_0, c_0)$ (it is in S_z as $T(u \vee u') \subseteq S_z$, and it is in $\text{Cube}(a_0, c_0)$ as any shortest path between a_0 and c_0 is in this cube). Based on this information, we would like to find a path (a, b, c) such that $a, c \in S_z$ and $b \notin S_z$.

Given a path (p, q, r) in a cube, there is a unique vertex $q' \neq q$ such that (p, q', r) is also a path. For example, if $p = 000, q = 100$ and $r = 110$ then $q' = 010$. A subset of the cube is *closed under switches* if for every path (p, q, r) in the set, the vertex q' also belongs to the set.

Proposition 15. *Assume that a set $B \subseteq \{0, 1\}^v$ contains a shortest path between two opposite vertices of $\{0, 1\}^v$, and is closed under switches. Then B is the whole cube.*

Proof The claim is trivial for $v = 2$. For $v \geq 3$, assume *w.l.o.g.* that the two opposite vertices are 0^v and 1^v , and that the shortest path is $1^i 0^{v-i}$ ($0 \leq i \leq v$). It follows by induction that $(a, 0) \in B$ for every $a \in \{0, 1\}^{v-1}$. Building a similar chain from the i 'th unit vector to 1^v , it again follows by

induction that every vector having 1 at the i 'th position ($1 \leq i \leq v-1$) is in B . Finally, $0^{v-1}1 \in B$ follows from $0^{v-3}101, 0^{v-3}111, 0^{v-3}011 \in B$. \square

Now let us apply this proposition to the set $S_z \cap \text{Cube}(a_0, c_0)$ in the cube $\text{Cube}(a_0, c_0)$. We know that it contains a shortest path between two opposite vertices, and it is not the whole cube. By the proposition, $S_z \cap \text{Cube}(a_0, c_0)$ is not closed under switches. Thus there is a path (a, b, c) in $S_z \cap \text{Cube}(a_0, c_0)$ such that $a, c \in S_z$ and $b \notin S_z$. Hence (a, b, c) is a path with the required properties. This completes the proof of Lemma 12. \square

Now we return to the proof of the Splitting Lemma. Consider a literal z such that $|S_z|$ is as large as possible. We show by induction that S_z is the entire z half cube, which, as noted above, implies the lemma. The statement is trivial for $n = 1, 2$. For $n > 2$, if $|S_z| = 2^{n-1}$ then we are done. Otherwise Lemma 12 implies that

$$|S_z| \leq 2^{n-2}. \quad (4)$$

Apply the induction hypothesis to the z -half cube. The restriction of the terms u_1, \dots, u_k to this cube is again a pairwise neighboring partition. Assume that every term of the restricted partition contains y or \bar{y} . Assume *w.l.o.g.* that $|S_y \cap S_z| \leq |S_{\bar{y}} \cap S_z|$ (where S_y refers to the original partition). By definition, S_y contains all the points in $S_y \cap S_z$. Also, by (3), for every $a \notin S_z$ in the quarter cube ($y = 1, z = 1$), S_y contains both a and $a^{(z)}$. Thus, using (4) one gets

$$|S_y| \geq |S_y \cap S_z| + 2(2^{n-2} - |S_y \cap S_z|) = 2^{n-1} - |S_y \cap S_z| \geq 2^{n-1} - 2^{n-3} > 2^{n-2},$$

which contradicts the choice of z . \square

6. The general splitting problem for cube partitions

According to the Splitting Lemma (Lemma 10), for every pairwise neighboring cube partition, the whole cube can be split into two halves in such a way that every cube of the partition is contained in one of the halves. In this section we consider the following question: what can be said without the pairwise neighboring property? Given an arbitrary cube partition of the whole cube and a split into two halves, let us say that a cube in the partition is good, if it is contained in either one of the halves. We would like to find a split such that the good cubes contain many points.

Thus we consider the following quantities. Given a cube partition φ over the variables x_1, \dots, x_n and a variable x_j , let

$$v_{\varphi, j} = \sum \left\{ 2^{-|t|} : t \in \varphi, x_j \in t \text{ or } \bar{x}_j \in t \right\}$$

be the fraction of the volume of good cubes in φ *w.r.to* the x_j split of the cube, and let

$$\alpha_n = \min_{\varphi} \max_{1 \leq j \leq n} v_{\varphi, j},$$

where φ ranges over all cube partitions, or in other words, over all disjoint DNF tautologies. Note that as φ is a partition it holds that

$$\sum_{t \in \varphi} 2^{-|t|} = 1. \quad (5)$$

Theorem 16.

$$\frac{\log n - \log \log n}{n} \leq \alpha_n \leq O\left(n^{-\frac{1}{5}}\right).$$

Proof Let $\varphi = t_1 \vee \dots \vee t_r$ be a disjoint DNF tautology over the variables x_1, \dots, x_n . If the term t_i contains x_j or \bar{x}_j , then t_i contributes $2^{-|t_i|}$ to $v_{\varphi,j}$. Thus

$$\sum_{j=1}^n v_{\varphi,j} = \sum_{i=1}^r |t_i| \cdot 2^{-|t_i|},$$

and there is a variable x_j with

$$v_{\varphi,j} \geq \frac{1}{n} \sum_{i=1}^r |t_i| \cdot 2^{-|t_i|}.$$

Let s denote the size of the shortest term in φ . As every term has size at least s , it follows from (5) that

$$\frac{1}{n} \sum_{i=1}^r |t_i| \cdot 2^{-|t_i|} \geq \frac{s}{n} \sum_{i=1}^r 2^{-|t_i|} = \frac{s}{n}.$$

On the other hand, for every variable x_j occurring in a shortest term t_i it holds that $v_{\varphi,j} \geq 2^{-s}$. Thus

$$\alpha_n \geq \min\left(\frac{s}{n}, 2^{-s}\right),$$

and the lower bound follows by taking $s = \log n - \log \log n$.

The upper bound follows from a construction of Savicky and Sgall [19]. They constructed a disjoint DNF tautology over $n = 4^\ell$ variables, having 2^{3^ℓ} terms of size 3^ℓ , such that every variable occurs in at most a

$$\left(\frac{3}{4}\right)^\ell$$

fraction of the terms. The bound then follows by a direct calculation. \square

In view of Theorems 1 and 16 it may be of interest to consider the quantity α_n^d , which is defined as α_n , except that φ is restricted to cube partitions with pairwise distances bounded by d . In the construction of [19] the maximal distance grows linearly with n .

7. Partitions of complete graphs into complete bipartite graphs

Given a set of pairwise disjoint cubes in $\{0, 1\}^n$, corresponding to terms t_1, \dots, t_r , one can construct a covering

$$\mathcal{G} = \{G_1, \dots, G_n\}$$

of the r -vertex complete graph K_r by complete bipartite graphs, where G_u has an edge connecting vertices v_i and v_j if terms t_i and t_j conflict in the variable x_u . If the set of cubes is pairwise neighboring, then this covering is a partition, as the complete bipartite graphs are edge disjoint.

Conversely, given a covering $\mathcal{G} = \{G_1, \dots, G_n\}$ of K_r by complete bipartite graphs, we can construct a set of pairwise disjoint cubes t_1, \dots, t_r of $\{0, 1\}^n$. For every G_u fix arbitrarily one of the sides as the left side. The term t_i contains x_u (resp. \bar{x}_u), if vertex v_i is contained in the left (resp. right) side of G_u . If \mathcal{G} is a partition, then it follows that the t_i 's are pairwise neighboring. The cubes thus constructed do not necessarily form a partition of $\{0, 1\}^n$.

The Graham - Pollak theorem [7] states that every partition of K_r into complete bipartite graphs consists of at least $r - 1$ graphs. A large class of such partitions, which can be called *recursive* partitions, is obtained as follows: take a complete bipartite graph on the whole vertex set, and recursively add similar partitions of the complete graphs formed by the two sides of this bipartite graph (see, e.g., [3]).

Consider a partition $\mathcal{G} = \{G_1, \dots, G_n\}$ of K_r into complete bipartite graphs. Let the degree of a vertex v w.r.to \mathcal{G} , denoted by $d_{\mathcal{G}}(v)$, be the number of G_i 's containing v , and let the *volume* $vol(\mathcal{G})$ of the partition be defined as

$$vol(\mathcal{G}) = \sum_v 2^{-d_{\mathcal{G}}(v)}.$$

In view of the translation into a set of pairwise disjoint cubes in $\{0, 1\}^n$ described above, $vol(\mathcal{G}) \leq 1$ for every \mathcal{G} , as $d_{\mathcal{G}}(v_i) = |t_i|$ for every $i = 1, \dots, r$, and $vol(\mathcal{G}) = 1$ if and only if the cubes form a partition of $\{0, 1\}^n$. For example, the partition of K_4 into the 3 complete bipartite graphs $(\{1\}, \{3, 4\})$, $(\{2\}, \{1, 4\})$, and $(\{3\}, \{2, 4\})$ (mentioned in [14]) has volume $\frac{7}{8}$. This partition of K_4 is not recursive. (It was actually this example which suggested Lemma 10.) As a corollary to the Splitting Lemma (Lemma 10) one gets the following characterization of recursive partitions.

Corollary 17. *A partition \mathcal{G} is recursive if and only if $vol(\mathcal{G}) = 1$.*

Proof The \Rightarrow direction follows directly by induction on the number of vertices by considering the bipartite graph from \mathcal{G} which contains all the vertices.

For the \Leftarrow direction, one only has to note that the set of terms t_1, \dots, t_r constructed above is pairwise neighboring, and by the volume condition it is also a partition of the whole cube.

Applying Lemma 10 we get that there is a variable which occurs (unnegated or negated) in every term. This means that the corresponding bipartite graph contains all the r vertices. The remaining

partitions of the two sides of this bipartite graph have total volume 2, and thus each side must have volume 1. The statement then follows by induction. \square

The corollary shows that among partitions of K_r into complete bipartite graphs, recursive ones have the largest possible volume. Among the partitions of K_r into $r - 1$ complete bipartite graphs, which ones have *minimal* volume?

8. Other open problems

The k -term DNF

$$x_1\bar{x}_2 \vee x_2\bar{x}_3 \vee \dots \vee x_{k-1}\bar{x}_k \vee x_k\bar{x}_1,$$

which is false for 0^k and 1^k , and true everywhere else, has $k(k - 1)$ prime implicants, namely $x_i\bar{x}_j$ for every $i \neq j$. These prime implicants are all *shortest* prime implicants. How many shortest prime implicants can a k -term DNF have in general?

Another question concerns the maximal number of prime implicants of a Boolean function which is true at a given number of points. As noted by Levin [15], every implicant is determined by the top and bottom of the corresponding subcube (which may also be identical). Thus if a function is true at m points, then it has $O(m^2)$ prime implicants. It is also noted in [15] that the n -variable function which is true for vectors of weight between $\frac{n}{3}$ and $\frac{2n}{3}$, has $m^{\log 3 - o(1)}$ prime implicants. (This is the function with the largest known number of prime implicants among n -variable functions.) Thus the maximal number of prime implicants is polynomial in m , and the question is to get sharper bounds for the exponent.

References

- [1] R. Aharoni, N. Linial: Minimal non-two-colorable hypergraphs and minimal unsatisfiable formulas, *J. Comb. Th. A* **43** (1986), 196–204.
- [2] N. H. Arai, T. Pitassi, A. Urquhart: The complexity of analytic tableaux, *33. STOC* (2001), 356–363.
- [3] L. Babai, P. Frankl: *Linear Algebra Methods in Combinatorics*. Preliminary version 2, Univ. of Chicago, 1992.
- [4] A. K. Chandra, G. Markowsky: On the number of prime implicants, *Discrete Math.* **24** (1978), 7–11.
- [5] G. Davydov, I. Davydova, H. Kleine Buning: An efficient algorithm for the minimal unsatisfiability problem for a subclass of CNF, *Ann. Math. and Artif. Intell.* **23** (1998), 229–245.

- [6] O. Ekin, P. L. Hammer, A. Kogan: On connected Boolean functions, *Discr. Appl. Math.* **96–97** (1999), 337–362.
- [7] R. L. Graham, H. O. Pollak: On the addressing problem for loop switching, *Bell Sys. Tech. J.* **50** (1971), 2459–2519.
- [8] P. L. Hammer, A. Kogan, U. G. Rothblum: Evaluation, strength and relevance of variables of Boolean functions, *SIAM J. Disc. Math.* **13** (2000), 302–312.
- [9] L. Hellerstein, V. Raghavan: Exact learning of DNF formulas using DNF hypothesis, *34. STOC* (2002), 465–473.
- [10] J. Kahn, G. Kalai, N. Linial: The influence of variables on Boolean functions, *29. FOCS* (1988), 68–80.
- [11] O. Kullmann: An application of matroid theory to the SAT problem, *15th IEEE Conf. on Computational Complexity* (2000), 116–124.
- [12] O. Kullmann: The combinatorics of conflicts between clauses, *Theory and Appl. of Satisfiability Testing* (2003), Springer LNCS 2919, 426–440.
- [13] O. Kullmann: On the conflict matrix of clause-sets, Tech. Rep. CSR 7-2003, Univ. Wales at Swansea, 2003.
- [14] J. M. Laborde: Sur le cardinal maximum de la base complete d’une fonction booleenne, en fonction du nombre de conjonctions de l’une de ses formes normales, *Discrete Math.* **32** (1980), 209–212.
- [15] A. A. Levin: Comparative complexity of DNF, *Methods of Discrete Analysis* **36** (1981), 23–38. (In Russian.)
- [16] L. Lovász, M. Naor, I. Newman, A. Wigderson: Search problems in the decision tree model, *SIAM J. Disc. Math.* **8** (1995), 119–132.
- [17] C. McMullen, J. Shearer: Prime implicants, minimum covers, and the complexity of logic simplification, *IEEE Trans. on Computers* **C-35** (1986), 761–762.
- [18] J. E. Savage: *Models of Computation: Exploring the Power of Computing*. Addison Wesley, 1998.
- [19] P. Savicky, J. Sgall: DNF tautologies with a limited number of occurrences of every variable, *Theor. Comp. Sci.* **238**, (2000) 495–498.
- [20] R. H. Sloan, B. Szörényi, Gy. Turán: Projective DNF formulae and their revision, *16. COLT/Kernel* (2003), 625–639.

- [21] B. Szörényi, Gy. Turán: Decision trees and cube partitions. In preparation.
- [22] A. Urquhart: The complexity of propositional proofs, *Bull. Symb. Logic* **1** (1995), 425–467.
- [23] I. Wegener: *The Complexity of Boolean Functions*. Wiley-Teubner, 1987.