

# From Weak to Strong LP Gaps for all CSPs

Mrinalkanti Ghosh\*      Madhur Tulsiani†

July 31, 2016

## Abstract

We study the approximability of constraint satisfaction problems (CSPs) by linear programming (LP) relaxations. We show that for every CSP, the approximation obtained by a basic LP relaxation, is no weaker than the approximation obtained using relaxations given by  $\Omega\left(\frac{\log n}{\log \log n}\right)$  levels of the Sherali-Adams hierarchy on instances of size  $n$ .

It was proved by Chan et al. [FOCS 2013] that any polynomial size LP extended formulation is no stronger than relaxations obtained by a super-constant levels of the Sherali-Adams hierarchy. Combining this with our result also implies that any polynomial size LP extended formulation is no stronger than the basic LP.

Using our techniques, we also simplify and strengthen the result by Khot et al. [STOC 2014] on (strong) approximation resistance by LPs. They provided a necessary and sufficient condition under which  $\Omega(\log \log n)$  levels of the Sherali-Adams hierarchy cannot achieve an approximation better than a random assignment. We simplify their proof and strengthen the bound to  $\Omega\left(\frac{\log n}{\log \log n}\right)$  levels.

## 1 Introduction

Given a finite alphabet  $[q] = \{0, \dots, q-1\}$  and a predicate  $f : [q]^k \rightarrow \{0, 1\}$ , an instance of the problem MAX k-CSP( $f$ ) consists of (say)  $m$  constraints over a set of  $n$  variables  $x_1, \dots, x_n$  taking values in the set  $[q]$ . Each constraint  $C_i$  is of the form  $f(x_{i_1} + b_{i_1}, \dots, x_{i_k} + b_{i_k})$  for some  $k$ -tuple of variables  $(x_{i_1}, \dots, x_{i_k})$  and  $b_{i_1}, \dots, b_{i_k} \in [q]$ , and the addition is taken to be modulo  $q$ . We say an assignment  $\sigma$  to the variables satisfying the constraint  $C_i$  if  $C_i(\sigma(x_{i_1}), \dots, \sigma(x_{i_k})) = 1$ . Given an instance  $\Phi$  of the problem, the goal is to find an assignment  $\sigma$  to the variables satisfying as many constraints as possible. The approximability of the MAX k-CSP( $f$ ) problem has been extensively studied for various predicates  $f$  (see e.g., [24] for a survey), and special cases include several interesting and natural problems such as MAX 3-SAT, MAX 3-XOR and MAX-CUT.

A topic of much recent interest has been the efficacy of Linear Programming (LP) and Semidefinite Programming (SDP) relaxations. For a given instance  $\Phi$  of MAX k-CSP( $f$ ), let  $\text{OPT}(\Phi)$  denote the *fraction* of constraints satisfied by an optimal assignment, and let  $\text{FRAC}(\Phi)$  denote the value of the convex (LP/SDP) relaxation for the problem. Then, the

---

\*Toyota Technological Institute at Chicago. mkgghosh@ttic.edu

†Toyota Technological Institute at Chicago. madhurt@ttic.edu.

performance guarantee of this algorithm is given by the integrality gap which equals the supremum of  $\frac{\text{FRAC}(\Phi)}{\text{OPT}(\Phi)}$ , over all instances  $\Phi$ .

The study of unconditional lower bounds for general families of LP relaxations was initiated by Arora, Bollobás and Lovász [2] (see also [3]). They studied the Lovász-Schrijver [19] LP hierarchy and proved lower bounds on the integrality gap for Minimum Vertex Cover (their technique also yields similar bounds for MAX-CUT). De la Vega and Kenyon-Mathieu [12] and Charikar, Makarychev and Makarychev [11] proved a lower bound of  $2 - o(1)$  for the integrality gap of the LP relaxations for MAX-CUT given respectively by  $\Omega(\log \log n)$  and  $n^{\Omega(1)}$  levels of the Sherali-Adams LP hierarchy [23]. Several follow-up works have also shown lower bounds for various other special cases of the MAX k-CSP problem, both for LP and SDP hierarchies [1, 22, 27, 21, 6, 4].

A recent result by Chan et al. [7] shows a connection between strong lower bounds for the Sherali-Adams hierarchy, and lower bounds on the size of LP extended formulations for the corresponding problem. In fact, their result proved a connection not only for a lower bound on the worst case integrality gap, but for the entire *approximability curve*. We say that  $\Phi$  is  $(c, s)$ -integrality gap instance for a relaxation of MAX k-CSP( $f$ ), if we have  $\text{FRAC}(\Phi) \geq c$  and  $\text{OPT}(\Phi) < s$ . They showed that for any fixed  $t \in \mathbb{N}$ , if there exist  $(c, s)$ -integrality gap instances of size  $n$  for the relaxation given by  $t$  levels of the Sherali-Adams hierarchy, then for all  $\varepsilon > 0$  and sufficiently large  $N$ , there exists a  $(c - \varepsilon, s + \varepsilon)$  integrality gap instance of size (number of variables)  $N$ , for any linear extended formulation of size at most  $N^{t/2}$ . They also give a tradeoff (described later) when  $t$  is a function of  $n$ , which was recently improved by Kothari et al. [17].

We strengthen the above results by showing that for all  $c, s \in [0, 1]$ ,  $(c, s)$ -integrality gap instances for a “basic LP” can be used to construct  $(c - \varepsilon, s + \varepsilon)$  integrality gap instances for  $\Omega_\varepsilon\left(\frac{\log n}{\log \log n}\right)$  levels of the Sherali-Adams hierarchy. The basic LP uses only a subset of the constraints in the relaxation given by  $k$  levels of the Sherali-Adams hierarchy for MAX k-CSP( $f$ ). In particular, this shows that a lower bound on the integrality gap for the basic LP, implies a similar lower bound on the integrality gap of any polynomial size extended formulation. We note that both the above results and our result apply to all  $f, q$  and all  $c, s \in [0, 1]$ .

**Comparison with (implications of) Raghavendra’s UGC hardness result.** A remarkable result by Raghavendra [20] shows that a  $(c, s)$ -integrality gap instance for a “basic SDP” relaxation of MAX k-CSP( $f$ ) implies hardness of distinguishing instances  $\Phi$  with  $\text{OPT}(\Phi) < s$  from instances with  $\text{OPT}(\Phi) \geq c$ , assuming the Unique Games Conjecture of Khot [13]. The basic SDP considered by Raghavendra involves moments for all pairs of variables, and all subsets of variables included in a constraint. The basic LP we consider is weaker than this SDP and does not contain the positive semidefiniteness constraint.

Combining Raghavendra’s result with known constructions of integrality gaps for Unique Games by Raghavendra and Steurer [21], and by Khot and Saket [14], one can obtain a result qualitatively similar to ours, for the mixed hierarchy. In particular, a  $(c, s)$  integrality gap for the basic SDP implies a  $(c - \varepsilon, s + \varepsilon)$  integrality gap for  $\Omega((\log \log n)^{1/4})$  levels of the mixed hierarchy.

Note however, that the above result is incomparable to our result, since it starts with stronger hypothesis (a basic SDP gap) and yields a gap for the mixed hierarchy as op-

posed to the Sherali-Adams hierarchy. While the above can also be used to derive lower bounds for linear extended formulations, one needs to start with an SDP gap instance to derive an LP lower bound. The basic SDP is known to be provably stronger than the basic LP for several problems including various 2-CSPs. Also, for the worst case  $f$  for  $q = 2$ , the integrality gap of the basic SDP is  $O(2^k/k)$  [10], while that of the basic LP is  $2^{k-1}$ .

**Other related work.** The power of the basic LP for solving valued CSPs *to optimality* has been studied in several previous works. These works consider the problem of minimizing the penalty for unsatisfied constraints, where the penalties take values in  $\mathbb{Q} \cup \{\infty\}$ . Also, they study the problem not only in terms of single predicate  $f$ , but rather in terms of the constraint language generated by a given set of (valued) predicates.

It was shown by Thapper and Živný [25] that when the penalties are finite-valued, if the problem of finding the optimum solution cannot be solved by the basic LP, then it is NP-hard. Kolmogorov, Thapper and Živný [16] give a characterization of CSPs where the problem of minimizing the penalty for unsatisfied constraints can be solved *exactly* by the basic LP. Also, a recent result by Thapper and Živný [26] shows the valued CSP problem for a constraint language can be solved to optimality by a bounded number of levels of the Sherali-Adams hierarchy if and only if it can be solved by a relaxation obtained by augmenting the basic LP with constraints implied by three levels of the Sherali-Adams hierarchy. However, the above works only consider the case when the LP gives an exact solution, and do not focus on approximation.

## Our results

Our main result is the following.

**Theorem 1.1** *Let  $f : [q]^k \rightarrow \{0, 1\}$  be any predicate. Let  $\Phi_0$  be a  $(c, s)$  integrality gap instance for basic LP relaxation of MAX  $k$ -CSP ( $f$ ). Then for every  $\varepsilon > 0$ , there exists  $c_\varepsilon > 0$  such that for infinitely many  $N \in \mathbb{N}$ , there exist  $(c - \varepsilon, s + \varepsilon)$  integrality gap instances of size  $N$  for the LP relaxation given by  $c_\varepsilon \cdot \frac{\log N}{\log \log N}$  levels of the Sherali-Adams hierarchy.*

Combining the above with the connection between Sherali-Adams gaps and extended formulations by Chan et al. [7] yields the following corollary. The improved tradeoff by Kothari et al. [17] gives a better exponent for  $\log N$  than  $3/2$ .

**Corollary 1.2** *Let  $f : [q] \rightarrow \{0, 1\}$  be any predicate. Let  $\Phi_0$  be a  $(c, s)$  integrality gap instance for basic LP relaxation of MAX  $k$ -CSP ( $f$ ). Then for every  $\varepsilon > 0$ , there exists  $c_\varepsilon > 0$  such that for infinitely  $N \in \mathbb{N}$ , there exist  $(c - \varepsilon, s + \varepsilon)$  integrality gap instances of size  $N$ , for every linear extended formulation of size at most  $\exp\left(c_\varepsilon \cdot \frac{(\log N)^{3/2}}{(\log \log N)^{1/2}}\right)$ .*

As an application of our methods, we also simplify and strengthen the approximation resistance results for LPs proved by Khot et al. [15]. They studied predicates  $f : \{0, 1\}^k \rightarrow \{0, 1\}$  and provided a necessary and sufficient condition for the predicate to be strongly approximation resistant for the Sherali-Adams LP hierarchy. We say a predicate is strongly approximation resistant if for all  $\varepsilon > 0$ , it is hard to distinguish instances  $\Phi$  for which  $|\text{OPT}(\Phi) - \mathbb{E}_{x \in \{0, 1\}^k} [f(x)]| \leq \varepsilon$  from instances with  $\text{OPT}(\Phi) \geq 1 - \varepsilon$ . In the context

of the Sherali-Adams hierarchy, they showed that when this condition is satisfied, there exist instances  $\Phi$  satisfying  $\left| \text{OPT}(\Phi) - \mathbb{E}_{x \in \{0,1\}^k} [f(x)] \right| \leq \varepsilon$  and  $\text{FRAC}(\Phi) \geq 1 - \varepsilon$ , where  $\text{FRAC}(\Phi)$  is the value of the relaxation given by  $O_\varepsilon(\log \log n)$  levels of the Sherali-Adams hierarchy. We strengthen their result (and provide a simpler proof) to prove the following.

**Theorem 1.3** *Let  $f : \{0,1\}^k \rightarrow \{0,1\}$  be any predicate satisfying the condition for strong approximation resistance for LPs, given by [15]. Then for every  $\varepsilon > 0$ , there exists  $c_\varepsilon > 0$  such that infinitely many  $N \in \mathbb{N}$ , there exists an instance  $\Phi$  of MAX  $k$ -CSP( $f$ ) of size  $N$ , satisfying*

$$\left| \text{OPT}(\Phi) - \mathbb{E}_{x \in \{0,1\}^k} [f(x)] \right| \leq \varepsilon \quad \text{FRAC}(\Phi) \geq 1 - \varepsilon,$$

where  $\text{FRAC}(\Phi)$  is the value of the relaxation given by  $c_\varepsilon \cdot \frac{\log N}{\log \log N}$  levels of the Sherali-Adams hierarchy.

As before, the above theorem also yields a corollary for extended formulations.

### Proof overview and techniques

**The gap instance.** The construction of our gap instances is inspired by the construction by Khot et al. [15]. They gave a generic construction to prove integrality gaps for any approximation resistant predicate (starting from certificates of hardness in form of certain “vanishing measures”), and we use similar ideas to give a construction which can start from a basic LP integrality gap instance as a certificate, to produce a gap instance for a large number of levels. This construction is discussed in [Section 5](#).

Given an integrality gap instance  $\Phi_0$  on  $n_0$  variables, we treat this as a “template” (as in Raghavendra [20]) and generate a random instance using this. Concretely, we generate a new instance  $\Phi$  on  $n_0$  sets of  $n$  variables each. To generate a constraint, we sample a random constraint  $C_0 \in \Phi_0$ , and pick a variable randomly from each of the sets corresponding to variables in  $C_0$ . Thus, the instances generated are  $n_0$ -partite random hypergraphs, with each edge being generated according to a specified “type” (indices of sets to chose vertices from). Previous instances of gap constructions for LP and SDP hierarchies were (hyper)graphs generated according to the model  $\mathcal{G}_{n,p}$ . However, properties of random  $\mathcal{G}_{n,p}$  hypergraphs easily carry over to our instances, and we collect these properties in [Section 3](#).

The above construction ensures that if the instance  $\Phi_0$  does not have an assignment satisfying more than an  $s$  fraction of the constraints, then  $\text{OPT}(\Phi) \leq s + \varepsilon$  with high probability. Also, it is well-known that providing a good LP solution to the relaxation given by  $t$  levels of the Sherali-Adams hierarchy is equivalent to providing distributions  $\mathcal{D}_S$  on  $[q]^S$  for all sets of variables  $S$  with  $|S| \leq t$ , such that the distributions are consistent restricted to subsets i.e., for all  $S$  with  $|S| \leq t$  and all  $T \subseteq S$ , we have  $\mathcal{D}_{S|T} = \mathcal{D}_T$ . Thus, in our case, we need to produce such consistent local distributions such that the expected probability that a random constraint  $C \in \Phi$  is satisfied by the local distribution on the set of variables involved in  $C$  (which we denote as  $S_C$ ) is at least  $c - \varepsilon$ .

**Local distributions from local structure.** Most works on integrality gaps for CSPs utilize the local structure of random hypergraphs to produce such distributions. Since the girth

of a sparse random hypergraph is  $\Omega(\log n)$ , any induced subgraph on  $o(\log n)$  vertices is simply a forest. In case the induced (hyper)graph  $G_S$  on a set  $S$  is a *tree*, there is an easy distribution to consider: simply choose an arbitrary root and propagate down the tree by sampling each child conditioned on its parent. It is also easy to see that for  $T \subseteq S$ , if the induced (hyper)graph  $G_T$  is a *subtree* of  $G_S$ , then the distributions  $\mathcal{D}_S$  and  $\mathcal{D}_T$  produced as above are consistent.

The extension of this idea to forests requires some care. One can consider extending the distribution to forests by propagating independently on each tree in the forest. However, if for  $T \subseteq S$   $G_T$  is a forest while  $G_S$  is a tree, then a pair of vertices disconnected in  $G_T$  will have no correlation in  $\mathcal{D}_T$  but may be correlated in  $\mathcal{D}_S$ . This was handled, for example, in [15] by adding noise to the propagation and using a large ball  $B(S)$  around  $S$  to define  $\mathcal{D}_S$ . Then, if two vertices of  $T$  are disconnected in  $B(T)$  but connected in  $B(S)$ , then they must be at a large distance from each other. Thus, because of the noise, the correlation between them (which is zero in  $\mathcal{D}_T$ ) will be very small in  $\mathcal{D}_S$ . However, correcting approximate consistency to exact consistency incurs a cost which is exponential in the number of levels (i.e., the sizes of the sets), which is what limits the results in [15, 12] to  $O(\log \log n)$  levels. This also makes the proof a bit more involved since it requires a careful control of the errors in consistency.

**Consistent partitioning schemes.** We resolve the above consistency issue by first partitioning the given set  $S$  into a set of clusters, each of which have diameter  $\Delta_H = o(\log n)$  in the underlying hypergraph  $H$ . Since each cluster has bounded diameter, it becomes a tree when we add all the missing paths between any two vertices in the cluster. We then propagate independently on each cluster (augmented with the missing paths). This preserves the correlation between any two vertices in the same cluster, even if the path between them was not originally present in  $G_S$ .

Of course, the above plan requires that the partition obtained for  $T \subseteq S$ , is consistent with the restriction to  $T$  of partition obtained for the set  $S$ . In fact, we construct distributions over partitions  $\{\mathcal{P}_S\}_{|S| \leq t}$ , which satisfy the consistency property  $\mathcal{P}_{S|T} = \mathcal{P}_T$ . These distributions over partitions, which we call consistent partitioning schemes, are constructed in [Section 4](#).

In addition to being consistent, we require that the partitioning scheme cuts only a small number of edges in expectation, since these contribute to a loss in the LP objective. We remark that such low-diameter decompositions (known as *separating* and *padded* decompositions) have been used extensively in the theory metric embeddings (see e.g., [18] and the references therein). The only additional requirement in our application is consistency.

We obtain the decompositions by proving the (easy) hypergraph extensions the results of Charikar, Makarychev and Makarychev [9], who exhibit a metric which is similar to the shortest path metric on graphs at small distances, and has the property that its restriction to any subset of size at most  $n^{\epsilon'}$  (for an appropriate  $\epsilon' < 1$ ) is  $\ell_2$  embeddable. This is proved in [Section 3](#). We then use these in [Section 4](#) to construct the consistent partitioning schemes as described above, by applying a result of Charikar et al. [8] giving separating decompositions for finite subsets of  $\ell_2$ .

We remark that it is the consistency requirement of the partitioning procedure that limits our results to  $O\left(\frac{\log n}{\log \log n}\right)$  levels. The separation probability in the decomposition proce-

dimension grows with the dimension of the  $\ell_2$  embedding, while (to the best of our knowledge) dimension reduction procedures seem to break consistency.

## 2 Preliminaries

We use  $[n]$  to denote the set  $\{1, \dots, n\}$ . The only exception is  $[q]$ , where we overload this notation to denote the set  $\{0, \dots, q-1\}$ , which corresponds to the the alphabet for the Constraint Satisfaction Problem under consideration. We use  $\mathcal{D}_S$  and  $\mathcal{P}_S$  to denote probability distributions over (assignments to or partitions of) a set  $S$ . For  $T \subseteq S$ , the notation  $\mathcal{D}_{S|T}$  is used to denote the restriction (marginal) of the distribution  $\mathcal{D}_S$  to the set  $T$  (and similarly for  $\mathcal{P}_{S|T}$ ).

### 2.1 Constraint Satisfaction Problems

**Definition 2.1** Let  $[q]$  denote the set  $\{0, \dots, q-1\}$ . For a predicate  $f : [q]^k \rightarrow \{0, 1\}$ , an instance  $\Phi$  of MAX  $k$ -CSP $_q$  ( $f$ ) consists of a set of variables  $\{x_1, \dots, x_n\}$  and a set of constraints  $C_1, \dots, C_m$  where each constraint  $C_i$  is over a  $k$ -tuple of variables  $\{x_{i_1}, \dots, x_{i_k}\}$  and is of the form

$$C_i \equiv f(x_{i_1} + b_{i_1}, \dots, x_{i_k} + b_{i_k})$$

for some  $b_{i_1}, \dots, b_{i_k} \in [q]$ , where the addition is modulo  $q$ . For an assignment  $\sigma : \{x_1, \dots, x_n\} \mapsto [q]$ , let  $\text{sat}(\sigma)$  denote the fraction of constraints satisfied by  $\sigma$ . The maximum fraction of constraints that can be simultaneously satisfied is denoted by  $\text{OPT}(\Phi)$ , i.e.

$$\text{OPT}(\Phi) = \max_{\sigma : \{x_1, \dots, x_n\} \mapsto [q]} \text{sat}(\sigma).$$

For a constraint  $C$  of the above form, we use  $x_C$  to denote the tuple of variables  $(x_{i_1}, \dots, x_{i_k})$  and  $b_C$  to denote the tuple  $(b_{i_1}, \dots, b_{i_k})$ . We then write the constraint as  $f(x_C + b_C)$ . We also denote by  $S_C$  the set of indices  $\{i_1, \dots, i_k\}$  of the variables participating in the constraint  $C$ .

### 2.2 The LP Relaxations for Constraint Satisfaction Problems

Below we present various LP relaxations for the MAX  $k$ -CSP $_q$  ( $f$ ) problem that are relevant in this paper.

We start with the level- $t$  Sherali-Adams relaxation. The intuition behind it is the following. Note that an integer solution to the problem can be given by an assignment  $\sigma : [n] \rightarrow [q]$ . Using this, we can define  $\{0, 1\}$ -valued variables  $x_{(S, \alpha)}$  for each  $S \subseteq [n], 1 \leq |S| \leq t$  and  $\alpha \in [q]^S$ , with the intended solution  $x_{(S, \alpha)} = 1$  if  $\sigma(S) = \alpha$  and 0 otherwise. We also introduce a variable  $x_{(\emptyset, \emptyset)}$ , which equals 1. We relax the integer program and allow variables to take real values in  $[0, 1]$ . Now the variables  $\{x_{(S, \alpha)}\}_{\alpha \in [q]^S}$  give a probability distribution  $\mathcal{D}_S$  over assignments to  $S$ . We can enforce consistency between these local distributions by requiring that for  $T \subseteq S$ , the distribution over assignments to  $S$ , when marginalized to  $T$ , is precisely the distribution over assignments to  $T$  i.e.,  $\mathcal{D}_{S|T} = \mathcal{D}_T$ . The relaxation is shown in Figure 1.

The basic LP relaxation is a reduced form of the above relaxation where only those variables  $x_{(S, \alpha)}$  are included for which  $S = S_C$  is the set of CSP variables for some constraint

$$\begin{aligned}
& \text{maximize } \mathbb{E}_{C \in \Phi} \left[ \sum_{\alpha \in [q]^k} f(\alpha \cdot b_C) \cdot x_{(S_C, \alpha)} \right] \\
& \text{subject to} \\
& \sum_{\substack{\alpha \in [q]^S \\ \alpha|_T = \beta}} x_{(S, \alpha)} = x_{(T, \beta)} & \forall T \subseteq S \subseteq [n], |S| \leq t, \forall \beta \in [q]^T \\
& x_{(S, \alpha)} \geq 0 & \forall S \subseteq [n], |S| \leq t, \forall \alpha \in [q]^S \\
& x_{(\emptyset, \emptyset)} = 1
\end{aligned}$$

Figure 1: Level- $t$  Sherali-Adams LP for MAX k-CSP $_q$  ( $f$ )

C. The consistency constraints are included only for singleton subsets of the sets  $S_C$ . Note that the all the constraints for the basic LP are implied by the relaxation obtained by level  $k$  of the Sherali-Adams hierarchy.

$$\begin{aligned}
& \text{maximize } \mathbb{E}_{C \in \Phi} \left[ \sum_{\alpha \in [q]^k} f(\alpha + b_C) \cdot x_{(S_C, \alpha)} \right] \\
& \text{subject to} \\
& \sum_{j \in [q]} x_{(i, j)} = 1 & \forall i \in [n] \\
& \sum_{\substack{\alpha \in [q]^{S_C} \\ \alpha(i) = b}} x_{(S_C, \alpha)} = x_{(i, b)} & \forall C \in \Phi, i \in S_C, b \in [q] \\
& x_{(S_C, \alpha)} \geq 0 & \forall C \in \Phi, \forall \alpha \in [q]^{S_C}
\end{aligned}$$

Figure 2: Basic LP relaxation for MAX k-CSP $_q$  ( $f$ )

For an LP/SDP relaxation of MAX k-CSP $_q$ , and for a given instance  $\Phi$  of the problem, we denote by  $\text{FRAC}(\Phi)$  the LP/SDP (fractional) optimum. A relaxation is said to have a  $(c, s)$ -integrality gap if there exists a CSP instance  $\Phi$  such that  $\text{FRAC}(\Phi) \geq c$  and  $\text{OPT}(\Phi) < s$ .

### 2.3 Hypergraphs

An instance  $\Phi$  of MAX k-CSP defines a natural associated hypergraph  $H = (V, E)$  with  $V$  being the set of variables in  $\Phi$  and  $E$  containing one  $k$ -hyperedge for every constraint  $C \in \Phi$ . We remind the reader of the familiar notions of degree, paths, and cycles for the case of ( $k$ -uniform) hypergraphs:

**Definition 2.2** Let  $H = (V, E)$  be a hypergraph.

- For a vertex  $v \in V$ , the degree of the vertex  $v$  is defined to be the number of distinct hyperedges containing it.

- A simple path  $P$  is a finite alternate sequence of distinct vertices and distinct edges starting and ending at vertices, i.e.,  $P = v_1, e_1, v_2, \dots, v_l, e_l, v_{l+1}$ , where  $v_i \in V \forall i \in [l+1]$  and  $e_i \in E \forall i \in [l]$ . Furthermore,  $e_i$  contains  $v_i, v_{i+1}$  for each  $i$ . Here  $l$  is called the length of the path  $P$ . All paths discussed in this paper will be simple paths.
- A sequence  $C = (v_1, e_1, v_2, \dots, v_l, e_l, v_1)$  is called a cycle of length  $l$  if the initial segment  $v_1, e_1, \dots, v_l$  is a (simple) path,  $e_{i+1} \neq e_i$  for all  $i \in [l]$ , and  $v_1 \in e_l$ . For a path  $P$  (or cycle  $C$ ), we use  $V(P)$  (or  $V(C)$ ) to denote the set of vertices all the vertices that occurs in the edges, i.e., the set  $\{v : (\exists i \in [h])(v \in e_i)\}$ , where  $e_1, \dots, e_h$  are the edges included in  $P$  (or  $C$ ).
- For a given hypergraph  $H$ , the length of the smallest cycle in  $H$  is called the girth of  $H$ .

To observe the difference the notions of cycle in graphs and hypergraphs, it is instructive to consider the following example: let  $u, v$  be two distinct vertices in a  $k$ -uniform hypergraph for  $k \geq 3$ , and let  $e_1, e_2$  be two distinct hyperedges both containing  $u$  and  $v$ . Then  $u, e_1, v, e_2, u$  is a cycle of length 2, which cannot occur in a graph.

We shall also need the following notion of the *closure* of a set  $S \subseteq V$  in a given hypergraph  $H$ , defined by [11] for the case of graphs. A stronger notion of closure was also considered by [4].

**Definition 2.3** For a given hypergraph  $H$  and  $R \in \mathbb{N}$ , and a set  $S \subseteq V(H)$ , we denote by  $\text{cl}_R(S)$  the  $R$ -closure of  $S$  obtained by adding all the vertices in all the paths of length at most  $R$  connecting two vertices of  $S$ , i.e.,

$$\text{cl}_R(S) = S \cup \bigcup_{\substack{P: P \text{ is a path in } H \\ P \text{ connects } u, v \in S \\ |P| \leq R}} V(P).$$

For ease of notation, we use  $\text{cl}(S)$  to denote  $\text{cl}_1(S)$ .

### 3 Properties of random hypergraphs

We collect here various properties of the hypergraphs corresponding to our integrality gap instances. The gap instances we generate contain several disjoint collections of variables. Each constraint in the instance has a specified “type”, which specifies which of the collections each of the participating  $k$  variables must be sampled from. The constraint is generated by randomly sampling each of the  $k$  variables, from the collections specified by its type. This is captured by the generative model described below.

In the model below and in the construction of the gap instance, the parameter  $n_0$  should be thought of as constant, while the parameters  $n$  and  $m$  should be thought of as growing to infinity. We will choose  $m = \gamma \cdot n$  for  $\gamma = O_{k,q}(1)$ .

**Definition 3.1** Let  $n_0, k \in \mathbb{N}$  with  $k \geq 2$ . Let  $m, n > 0$  and let  $\Gamma$  be a distribution on  $[n_0]^k$ . We define a distribution  $\mathcal{H}_k(m, n, n_0, \Gamma)$  on  $k$ -uniform  $n_0$ -partite hypergraphs with  $m$  edges and  $N = n_0 \cdot n$  vertices, divided in  $n_0$  sets  $X_1, \dots, X_{n_0}$  of size  $n$  each. A random hypergraph  $H \sim \mathcal{H}_k(m, n, n_0, \Gamma)$  is generated by sampling  $m$  random hyperedges independently as follows:

- Sample a random type  $(i_1, \dots, i_k) \in [n_0]^k$  from the distribution  $\Gamma$ .



- For all  $j \in [k]$ , sample  $v_{i_j}$  independently and uniformly in  $X_{i_j}$ .
- Add the edge  $e_i = \{v_{i_1}, \dots, v_{i_k}\}$  to  $H$ .

Note that as specified above, the model may generate a multi-hypergraph. However, the number of such repeated edges is likely to be small, and we will bound these, and in fact the number of cycles of size  $o(\log n)$  in [Lemma 3.5](#).

We will study the following metrics (similar to the ones defined in [9]) in this section:

**Definition 3.2** Given a hypergraph  $H$  with vertex set  $V$ , we define two metrics  $d_\mu^H(\cdot, \cdot), \rho_\mu^H(\cdot, \cdot)$  on  $V$  as

$$d_\mu^H(u, v) := 1 - (1 - \mu)^{2 \cdot d_H(u, v)} \quad \text{and} \quad \rho_\mu^H(u, v) := \sqrt{\frac{2 \cdot d_\mu^H(u, v) + \mu}{1 + \mu}},$$

for  $u \neq v$ , where  $d_H(\cdot, \cdot)$  denotes the shortest path distance in  $H$ .

The goal of this section is to prove the following result about the local  $\ell_2$ -embeddability of the metric  $\rho_\mu$ . The proof of the theorem heavily uses results proved in [3] and [11].

**Theorem 3.3** Let  $H' \sim \mathcal{H}_k(m, n, n_0, \Gamma)$  with  $m = \gamma \cdot n$  edges and let  $\varepsilon > 0$ . Then for large enough  $n$ , with high probability (at least  $1 - \varepsilon$ , over the choice of  $H'$ ), there exists  $\delta > 0$ , constant  $c = c(k, \gamma, n_0, \varepsilon)$ ,  $\theta = \theta(k, \gamma, n_0, \varepsilon)$  and a subhypergraph  $H \subset H'$  with  $V(H) = V(H')$  satisfying the following:

- $H$  has girth  $g \geq \delta \cdot \log n$ .
- $|E(H') \setminus E(H)| \leq \varepsilon \cdot m$ .
- For all  $t \leq n^\theta$ , for  $\mu \geq c \cdot \frac{\log t + \log \log n}{\log n}$ , for all  $S \subseteq V(H')$  with  $|S| \leq t$ , the metric  $\rho_\mu^H$  restricted to  $S$  is isometrically embeddable into the unit sphere in  $\ell_2$ ,

To prove the above theorem, we will use the local structure of random (hyper)graphs. We first prove that with high probability random hypergraphs (sampled from  $\mathcal{H}_k(m, n, n_0, \Gamma)$ ) can be modified by removing a few edges to a hypergraph whose girth is  $\Omega(\log n)$  and the degree of the resulting hypergraph is bounded. The following lemma shows a possible trade-off between the degree of the hypergraph vs the number of edges required to be removed.

**Lemma 3.4** Let  $H' \sim \mathcal{H}_k(m, n, n_0, \Gamma)$  be a random hypergraph with  $m = \gamma \cdot n$  edges. Then for any  $\varepsilon > 0$ , with probability  $1 - \varepsilon$  there exists a sub-hypergraph  $H$  with  $V(H) = V(H')$  such that  $\forall u \in V(H)$ ,  $\deg_H(u) \leq 100 \cdot \log\left(\frac{n_0}{\varepsilon}\right) \cdot k \cdot \gamma$  and  $|E(H') \setminus E(H)| \leq \varepsilon \cdot m$ .

**Proof:** By linearity of expectation, the expected degree of any vertex  $v$  in  $H'$  is at most  $k \cdot \gamma$ . Let  $D = 100 \cdot \log\left(\frac{n_0}{\varepsilon}\right) \cdot k \cdot \gamma$ . Let  $S$  be the set of all vertices  $u$  such that  $\deg_{H'}(u) > D$ . Let  $E_S$  be the set of all hyperedges with one vertex in  $S$ . We shall take  $E(H) = E(H') \setminus E_S$ .

Note that for any  $u \in V(H')$ ,  $\mathbb{P}[u \in S] = \mathbb{P}[\deg_{H'}(u) \geq D] \leq \exp(-D/4)$  by a Chernoff-Hoeffding bound. We use this to bound the expected number of edges deleted.

$$\begin{aligned} \mathbb{E}[E_S] &\leq \sum_{u \in V(H')} \mathbb{E}[\deg(u) \cdot \mathbf{1}_{\{u \in S\}}] = \sum_{u \in V(H')} \mathbb{E}[\deg(u) \mid u \in S] \cdot \mathbb{P}[u \in S] \\ &\leq \sum_{u \in V(H')} \mathbb{E}[\deg(u) \mid u \in S] \cdot \exp(-D/4) \\ &\leq \sum_{u \in V(H')} (D + k\gamma) \cdot \exp(-D/4) \\ &\leq (n \cdot n_0) \cdot 2D \cdot \exp(-D/4). \end{aligned}$$

The penultimate inequality uses the independence of the hyper-edges in the generation process, which gives  $\mathbb{E}[\deg_{H'}(u) \mid \deg_{H'}(u) \geq D] \leq D + \mathbb{E}[\deg_{H'}(u)]$ . From our choice of the parameter  $D$ , we get that  $\mathbb{E}[E_S] \leq \varepsilon^2 \cdot \gamma \cdot n = \varepsilon^2 \cdot m$ . Thus, the number of edges deleted is at most  $\varepsilon \cdot m$  with probability at least  $1 - \varepsilon$ .  $\blacksquare$

The following lemma shows that the expected number of small cycles in random hypergraph is small.

**Lemma 3.5** *Let  $H \sim \mathcal{H}_k(m, n, n_0, \Gamma)$  be a random hypergraph and for  $l \geq 2$ , let  $Z_l(H)$  denote the number of cycles of length at most  $l$  in  $H$ . For  $m, n$  and  $k$  such that  $k^2 \cdot (m/n) > 1$ , we have*

$$\mathbb{E}_{H \sim \mathcal{H}_k(m, n, n_0, \Gamma)} [Z_l(H)] \leq \left(k^2 \cdot \frac{m}{n}\right)^{2l}.$$

**Proof:** Let the vertices of  $H$  correspond to the set  $[n_0] \times [n]$ . Suppose we contract the set of  $[n_0] \times \{j\}$  vertices into a single vertex  $j \in [n]$  to get a random multi-hypergraph  $H'$  on vertex set  $[n]$ . An equivalent way to view the sampling to  $H'$  is: for each  $i \in [m]$ , the  $i$ -th hyperedge  $e_i$  of  $H'$  is sampled by independently sampling  $k$  vertices (with replacement) uniformly at random from  $[n]$ . Note that the sampling of  $H'$  is independent of  $\Gamma$  in the definition of  $\mathcal{H}_k(m, n, n_0, \Gamma)$ . Clearly, a cycle of length at most  $l$  in  $H$  produces a cycle of length at most  $l$  in  $H'$ . Hence, suffices to bound the expected number of cycles in  $H'$

Given any pair  $(u', v')$  of vertices of  $H'$ , for  $u' \neq v'$ , the probability of the pair  $(u', v')$  belonging together in some edge of  $H'$  is at most  $\frac{mk^2}{n^2}$ . Consider a given  $h$ -tuple of vertices  $\mathbf{u} = (u_{i_1}, \dots, u_{i_h})$ . Note that we require that edges participating in a cycle be distinct. So, the probability that  $\mathbf{u}$  is part of a cycle in  $H'$ , i.e., there exists distinct edges  $e_j \in H'$  for  $j \in [h]$  such that  $u_{i_j}, u_{i_{j+1}} \in e_j$  for  $j \in [h-1]$ , and  $u_{i_1}, u_{i_h} \in e_h$  is at most  $\left(\frac{mk^2}{n^2}\right)^h$ . As a result, expected number of cycles of length  $h$  in  $H'$  is bounded above by:

$$\binom{n}{h} \left(\frac{mk^2}{n^2}\right)^h \leq n^h \left(\frac{mk^2}{n^2}\right)^h = \left(k^2 \cdot \frac{m}{n}\right)^h$$

From the geometric form of the bound, it follows that expected number of cycles of length at most  $l$  in  $H'$  is at most  $\frac{\left(k^2 \cdot \frac{m}{n}\right)^{l+1}}{\left(k^2 \cdot \frac{m}{n}\right) - 1} < \left(k^2 \cdot \frac{m}{n}\right)^{2l}$ .  $\blacksquare$

Using the above lemma, it is easy to show that one can remove all small cycles in a random hypergraph by deleting only a small number of edges.

**Corollary 3.6** *Let  $H \sim \mathcal{H}_k(m, n, n_0, \Gamma)$  be a random hypergraph with  $m = \gamma \cdot n$  for  $\gamma > 1$  and  $k \geq 2$ . Then, there exists  $\delta = \delta(\gamma) > 0$  such that with probability  $1 - n^{-1/6}$ , all cycles of length at most  $\delta \cdot \log n$  in  $H$  can be removed by deleting at most  $n^{2/3}$  edges.*

**Proof:** As above, let  $Z_l$  denote the number of cycles of length at most  $l$ . With the choice of  $m, n$ , and  $k$ , we have  $k^2 \cdot \frac{m}{n} \geq 2$ . By [Lemma 3.5](#),  $\mathbb{E}[Z_l] \leq (k^2 \cdot \frac{m}{n})^{2l}$ . Since  $m = \gamma \cdot n$ , there exists a  $g = \delta \cdot \log n$  such that  $\mathbb{E}[Z_l] \leq \sqrt{n}$ . By Markov's inequality,  $\mathbb{P}[Z_l \geq n^{2/3}] \leq n^{-1/6}$ . Thus, with probability  $1 - n^{-1/6}$ , one can remove all cycles of length at most  $\delta \cdot \log n$  by deleting at most  $n^{2/3}$  edges.  $\blacksquare$

One can also extend the analysis in [\[3\]](#) to show that the hypergraphs are locally sparse i.e., the number of edges contained in a small set of vertices is small. For a hypergraph  $H$  and a set  $S \subseteq V(H)$ , we use  $E(S)$  to denote the edges contained in the set  $S$ .

**Definition 3.7** *We say that  $S \subseteq V(H)$  is  $\eta$ -sparse if  $|E(S)| \leq \frac{|S|}{k-1-\eta}$ . We call an  $k$ -uniform hypergraph  $H$  on  $N$  vertices to be  $(\tau, \eta)$ -sparse if all subsets  $S \subset V(H)$ ,  $|S| \leq \tau \cdot |V(H)|$ ,  $S$  is  $\eta$ -sparse. We call  $H$  to be  $\eta$ -sparse if it is  $(1, \eta)$ -sparse, i.e., all subsets of vertices of  $H$  are sparse.*

We note here that while this notion of sparsity is a generalization of that considered in [\[3\]](#), it is also identical to the notions of expansion considered in works in proof complexity (see e.g., [\[5\]](#)) and later in works on integrality gaps [\[1, 6, 4\]](#). We prove that random hypergraphs generated with our model are locally sparse:

**Lemma 3.8** *Let  $\eta < 1/4$  and  $m = \gamma \cdot n$  for  $\gamma > 1$ . Then for  $\tau \leq \frac{1}{n_0} \cdot \left(\frac{1}{e \cdot k^{3k} \cdot \gamma}\right)^{1/\eta}$  the following holds:*

$$\mathbb{P}_{H \sim \mathcal{H}_k(m, n, n_0, \Gamma)} [H \text{ is not } (\tau, \eta)\text{-sparse}] \leq 3 \cdot \left(\frac{k^{3k} \cdot \gamma}{n^{\eta/4}}\right)^{1/k}.$$

We note that we will require the sparsity  $\eta$  to be  $O_{k, \gamma}(1/(\log n))$ . This gives sparsity only for sublinear size sets, as compared to sets of size  $\Omega(n)$  in previous works where  $\eta$  is a constant. For the proof of the lemma, we follow an approach similar to that of [Lemma 3.5](#): we collapse the vertices of  $H$  of the form  $[n_0] \times \{j\}$  to vertex  $j \in [n]$  to construct  $H'$ , and thus reducing the problem to random multi-hypergraph form a random multipartite hypergraph. The rest proof of the lemma is along the lines of several known proofs [\[1, 6\]](#) and we defer the details to [Appendix A](#).

Charikar et al. [\[9\]](#) prove an analogue of [Theorem 3.3](#) for metrics defined on locally-sparse graphs. In fact, they use a consequence of sparsity, which they call  $\ell$ -path decomposability. To this end, we define the *incidence graph*<sup>1</sup> associated with a hypergraph, on which we will apply their result.

**Definition 3.9** *Let  $H = (V(H), E(H))$  be a  $k$ -uniform hypergraph. We define its incidence graph as the bipartite graph  $G_H$  defined on vertex sets  $V(H)$  and  $E(H)$ , and edge set  $\mathcal{E}$  defined as*

$$\mathcal{E} := \{(v, e) \mid v \in V(H), e \in E(H), v \in e\}.$$

<sup>1</sup>This is the same notion as the constraint-variable graph considered in various works on lower bounds for CSPs.

Note that for any  $u, v \in V(H)$ , we have  $d_{G_H}(u, v) = 2 \cdot d_H(u, v)$ . We prove that for a locally sparse hypergraph  $H$ , its incidence graph  $G_H$  is also locally sparse.

**Lemma 3.10** *Let  $H$  be a  $k$ -uniform  $(\tau, \eta)$ -sparse hypergraph on  $N$  vertices with  $m = \gamma \cdot n$  hyperedges. Then the incidence graph  $G_H$  is  $(\tau', \eta')$  sparse for  $\tau' = \tau/k \cdot (1+\gamma)$  and  $\eta' = \eta/(1+\eta)$ .*

**Proof:** Let  $\tau' = \tau/k \cdot (1+\gamma)$  and let  $G_H$  be the incidence graph with  $N + m = (1 + \gamma) \cdot N$  vertices. Let  $G'$  be the densest subgraph of  $G_H$ , among all subgraphs of size at most  $\tau' \cdot (N + m)$ . Let the vertex set of  $G'$  be  $V' \cup E'$  where  $V' \subseteq V(H)$  and  $E' \subseteq E(H)$ , and let the edge-set be  $\mathcal{E}'$ . There cannot be any isolated vertices in  $G'$  since removing those will only increase the density.

Let  $S \subseteq V(H)$  be the set of all vertices contained in all edges in  $E'$  i.e.,  $S := \{v \in V(H) \mid \exists e \in E' \text{ s.t. } v \in e\}$ . Note that  $V' \subseteq S$ , since there are no isolated vertices, and  $E' \subseteq E(S)$ , where  $E(S)$  denotes the set of hyperedges contained in  $S$ .

By our choice of parameters,  $|S| \leq k \cdot |E'| \leq k \cdot \tau' \cdot (N + m) \leq \tau \cdot N$ . Thus, using the sparsity of  $H$ , we have

$$|E'| \leq |E(S)| \leq \frac{|S|}{k-1-\eta}.$$

Also, since each hyperedge of  $E'$  can include at most  $k$  vertices in  $S$ , and since each edge in  $\mathcal{E}'$  is incident on a vertex in  $V'$ , we have

$$|S| - |V'| \leq k \cdot |E'| - |\mathcal{E}'|.$$

Combining the two inequalities gives

$$(k-1-\eta) \cdot |E'| \leq |V'| + k \cdot |E'| - |\mathcal{E}'| \Rightarrow |\mathcal{E}'| \leq (1+\eta) \cdot |E'| + |V'|.$$

Hence, we get that  $|\mathcal{E}'| \leq \frac{|V'|+|E'|}{(1-\eta')}$  for  $\eta' = \frac{\eta}{(1+\eta)}$ . ■

Charikar et al. [9] defined the following structural property of a graph.

**Definition 3.11 ([9])** *A graph  $G$  is  $\ell$ -path decomposable if every 2-connected subgraph  $G'$  of  $G$ , such that  $G'$  is not an edge, contains a path of length  $\ell$  such that every vertex of the path has degree at most 2 in  $G'$ .*

The above property was also implicitly used by Arora et al. ([3]), who proved the following (see Lemma 2.12 in [3]):

**Lemma 3.12** *Let  $\ell > 0$  be an integer and  $0 < \eta < \frac{1}{3\ell-1} < 1$ . Let  $G$  be a  $\eta$ -sparse graph with girth  $g > \ell$ . Then  $G$  is  $\ell$ -path decomposable.*

Recall that we defined the metrics  $d_\mu$  and  $\rho_\mu$  on  $H$  as (for  $u \neq v$ ):

$$d_\mu^H(u, v) := 1 - (1 - \mu)^{2 \cdot d_H(u, v)} \quad \text{and} \quad \rho_\mu^H(u, v) := \sqrt{\frac{2 \cdot d_\mu^H(u, v) + \mu}{1 + \mu}},$$

For a graph  $G$ , we define the following two metrics, for  $u \neq v$ :

$$d_\mu^G(u, v) := 1 - (-1)^{d_G(u, v)} (1 - \mu)^{d_G(u, v)} \quad \text{and} \quad \rho_\mu^G(u, v) := \sqrt{\frac{2 \cdot d_\mu^G(u, v) + \mu}{1 + \mu}}.$$

We note that if  $H$  is a hypergraph and  $G_H$  is its incidence graph, then the metrics  $d_\mu^{G_H}$  and  $\rho_\mu^{G_H}$  restricted to  $V(H)$ , coincide with the metrics  $d_\mu$  and  $\rho_\mu$  defined on  $H$ . Charikar et al. proved the following theorem (see Theorem 5.2) in [11].

**Theorem 3.13 ([11])** *Let  $G$  be a graph on  $n'$  vertices with maximum degree  $D$ . Let  $t < \sqrt{n'}$  and  $\ell > 0$  be such that for  $t' = D^{\ell+1} \cdot t$ , every subgraph of  $G$  on at most  $t'$  vertices is  $\ell$ -path decomposable. Also, let  $\mu$ ,  $t$  and  $\ell$  satisfy the relation  $(1 - \mu)^{\ell/9} \leq \frac{\mu}{2^{(t+1)}}$ . Then for every subset  $S$  of at most  $t$  vertices there exists a mapping  $\psi_S$  from  $S$  to unit sphere in  $\ell_2$  such that all  $u, v \in S$ :*

$$\|\psi_S(u) - \psi_S(v)\|_2 = \rho_\mu^G(u, v).$$

We use this theorem to prove the main theorem of the section.

**Proof of Theorem 3.3:** Let  $H' \sim \mathcal{H}_k(m, n, n_0, \Gamma)$  with  $m = \gamma \cdot n$  hyperedges and  $N = n_0 \cdot n$  vertices. Given  $\varepsilon > 0$ , from Lemma 3.4 we have that with high probability at least  $1 - \varepsilon/2$ , there exists  $H_1$  such that the maximum degree of  $H_1$  is at most  $D = 100 \cdot \log\left(\frac{2n_0}{\varepsilon}\right) \cdot k \cdot \gamma$  with  $|E(H') \setminus E(H_1)| \leq (\varepsilon/2) \cdot m$ .

Using Corollary 3.6 we also have that there exists  $\delta > 0$ , such that with probability at least  $1 - \varepsilon/4$  (for large enough  $n$ )  $H'$  has a sub-hypergraph  $H_2$  with  $g \geq \delta \cdot \log n$  and  $|E(H') \setminus E(H_2)| \leq (\varepsilon/4) \cdot m$ . By Lemma 3.8, there exists  $\eta = \Omega_{n_0, k, \gamma, \varepsilon}(1/(\log n))$  such that  $H'$  is  $(\tau, \eta)$ -sparse with probability at least  $1 - \varepsilon/4$ , for  $\tau \geq n^{-1/4}$ .

Hence with probability  $1 - \varepsilon$ , we have that  $H = (V(H'), E(H_1) \cap E(H_2))$  satisfies:

- Degree of  $H$  is bounded above by  $D$ .
- $H$  is  $(\tau, \eta)$ -sparse (for  $\tau \geq n^{-1/4}$  and  $\eta = \Omega_{n_0, k, \gamma, \varepsilon}(1/(\log n))$ ).
- Girth of  $H$  is at least  $g > \delta \cdot \log n$ .
- $|E(H') \setminus E(H)| \leq \varepsilon \cdot m$ .

We now show that the metric  $\rho_\mu^H$  is locally  $\ell_2$  embeddable.

Let  $G = G_H$  be the incidence graph for the hypergraph  $H$ . Note that  $N \leq |V(G)| \leq N \cdot (1 + \gamma)$  and degree of  $G$  is also bounded by  $D$ . Since a cycle in  $G$  is also a cycle in  $H$ , the girth of  $G$  is at also least  $g \geq \delta \cdot \log n$ .

By Lemma 3.10, we have  $G$  is  $(\frac{\tau}{k(1+\gamma)}, \frac{\eta}{1+\eta})$ -sparse. By Lemma 3.12, any subgraph of  $G$  on at most  $\frac{\tau}{k(1+\gamma)} \cdot (N + m)$  vertices is  $\ell$ -path decomposable for any  $\ell \leq \min\{g, 1/(4\eta)\}$ . Since  $D = 100 \cdot k\gamma \cdot \log(2n_0/\varepsilon)$ , there exists  $\ell_0 = \Omega_{k, \gamma, n_0, \varepsilon}(\log n)$  such that  $D^{\ell_0+1} \leq n^{1/6}$ . We choose  $\ell = \min\{g, 1/(4\eta), \ell_0\}$ .

Let  $\mu_0$  be the smallest  $\mu$  such that  $\exp(-\mu\ell/9) \leq \frac{\mu}{2^{(t+1)}}$  (note that  $\frac{1}{\mu} \cdot \exp(-\mu\ell/9)$  is decreasing in  $\mu$ ). Since we must have  $\mu \geq 1/\ell$ , there exists a  $\mu_0$  satisfying

$$\mu_0 \leq \frac{9}{\ell} \cdot (\ln(2(t+1)) + \ln \ell).$$

From our choice of  $\ell$ , there exist constants  $c = c(k, \gamma, n_0, \varepsilon)$  and  $\theta = \theta(k, \gamma, n_0, \varepsilon) < 1/2$  such that  $\mu_0 \leq c \cdot \frac{\log t + \log \log n}{\log n} < 1$  when  $t \leq n^\theta$ . Then, for any  $\mu \in [\mu_0, 1)$ , we have  $(1 - \mu)^{\ell/9} \leq \exp(-\mu\ell/9) \leq \frac{\mu}{2^{(t+1)}}$ .

We can now apply [Theorem 3.13](#) to construct the embedding. Given any subset  $S$  of  $V(H)$  of size at most  $t \leq n^\theta$ , note that  $S$  is also a subset of  $V(G)$ . Moreover, we have  $t \leq n^\theta \leq (N+m)^{1/2}$ . Also, we have  $t \cdot D^{\ell+1} \leq n^{1/2} \cdot n^{1/6} = n^{2/3} \leq \frac{\tau}{k(\gamma+1)} \cdot (N+m)$ . Thus, any subgraph of  $G$  on  $t \cdot D^{\ell+1}$  vertices is  $\ell$ -path decomposable.

Thus, when  $\mu \geq \mu_0$ , by [Theorem 3.13](#) there exists a mapping  $\psi_S$  from  $S$  to the unit sphere, such that for all  $u, v \in S$ , we have

$$\|\psi_S(u) - \psi_S(v)\|_2 = \rho_\mu^G(u, v) = \rho_\mu^H(u, v),$$

where the last equality uses the fact that for all  $u, v \in V(H)$ ,  $\rho_\mu^H(u, v) = \rho_\mu^G(u, v)$  since  $d_G(u, v) = 2 \cdot d_H(u, v)$ .  $\blacksquare$

## 4 Decompositions of hypergraphs from local geometry

We will construct the Sherali-Adams solution by partitioning the given subset of vertices into trees, and then creating a natural distribution over satisfying assignments on trees. We define below the kind of partitions we need.

**Definition 4.1** *Let  $X$  be a finite set. For a set  $S$ , let  $\mathcal{P}_S$  denote a distribution over partitions of  $S$ . For  $T \subseteq S$ , let  $\mathcal{P}_{S|T}$  be the distribution over partitions of  $T$  obtained by restricting the partitions in  $\mathcal{P}_S$  to the set  $T$ . We say that a collection of distributions  $\{\mathcal{P}_S\}_{|S| \leq t}$  forms a consistent partitioning scheme of order  $t$ , if*

$$\forall S \subseteq X, |S| \leq t \text{ and } \forall T \subseteq S \quad \mathcal{P}_T = \mathcal{P}_{S|T}.$$

In addition to being consistent as described above, we also require the distributions to have small probability of cutting the edges for the hypergraphs corresponding to our CSP instances. We define this property below.

**Definition 4.2** *Let  $H = (V, E)$  be a  $k$ -uniform hypergraph. Let  $\{\mathcal{P}_S\}_{|S| \leq t}$  be a consistent partitioning scheme of order  $t$  for the vertex set  $V$ , with  $t \geq k$ . We say the scheme  $\{\mathcal{P}_S\}_{|S| \leq t}$  is  $\varepsilon$ -sparse for  $H$  if*

$$\forall e \in E \quad \mathbb{P}_{P \sim \mathcal{P}_e} [P \neq \{e\}] \leq \varepsilon.$$

In this section, we will prove that the hypergraphs arising from random CSP instances admit sparse and consistent partitioning schemes. Recall that for a hypergraph  $H$ , we define ([Theorem 3.2](#)) the metrics  $d_\mu^H$  and  $\rho_\mu^H$  as:

$$d_\mu^H(u, v) := 1 - (1 - \mu)^{2 \cdot d_H(u, v)} \quad \text{and} \quad \rho_\mu^H(u, v) := \sqrt{\frac{2 \cdot d_\mu^H(u, v) + \mu}{1 + \mu}},$$

**Lemma 4.3** *Let  $H = (V, E)$  be  $k$ -uniform hypergraph and let  $d_\mu$  be the metric as defined above. Let  $H$  be such that for all sets  $S \subseteq V$  with  $|S| \leq t$ , the metric induced on  $\rho_\mu$  on  $S$  is isometrically embeddable into  $\ell_2$ . Then, there exists  $\varepsilon \leq 10k \cdot \sqrt{\mu} \cdot t$  and  $\Delta_H = O(1/\mu)$  such that  $H$  admits an  $\varepsilon$ -sparse consistent partitioning scheme of order  $t$ , with each partition consisting of clusters of diameter at most  $\Delta_H$  in  $H$ .*

We use the following result of Charikar et al. [8] which shows that low-dimensional metrics have good *separating decompositions* with bounded diameter i.e., decompositions which have a small probability of separating points at a small distance.

**Theorem 4.4 ([8])** *Let  $W$  be a finite collection of points in  $\mathbb{R}^d$  and let  $\Delta > 0$  be given. Then there exists a distribution  $\mathcal{P}$  over partitions of  $W$  such that*

- $\forall P \in \text{Supp}(\mathcal{P})$ , each cluster in  $P$  has  $\ell_2$  diameter at most  $\Delta$ .
- For all  $x, y \in W$

$$\mathbb{P}_{P \sim \mathcal{P}} [P \text{ separates } x \text{ and } y] \leq 2\sqrt{d} \cdot \frac{\|x - y\|_2}{\Delta}.$$

We also need the observation that the partitions produced by the above theorem are consistent, assuming the set  $S$  considered above lie in a fixed bounded set (using a trivial modification of the procedure in [8]). For the sequel, we use  $B(x, \delta)$  to denote the  $\ell_2$  ball around  $x$  of radius  $\delta$  and  $B_H(u, r)$  to denote a ball of radius  $r$  around a vertex  $u \in V(H)$ . Thus,

$$B(x, \delta) := \{y \mid \|x - y\|_2 \leq \delta\} \quad \text{and} \quad B_H(u, r) := \{v \in V \mid d_H(u, v) \leq r\}.$$

The balls  $B(S, \delta)$  and  $B_H(S, r)$  are defined similarly.

**Claim 4.5** *Let  $S$  and  $T$  be sets such that  $T \subseteq S$ . Let  $W_S = \{w_u\}_{u \in S}$  and  $W_T = \{w'_u\}_{u \in T}$  be  $\ell_2$ -embeddings of  $S$  and  $T$  satisfying  $\varphi(W_T) \subseteq W_S \subseteq B(0, R_0) \subset \mathbb{R}^d$ , for some unitary transformation  $\varphi$  and  $R_0 > 0$ . Let  $\mathcal{P}_S$  and  $\mathcal{P}_T$  be distributions over partitions of  $S$  and  $T$  respectively, induced by partitions on  $W_S$  and  $W_T$  as given by [Theorem 4.4](#). Then*

$$\mathcal{P}_{S|T} = \mathcal{P}_T.$$

**Proof:** The claim follows simply by considering (a trivial modification of) the algorithm of [8]. For a given set  $W$  and a parameter  $\Delta$ , they produce a partition using the following procedure:

- Let  $W' = W$ .
- Repeat until  $W' = \emptyset$ 
  - Pick a random point  $x$  in  $B(W, \Delta/2)$  according to the Haar measure. Let  $C_x = B(x, \Delta/2) \cap W'$ .
  - If  $C_x \neq \emptyset$ , set  $W' = W' \setminus C_x$ . Output  $C_x$  as a cluster in the partition.

[8] show that the above procedure produces a distribution over partitions satisfying the conditions in [Theorem 4.4](#). We simply modify the procedure to sample a random point  $x$  in  $B(0, R_0 + \Delta/2)$  instead of  $B(S, \Delta/2)$ . This does not affect the separation probability of any two points, since the only non-empty clusters are still produced by the points in  $B(S, \Delta/2)$ .

Let  $P$  be a partition of  $S$  produced by the above procedure when applied to the point set  $W_S$ , and let  $P'$  be a random partition produced when applied to the point set  $\varphi(W_T)$ . It is easy to see from the above procedure that the distribution  $\mathcal{P}_T$  is invariant under a unitary transformation of  $W_T$ . By coupling the random choice of a point in  $B(0, R_0 + \Delta/2)$  chosen at each step in the procedures applied to  $W_S$  and  $\varphi(W_T) \subseteq W_S$ , we get that  $P(T) = P'$  i.e., the partition  $P$  restricted to  $T$  equals  $P'$ . Thus, we get  $\mathcal{P}_{S|T} = \mathcal{P}_T$ .  $\blacksquare$

We can use the above to prove [Lemma 4.3](#).

**Proof of Lemma 4.3:** Given a set  $S$ , let  $W_S$  be an  $\ell_2$  embedding of the metric  $\rho_\mu$  restricted to  $S$ . Since,  $|S| \leq t$ , we can assume  $W_S \in \mathbb{R}^t$ . We apply partitioning procedure of Charikar et al. from [Theorem 4.4](#) with  $\Delta = 1/2$ . From the definition of the metric  $\rho_\mu^H$ , we get that there exists a  $\Delta_H = O(1/\mu)$  such that  $\rho_{u,v}^H \leq 1/2 \Rightarrow d_H(u,v) \leq \Delta_H$ . Moreover, for  $u, v$  contained in an edge  $e$ , we have that  $\rho_\mu(u,v) \leq \sqrt{5\mu}$  and hence the probability that  $u$  and  $v$  are separated is at most  $10\sqrt{\mu \cdot t}$ . Thus, the probability that any vertex in  $e$  is separated from  $u$  is at most  $10k \cdot \sqrt{\mu \cdot t}$ .

Finally, for any  $S \subseteq T$ , if  $W_S$  and  $W_T$  denote the corresponding  $\ell_2$  embeddings, by the rigidity of  $\ell_2$  we have that for  $\varphi(W_T) \subseteq W_S$  for some unitary transformation  $\varphi$ . Thus, by [Claim 4.5](#), we get that this is a consistent partitioning scheme of order  $t$ . ■

## 5 The Sherali-Adams Integrality Gaps construction

### 5.1 Integrality Gaps from the Basic LP

Recall that the basic LP relaxation for MAX k-CSP $_q$  ( $f$ ) as given in [Fig. 2](#). In this section, we will prove [Theorem 1.1](#). We recall the statement below.

**Theorem 1.1** *Let  $f : [q]^k \rightarrow \{0, 1\}$  be any predicate. Let  $\Phi_0$  be a  $(c, s)$  integrality gap instance for basic LP relaxation of MAX k-CSP ( $f$ ). Then for every  $\varepsilon > 0$ , there exists  $c_\varepsilon > 0$  such that for infinitely many  $N \in \mathbb{N}$ , there exist  $(c - \varepsilon, s + \varepsilon)$  integrality gap instances of size  $N$  for the LP relaxation given by  $c_\varepsilon \cdot \frac{\log N}{\log \log N}$  levels of the Sherali-Adams hierarchy.*

Let  $\Phi_0$  be a  $(c, s)$  integrality gap instance for the basic LP relaxation for MAX k-CSP $_q$  ( $f$ ) with  $n_0$  variables and  $m_0$  constraints. We use it to construct a new integrality gap instance  $\Phi$ . The construction is similar to the gap instances constructed by Khot et al. [[15](#)] discussed in the next section. However, we describe this construction first since it's simpler. The procedure for constructing the instance  $\Phi$  is described in [Fig. 3](#).

Given: A  $(c, s)$  gap instance  $\Phi_0$  on  $n_0$  variables, for the basic LP.

Output: An instance  $\Phi$  with  $N = n \cdot n_0$  variables and  $m$  constraints.

The variables are divided into  $n_0$  sets  $X_1, \dots, X_{n_0}$ , one for each variable in  $\Phi_0$ . We generate  $m$  constraints independently at random as follows:

1. Sample a random constraint  $C_0 \sim \Phi_0$ . Let  $S_{C_0} = \{i_1, \dots, i_k\} \subseteq [n_0]$  denote the set of variables in this constraint.
2. For each  $j \in [k]$ , sample a random variable  $x_{i_j} \in X_{i_j}$ .
3. Add the constraint  $f((x_{i_1}, \dots, x_{i_k}) + b_{C_0})$  to the instance  $\Phi$ .

Figure 3: Construction of the gap instance  $\Phi$



## Soundness

We first prove that no assignment satisfies more than  $s + \varepsilon$  fraction of constraints for the above instance.

**Lemma 5.1** *For every  $\varepsilon > 0$ , there exists  $\gamma = \gamma(\varepsilon, n_0, q)$  such that for an instance  $\Phi$  generated by choosing at least  $\gamma \cdot n$  constraints independently at random as above, we have with probability  $1 - \exp(-\Omega(n))$ ,  $\text{OPT}(\Phi) < s + \varepsilon$ .*

**Proof:** Fix an assignment  $\sigma \in [q]^N$ . We will first consider  $\mathbb{E}[\text{sat}_\Phi(\sigma)]$  for a randomly generated  $\Phi$  as above.

$$\begin{aligned} \mathbb{E}_\Phi[\text{sat}_\Phi(\sigma)] &= \mathbb{E}_{C_0 \in \Phi_0} \mathbb{E}_{x_{i_1} \in X_{i_1}} \cdots \mathbb{E}_{x_{i_k} \in X_{i_k}} [f(\sigma(x_{i_1}) + b_{i_1}, \dots, \sigma(x_{i_k}) + b_{i_k})] \\ &= \mathbb{E}_{C_0 \in \Phi_0} \mathbb{E}_{Z_1, \dots, Z_{n_0}} [f(Z_{C_0} + b_{C_0})], \end{aligned}$$

where for each  $i \in [n_0]$ ,  $Z_i$  is an independent random variable with the distribution

$$\mathbb{P}[Z_i = b] := \mathbb{E}_{x \in X_i} [\mathbb{1}_{\{\sigma(x)=b\}}],$$

and  $Z_{C_0}$  denotes the collection of variables in the constraint  $C_0$  i.e.,  $Z_{C_0} = \{Z_i\}_{i \in S_{C_0}}$ . Thus, the random variables  $Z_1, \dots, Z_{n_0}$  define a random assignment to the variables in  $\Phi_0$ , which gives, for any  $\sigma$

$$\mathbb{E}_\Phi[\text{sat}_\Phi(\sigma)] = \mathbb{E}_{C_0 \in \Phi_0} \mathbb{E}_{Z_1, \dots, Z_{n_0}} [f(Z_{C_0} + b_{C_0})] < s.$$

Consider a randomly added constraint  $C$  to the instance  $\Phi$ . We have that

$$\mathbb{P}[C(\sigma) = 1] = \mathbb{E}_\Phi[\text{sat}_\Phi(\sigma)] < s,$$

for any fixed  $\sigma$  over a random choice of the constraint  $C$ . Thus, for an instance  $\Phi$  with  $m$  independently and randomly generated constraints, we have

$$\begin{aligned} \mathbb{P}_\Phi[\text{sat}_\Phi(\sigma) \geq s + \varepsilon] &\leq \mathbb{P}_\Phi[\text{sat}_\Phi(\sigma) \geq \mathbb{E}_\Phi[\text{sat}_\Phi(\sigma)] + \varepsilon] \\ &= \mathbb{P}_\Phi \left[ \mathbb{E}_{C \in \Phi} [\mathbb{1}_{\{C(\sigma)=1\}}] \geq \mathbb{E}_\Phi[\text{sat}_\Phi(\sigma)] + \varepsilon \right] \\ &\leq \exp(-\Omega(\varepsilon^2 \cdot m)). \end{aligned}$$

Taking a union bound over all assignments, we get

$$\mathbb{P}_\Phi[\exists \sigma \text{ sat}_\Phi(\sigma) \geq s + \varepsilon] \leq q^{n \cdot n_0} \cdot \exp(-\varepsilon^2 \cdot m),$$

which is at most  $\exp(-\Omega(n))$  for  $m = O(((\log q)/\varepsilon^2) \cdot n \cdot n_0)$ . ■

## Completeness

To prove the completeness, we first observe that the instance  $\Phi$  as constructed above is also a gap instance for the basic LP. We will then “boost” this hardness to many levels of the Sherali-Adams hierarchy.

**Lemma 5.2** For every  $\varepsilon > 0$ , there exists  $\gamma = \gamma(\varepsilon)$  such that for an instance  $\Phi$  generated by choosing at least  $\gamma \cdot n$  constraints independently at random as above, with probability  $1 - \exp(-\Omega(n))$  there exist distributions  $\overline{\mathcal{D}}_{S_C}$  over  $[q]^{S_C}$  for each  $C \in \Phi$ , and distributions  $\overline{\mathcal{D}}_i$  over  $[q]$  for each variable  $x_i \in [n \cdot n_0]$ , satisfying

- For all  $C \in \Phi$  and all  $i \in S_C$ ,  $\overline{\mathcal{D}}_{S_C|_{\{i\}}} = \overline{\mathcal{D}}_i$ .
- The distributions satisfy  $\mathbb{E}_{C \in \Phi} \mathbb{E}_{\alpha \sim \overline{\mathcal{D}}_{S_C}} [f(\alpha + b_C)] \geq c - \frac{\varepsilon}{10}$ .

**Proof:** For each  $C_0 \in \Phi_0$  and each  $j \in [n_0]$ , let  $\mathcal{D}_{S_{C_0}}^{(0)}$  and  $\mathcal{D}_j^{(0)}$  denote the basic LP solution satisfying

$$\mathcal{D}_{S_{C_0}|j}^{(0)} = \mathcal{D}_j^{(0)} \quad \forall C_0 \in \Phi_0 \quad \forall j \in S_{C_0} \quad \text{and} \quad \mathbb{E}_{C_0 \in \Phi_0} \mathbb{E}_{\alpha \sim \mathcal{D}_{S_{C_0}}^{(0)}} [f(\alpha + b_{C_0})] \geq c.$$

Each constraint  $C \in \Phi$  is sampled according to some constraint  $C_0 \in \Phi_0$ , and we take  $\overline{\mathcal{D}}_{S_C} := \mathcal{D}_{S_{C_0}}^{(0)}$  for the corresponding constraint  $C_0 \in \Phi_0$ . Also, each variable  $x_i$  for  $i \in [n_0 \cdot n]$ , belongs to one of the sets  $X_j$  for  $j \in [n_0]$ , and we take  $\overline{\mathcal{D}}_i := \mathcal{D}_j^{(0)}$  for the corresponding  $j \in [n_0]$ .

The consistency of the distributions follows immediately from the construction of the instance  $\Phi$ . Let  $C \in \Phi$  be any constraint and let  $C_0$  be the corresponding constraint in  $\Phi_0$ . If  $S_{C_0} = (j_1, \dots, j_k)$ , then  $S_C = (i_1, \dots, i_k)$  where each  $i_r \in \{j_r\} \times [n]$  for all  $r \in [k]$ . Thus, for any  $r \in [k]$ ,

$$\overline{\mathcal{D}}_{S_C|i_r} = \mathcal{D}_{S_{C_0}|j_r}^{(0)} = \mathcal{D}_{j_r}^{(0)} = \overline{\mathcal{D}}_{i_r}.$$

To bound the objective value, we again consider its expectation over a randomly generated instance  $\Phi$ . Let  $C$  be a random constraint added to  $\Phi$ . Then, if we define  $\overline{\mathcal{D}}_{S_C}$  as above for this constraint, we have

$$\mathbb{E}_C \mathbb{E}_{\alpha \sim \overline{\mathcal{D}}_{S_C}} [f(\alpha + b_C)] = \mathbb{E}_{C_0 \in \Phi_0} \mathbb{E}_{\alpha \sim \mathcal{D}_{S_{C_0}}^{(0)}} [f(\alpha + b_{C_0})] \geq c.$$

Thus, the expected contribution of each constraint is at least  $c$ . The probability that the average of  $m$  constraints deviates by at least  $\varepsilon/10$  from the expectation, is at most  $\exp(-\Omega(\varepsilon^2 \cdot m))$ . There exists  $\gamma = O(1/\varepsilon^2)$  such that for  $m \geq \gamma \cdot n$ , the probability is at most  $\exp(-\Omega(n))$ .  $\blacksquare$

To construct local distributions for the Sherali-Adams hierarchy, we will consider (a slight modification) the hypergraph  $H$  corresponding to the instance  $\Phi$ . We first show that distributions on edges of this hypergraph can be consistently propagated in a tree, provided they agree on intersecting vertices.

For a set  $U \subseteq V(H)$  in a hypergraph  $H$ , recall that  $\text{cl}(U)$  includes all paths of lengths at most 1 between any two vertices in  $U$ . Thus,  $E(\text{cl}(U)) = \{e \in E \mid |e \cap U| \geq 2\}$ . Note that **Lemma 5.2** implies that edges forming a tree in  $H$  satisfy the hypothesis of **Lemma 5.3** below.

**Lemma 5.3** Let  $H = (V, E)$  be a  $k$ -uniform hypergraph. Let  $U \subseteq V$  and let the set of edges  $E(\text{cl}(U))$  form a tree. For each  $e \in E(\text{cl}(U))$ , let  $\overline{\mathcal{D}}_e$  be a distribution on  $[q]^e$  such that for any  $u \in U$  and  $e_1, e_2 \in E(\text{cl}(U))$  such that  $e_1 \cap e_2 = \{u\}$ , we have  $\overline{\mathcal{D}}_{e_1|u} = \overline{\mathcal{D}}_{e_2|u} = \overline{\mathcal{D}}_u$ . Then,

- there exists a distribution  $\bar{\mathcal{D}}_U$  on  $[q]^U$  such that  $\bar{\mathcal{D}}_{U|e \cap U} = \bar{\mathcal{D}}_{e|e \cap U}$  for all  $e \in E(U)$ .
- If  $U' \subseteq U$  is such that the edges in  $E(\text{cl}(U'))$  form a subtree of  $E(\text{cl}(U))$ , then  $\bar{\mathcal{D}}_{U|U'} = \bar{\mathcal{D}}_{U'}$ .

**Proof:** We define the distribution by starting with an arbitrary edge and traversing the tree in an arbitrary order. Let  $e_1, \dots, e_r$  be a traversal of the edges in  $E(\text{cl}(U))$  such that for all  $i$ ,  $|(\cup_{j < i} e_j) \cap e_i| = 1$ . Let  $U_0 = \cup_{j < i} e_j$  be the set of vertices for which we have already sampled an assignment and let  $e_i$  be the next edge in the traversal, with  $u$  being the unique vertex in  $e_i \cap U_0$ . We sample an assignment to the vertices in  $e$ , conditioned on the value for the vertex  $u$ . Formally, we extend the distribution  $\bar{\mathcal{D}}_{U_0}$  to  $U_0 \cup e$  by taking, for any  $\alpha \in [q]^{U_0 \cup e}$

$$\bar{\mathcal{D}}_{U_0 \cup e}(\alpha) = \bar{\mathcal{D}}_{U_0}(\alpha(U_0)) \cdot \frac{\bar{\mathcal{D}}_e(\alpha(e))}{\bar{\mathcal{D}}_{e|u}(\alpha(u))} = \bar{\mathcal{D}}_{U_0}(\alpha(U_0)) \cdot \frac{\bar{\mathcal{D}}_e(\alpha(e))}{\bar{\mathcal{D}}_u(\alpha(u))}.$$

The above process defines a distribution  $\bar{\mathcal{D}}_{\text{cl}(U)}$  on  $\text{cl}(U)$ , with

$$\bar{\mathcal{D}}_{\text{cl}(U)}(\alpha) = \frac{\prod_{e \in E(U)} \bar{\mathcal{D}}_e(\alpha(e))}{\prod_{u \in \text{cl}(U)} (\bar{\mathcal{D}}_u(\alpha(u)))^{\deg(u)-1}}.$$

In the above expression, we use  $\deg(u)$  to denote the degree of vertex  $u$  in tree formed by the edges in  $E(\text{cl}(U))$  i.e.,  $\deg(u) = |\{e \in E(\text{cl}(U)) \mid u \in e\}|$ . We then define the distribution  $\bar{\mathcal{D}}_U$  as the marginalized distribution  $\bar{\mathcal{D}}_{\text{cl}(U)|U}$  i.e.,

$$\bar{\mathcal{D}}_U(\alpha) = \sum_{\substack{\beta \in [q]^{\text{cl}(U)} \\ \beta(U) = \alpha}} \bar{\mathcal{D}}_{\text{cl}(U)}(\beta).$$

Note that the distribution  $\bar{\mathcal{D}}_{\text{cl}(U)}$  and hence also the distribution  $\bar{\mathcal{D}}_U$  are independent of the order in which we traverse the edges in  $E(\text{cl}(U))$ . Also, since the above process samples each edge according to the distribution  $\bar{\mathcal{D}}_e$ , we have that for any  $e \in E(U)$ ,  $\bar{\mathcal{D}}_{\text{cl}(U)|e} = \bar{\mathcal{D}}_e$ . Thus, also for any  $e \in E(U)$ ,  $\bar{\mathcal{D}}_{U|e \cap U} = \bar{\mathcal{D}}_{e|e \cap U}$ .

Let  $U' \subseteq U$  be any set such that  $E(\text{cl}(U'))$  forms a subtree of  $E(\text{cl}(U))$ . Then there exists a traversal  $e_1, \dots, e_r$ , and  $i \in [r]$  such that  $e_j \in E(\text{cl}(U')) \forall j \leq i$  and  $e_j \notin E(\text{cl}(U')) \forall j > i$ . However, the distribution defined by the partial traversal  $e_1, \dots, e_i$  is precisely  $\bar{\mathcal{D}}_{\text{cl}(U')}$ . Thus, we get that  $\bar{\mathcal{D}}_{\text{cl}(U)|\text{cl}(U')} = \bar{\mathcal{D}}_{\text{cl}(U')}$  which implies  $\bar{\mathcal{D}}_{U|U'} = \bar{\mathcal{D}}_{U'}$ . ■

We can now prove the completeness for our construction using consistent decompositions.

**Lemma 5.4** *Let  $\varepsilon > 0$  and let  $\Phi$  be a random instance of MAX  $k$ -CSP $_q$  ( $f$ ) generated by choosing  $\gamma \cdot n$  constraints independently at random as above. Then, there is a  $t = \Omega_{\varepsilon, k, n_0} \left( \frac{\log n}{\log \log n} \right)$ , such that with probability  $1 - \varepsilon$  over the choice of  $\Phi$ , there exist distributions  $\{\mathcal{D}_S\}_{|S| \leq t}$  satisfying:*

- For all  $S \subseteq V$  with  $|S| \leq t$ ,  $\mathcal{D}_S$  is a distribution on  $[q]^S$ .
- For all  $T \subseteq S \subseteq V$  with  $|S| \leq t$ ,  $\mathcal{D}_{S|T} = \mathcal{D}_T$ .

- The distributions satisfy

$$\mathbb{E}_{C \in \Phi} \mathbb{E}_{\alpha_C \sim \mathcal{D}_{S_C}} [f(\alpha_C + b_C)] \geq c - \varepsilon.$$

**Proof:** By [Theorem 3.3](#), we know that there exists  $\delta$  such that with probability  $1 - \varepsilon/4$ , after removing a set of constraints  $C_B$  of size at most  $(\varepsilon/4) \cdot m$ , we can assume that the remaining instance has girth at least  $g = \delta \cdot \log n$ . Also, there exists  $\theta, c > 0$  such that for all  $t \leq n^\theta$ , the metric  $\rho_\mu^H$  restricted to any set  $S$  of size at most  $t$  embeds isometrically into the unit sphere in  $\ell_2$ , for all  $\mu \geq c \cdot \frac{\log t + \log \log n}{\log n}$ .

We choose  $\mu = 2c \cdot \frac{\log \log n}{\log n}$  and  $t = \frac{\varepsilon^2}{400k^2} \cdot \frac{1}{\mu}$  so that

$$\mu \geq c \cdot \frac{\log t + \log \log n}{\log n} \quad \text{and} \quad \sqrt{\mu \cdot t} \leq \frac{\varepsilon}{20k}.$$

Thus, by [Lemma 4.3](#),  $H$  admits an  $(\varepsilon/2)$ -sparse partitioning scheme of order  $t$  with each cluster in the partition having diameter at most  $\Delta_H = O(1/\mu)$ . Let  $\{\mathcal{P}_S\}_{|S| \leq t}$  denote this partitioning scheme.

Given a set  $S$ , the distribution  $\mathcal{D}_S$  is a convex combination of several distributions  $\mathcal{D}_{S,P}$ , corresponding to different partitions  $P$  sampled from  $\mathcal{P}_S$ . We describe the distribution  $\mathcal{D}_S$  by giving the procedure to sample an  $\alpha \in [q]^S$ . Given the set  $S$  with  $|S| \leq t$ :

- Sample a partition  $P = (U_1, \dots, U_r)$  from the distribution  $\mathcal{P}_S$ .
- For each set  $U_i$ , consider the set  $\mathcal{C}(U_i)$  obtained by including the vertices contained in all the edges in the shortest path between all  $u, v \in U_i$ . Note that since  $U_i$  has diameter at most  $\Delta_H$  in  $H$ ,  $\mathcal{C}(U_i)$  is connected and in fact  $\mathcal{C}(U_i) = \text{cl}_{\Delta_H}(U_i)$ . Also, since the each vertex in an included path is within distance at most  $\Delta_H/2$  of an endpoint, and  $U_i$  has diameter at most  $\Delta_H$ , we know that the diameter of  $\mathcal{C}(U_i)$  is at most  $2 \cdot \Delta_H$ . Hence,  $\mathcal{C}(U_i)$  is a tree. Finally, we must have  $\text{cl}(\mathcal{C}(U_i)) = \mathcal{C}(U_i)$  since any additional path of length 1 would create a cycle of length at most  $2 \cdot \Delta_H + 1$ .

Thus, by [Lemma 5.2](#) and [Lemma 5.3](#) (with probability at least  $1 - \varepsilon/4$ ) there exists a distribution  $\overline{\mathcal{D}}_{\mathcal{C}(U_i)}$  for each  $U_i$ , satisfying  $\overline{\mathcal{D}}_{\mathcal{C}(U_i)|e} = \overline{\mathcal{D}}_e$  for all  $e \in E(\mathcal{C}(U_i))$ . Here,  $\overline{\mathcal{D}}_e$  are the distributions given by [Lemma 5.2](#), which form a solution to the basic LP for  $\Phi$ , with value at least  $c - \varepsilon/4$ . For each  $U_i$ , define the distribution

$$\mathcal{D}'_{U_i} := \overline{\mathcal{D}}_{\mathcal{C}(U_i)|U_i}.$$

- Sample  $\alpha \in [q]^S$  according to the distribution

$$\mathcal{D}_{S,P} := \mathcal{D}'_{U_1} \times \dots \times \mathcal{D}'_{U_r}.$$

Thus, we have

$$\mathcal{D}_S := \mathbb{E}_{P=(U_1, \dots, U_r) \sim \mathcal{P}_S} \left[ \prod_{i=1}^r \mathcal{D}'_{U_i} \right],$$

where the distributions  $\mathcal{D}'_{U_i}$  are defined as above.

We first prove the distributions are consistent on intersections i.e.,  $\mathcal{D}_{S|T} = \mathcal{D}_T$  for any  $T \subseteq S$ . Note that by [Lemma 4.3](#), the distributions  $\mathcal{P}_S$  and  $\mathcal{P}_T$  satisfy  $\mathcal{P}_{S|T} = \mathcal{P}_T$ . Each partition  $(U_1, \dots, U_r)$  also produces a partition  $T$ . For ease of notation, we assume that the first (say)  $r'$  clusters have non-empty intersection with  $S$ . Let  $V_i = U_i \cap T$  for  $1 \leq i \leq r'$  ( $V_i = \emptyset$  for  $i > r'$ ). Then, we have

$$\begin{aligned} \mathcal{D}_{S|T} &= \mathbb{E}_{P=(U_1, \dots, U_r) \sim \mathcal{P}_S} \left[ \prod_{i=1}^r \mathcal{D}'_{U_i|V_i} \right] = \mathbb{E}_{P=(U_1, \dots, U_r) \sim \mathcal{P}_S} \left[ \prod_{i=1}^{r'} \overline{\mathcal{D}}_{\mathcal{C}(U_i)|V_i} \right] \\ &= \mathbb{E}_{P=(U_1, \dots, U_r) \sim \mathcal{P}_S} \left[ \prod_{i=1}^{r'} \overline{\mathcal{D}}_{\mathcal{C}(V_i)|V_i} \right] \\ &= \mathbb{E}_{P'=(V_1, \dots, V_{r'}) \sim \mathcal{P}_T} \left[ \prod_{i=1}^{r'} \overline{\mathcal{D}}_{\mathcal{C}(V_i)|V_i} \right] \end{aligned}$$

The second to last equality above uses the fact that  $\mathcal{C}(V_i)$  is a subtree of  $\mathcal{C}(U_i)$  and thus  $\overline{\mathcal{D}}_{\mathcal{C}(U_i)|\mathcal{C}(V_i)} = \overline{\mathcal{D}}_{\mathcal{C}(V_i)}$  by [Lemma 5.3](#). The last equality uses the fact that  $\mathcal{P}_{S|T} = \mathcal{P}_T$  by [Lemma 4.3](#).

We now argue that the LP solution corresponding to the above distributions  $\{\mathcal{D}_S\}_{|S| \leq t}$  has value at least  $c - \varepsilon$ . Recall that the value of the LP solution is given by

$$\mathbb{E}_{C \in \Phi} \mathbb{E}_{\alpha \sim \mathcal{D}_{S_C}} [f(\alpha + b_C)].$$

Consider any constraint  $C$  in  $\Phi$ , with the corresponding set of variables  $S_C$  and the corresponding hyperedge  $e$ . When defining the distribution  $\mathcal{D}_{S_C}$ , we will partition  $S_C$  according to the distribution  $\mathcal{P}_{S_C}$ . By [Lemma 4.3](#) and our choice of parameters

$$\mathbb{P}_{P \sim \mathcal{P}_{S_C}} [P \neq \{S_C\}] \leq 10k \cdot \sqrt{\mu \cdot t} \leq \frac{\varepsilon}{2}.$$

For a constraint set which is not in the deleted set  $C_B$ , if the edge  $e$  corresponding to the constraint  $C$  is not split by a partition  $P$  sampled according to  $\mathcal{P}_{S_C}$ , then by [Lemma 5.3](#)  $\mathcal{D}_{S_C, P} = \overline{\mathcal{D}}_{S_C}$ . Here,  $\overline{\mathcal{D}}_{S_C}$  is the distribution given by [Lemma 5.2](#). Since  $f$  is Boolean, we have that for  $C \notin C_B$ ,

$$\mathbb{E}_{\alpha \sim \mathcal{D}_{S_C}} [f(\alpha + b_C)] \geq \mathbb{E}_{\alpha \sim \overline{\mathcal{D}}_{S_C}} [f(\alpha + b_C)] - \frac{\varepsilon}{2}.$$

Using [Lemma 5.2](#) again, we get

$$\begin{aligned} \mathbb{E}_{C \sim \Phi} \mathbb{E}_{\alpha \sim \mathcal{D}_{S_C}} [f(\alpha + b_C)] &\geq \mathbb{E}_{C \sim \Phi} \left[ (1 - \mathbb{1}_{\{C \in C_B\}}) \cdot \left( \mathbb{E}_{\alpha \sim \overline{\mathcal{D}}_{S_C}} [f(\alpha + b_C)] - \frac{\varepsilon}{2} \right) \right] \\ &\geq \mathbb{E}_{C \sim \Phi} \mathbb{E}_{\alpha \sim \overline{\mathcal{D}}_{S_C}} [f(\alpha + b_C)] - \frac{\varepsilon}{2} - \mathbb{E}_{C \sim \Phi} [\mathbb{1}_{\{C \in C_B\}}] \\ &\geq c - \frac{\varepsilon}{4} - \frac{\varepsilon}{2} - \frac{\varepsilon}{4} \\ &\geq c - \varepsilon, \end{aligned}$$

where the penultimate inequality uses the fact that the fraction of constraints in the initially deleted set  $C_B$  is at most  $\varepsilon/4$  (for large enough  $n$ ).  $\blacksquare$

## 5.2 Integrality Gaps for resistant predicates

Let  $f : \{0,1\}^k \rightarrow \{0,1\}$  be a boolean predicate and let  $\rho(f) = \frac{f^{-1}(1)}{2^k}$  be the fractions of satisfying assignments to  $f$ . Then  $f$  is approximation resistant if it is hard to distinguish the MAX-CSP instances on  $f$  between which are at least  $1 - o(1)$  satisfiable vs which are at most  $\rho(f) + o(1)$  satisfiable.

In [15] the authors introduce the notion of vanishing measure (on a polytope defined by  $f$ ) and use it to characterize a variant of approximation resistance, called strong approximation resistance, assuming the Unique Games conjecture. They also show gave a *weaker* notion of vanishing measures, which they used to characterize strong approximation resistance for LP hierarchies. In particular, they proved that when the condition in their characterization is satisfied, there exists a  $(1 - o(1), \rho(f) + o(1))$  integrality gap for  $O(\log \log n)$  levels of Sherali-Adams hierarchy for predicates  $f$ . Here, we show that using [Theorem 1.1](#), their result can be simplified and strengthened <sup>2</sup> to  $O\left(\frac{\log n}{\log \log n}\right)$  levels.

Let us first recall some useful notation defined by Khot et al. [15] before we define the notion of vanishing measure:

**Definition 5.5** For a predicate  $f : \{0,1\}^k \rightarrow \{0,1\}$ , let  $\mathcal{C}(f)$  be the convex polytope of first moments (biases) of distributions supported on satisfying assignments of  $f$  i.e.,

$$\mathcal{C}(f) := \left\{ \zeta \in \mathbb{R}^k \mid \forall i \in [k], \zeta_i = \mathbb{E}_{\alpha \sim \nu} [(-1)^{\alpha_i}], \text{ Supp}(\nu) \subseteq f^{-1}(1) \right\}.$$

For a measure  $\Lambda$  on  $\mathcal{C}(f)$ ,  $S \subseteq [k]$ ,  $b \in \{0,1\}^S$  and permutation  $\pi : S \rightarrow S$ , let  $\Lambda_{S,\pi,b}$  denote the induced measure on  $\mathbb{R}^S$  by considering vectors with coordinates  $\left\{ (-1)^{b_{\pi(i)}} \cdot \zeta_{\pi(i)} \right\}_{i \in S}$ , where  $\zeta \sim \Lambda$ .

We recall below the definition of vanishing measure for LPs from [15] (see Definition 1.3) :

**Definition 5.6** A measure  $\Lambda$  on  $\mathcal{C}(f)$  is called vanishing (for LPs) if for every  $1 \leq t \leq k$ , the following signed measure

$$\mathbb{E}_{|S|=t} \mathbb{E}_{\pi: S \rightarrow S} \mathbb{E}_{b \in \{0,1\}^t} \left[ \left( \prod_{i=1}^t (-1)^{b_i} \right) \cdot \hat{f}(S) \cdot \Lambda_{S,\pi,b} \right]$$

is identically 0. We say  $f$  has a vanishing measure if there exists a vanishing measure  $\Lambda$  on  $\mathcal{C}(f)$ .

In particular, they prove the following theorem:

**Theorem 5.7** Let  $f : \{0,1\}^k \rightarrow \{0,1\}$  be a  $k$ -ary boolean predicate that has a vanishing measure. Then for every  $\varepsilon > 0$ , there is a constant  $c_\varepsilon > 0$  such that for infinitely many  $N \in \mathbb{N}$ , there exists an instance  $\Phi$  of MAX  $k$ -CSP( $f$ ) on  $N$  variables satisfying the following:

$$- \text{OPT}(\Phi) \leq \rho(f) + \varepsilon.$$

<sup>2</sup>The LP integrality gap result of Khot et al. is in fact slightly stronger than stated above. They show that LP value is at least  $1 - o(1)$  while there is no integer solution achieving a value outside the range  $[\rho(f) - o(1), \rho(f) + o(1)]$ . It is easy to see that the same also holds for the instance constructed here.

- The optimum for the LP relaxation given by  $c_\varepsilon \cdot \log \log N$  levels of Sherali-Adams hierarchy has  $\text{FRAC}(\Phi) \geq 1 - O(k \cdot \sqrt{\varepsilon})$ .

Combining this with our [Theorem 1.1](#) already gives us the following stronger result:

**Corollary 5.8** *Let  $f : \{0, 1\}^k \rightarrow \{0, 1\}$  be a  $k$ -ary boolean predicate that has a vanishing measure. Then for every  $\varepsilon > 0$ , there is a constant  $c_\varepsilon > 0$  such that for infinitely many  $N \in \mathbb{N}$ , there exists an instance  $\Phi$  of MAX  $k$ -CSP( $f$ ) on  $N$  variables satisfying the following:*

- All integral assignment of  $\Phi$  satisfies at most  $\rho(f) + \varepsilon$  fraction of constraints.
- The LP relaxation given by  $c_\varepsilon \cdot \frac{\log N}{\log \log N}$  levels of Sherali-Adams hierarchy has  $\text{FRAC}(\Phi) \geq 1 - O(k\sqrt{\varepsilon})$ .

However, note that to apply [Theorem 1.1](#), one only needs a gap for the basic LP, which is much weaker requirement than the  $O(\log \log N)$ -level gap given by [Theorem 5.7](#). We observe below that the gap for the basic LP follows very simply from the construction by Khot et al. [15]. One can then directly use this gap for applying [Theorem 1.1](#) instead of going through [Theorem 5.7](#).

Khot et al. [15] use the probabilistic construction given in [Fig. 4](#), for a given  $\varepsilon > 0$ . The construction actually requires  $\Lambda$  to be a vanishing measure over the polytope  $\mathcal{C}_\delta(f) := (1 - \delta) \cdot \mathcal{C}(f)$ , for  $\delta = \sqrt{\varepsilon}$ . However, since  $\mathcal{C}_\delta(f)$  is simply a scaling of  $\mathcal{C}(f)$ , a vanishing measure over  $\mathcal{C}(f)$  also gives a vanishing measure over  $\mathcal{C}_\delta(f)$ . Note that each  $\zeta_0 \in \mathcal{C}(f)$  corresponds to a distribution  $\nu_0$  supported in  $f^{-1}(1)$ . For each  $\zeta \in \mathcal{C}_\delta$ , let  $\zeta_0 = \frac{1}{1-\delta} \cdot \zeta$  be the point in  $\mathcal{C}(f)$  with distribution  $\nu_0$ . Then the distribution  $\nu = (1 - \delta) \cdot \nu_0 + \delta \cdot U_k$  (where  $U_k$  denotes the uniform distribution on  $\{0, 1\}^k$ ) satisfies  $\forall i \in [k] \mathbb{E}_{\alpha \sim \nu} [(-1)^{\alpha_i}] = \zeta_i$ .

Let  $n_0 = \lceil \frac{1}{\varepsilon} \rceil$ . Partition the interval  $[0, 1]$  into  $n_0 + 1$  disjoint intervals  $I_0, I_1, \dots, I_{n_0}$  where  $I_0 = \{0\}$  and  $I_i = (i^{-1}/n_0, i/n_0]$  for  $1 \leq i \leq n_0$ . For each interval  $I_i$ , let  $X_i$  be a collection of  $n$  variables (disjoint from all  $X_j$  for  $j \neq i$ ).

Generate  $m$  constraints independently according to the following procedure:

- Sample  $\zeta \sim \Lambda$ .
- For each  $j \in [k]$ , let  $i_j$  be the index of the interval which contains  $|\zeta(j)|$ . Sample uniformly a variable  $y_j$  from the set  $X_{i_j}$ .
- If  $\zeta(j) < 0$ , then negate  $y_j$ . If  $\zeta(j) = 0$ , then negate  $y_j$  w.p.  $\frac{1}{2}$ .
- Introduce the constraint  $f$  on the sampled  $k$  tuple of literals.

Figure 4: Sherali-Adams integrality gap instance for vanishing measure

They show for a sufficiently large constant  $\gamma$ , an instance  $\Phi$  with  $m = \gamma \cdot n$  constraints satisfies with high probability, that for all assignments  $\sigma$ ,  $|\text{sat}_\Phi(\sigma) - \rho(f)| \leq \varepsilon$  (see Lemma 4.4 in [15]). The proof is similar to that of [Lemma 5.1](#).

Additionally, we need the following claim from [15] (see Claim 4.7 there), which allows one to “round” coordinates of the vectors  $\zeta \in \mathcal{C}_\delta(f)$  to the end-points of the intervals

$I_0, \dots, I_{n_0}$ . This ensures that any two variables in the same collection  $X_i$  have the same bias. The proof of the claim follows simply from a hybrid argument. We include it in the appendix for completeness.

**Claim 5.9** *Let  $\zeta \in \mathcal{C}_\delta(f)$  and let  $\nu$  be the corresponding distribution supported in  $f^{-1}(1)$  such that for all  $i \in [k]$ , we have  $\zeta_i = \mathbb{E}_{\alpha \sim \nu} [(-1)^{\alpha_i}]$ . Let  $t_1, \dots, t_k \in [0, 1]$  be such that for all  $i \in [k]$ ,  $|t_i - |\zeta_i|| \leq \varepsilon$  for  $\varepsilon < \delta/2$ . Then there exists a distribution  $\nu'$  on  $\{0, 1\}^k$  such that*

$$\|\nu - \nu'\|_1 = O(k \cdot (\varepsilon/\delta)) \quad \text{and} \quad \forall i \in [k], \quad \mathbb{E}_{\alpha \sim \nu'} [(-1)^{\alpha_i}] = \text{sign}(\zeta_i) \cdot t_i.$$

We can now use the above to give a simplified proof of [Corollary 5.8](#).

**Proof of Corollary 5.8:** Here we exhibit a solution of the basic LP [Fig. 2](#) for the instance given in [Fig. 4](#). For each variable  $y_j$  coming from the set  $X_j$  for  $j \in \{0, 1, \dots, n_0\}$ , we set the bias  $t_j$  of the variable to be the rightmost point of the interval  $I_j$  i.e., set  $x_{(y_j, -1)} = \frac{1}{2} \cdot \left(1 - \frac{i}{n_0}\right)$  and  $x_{(y_j, 1)} = \frac{1}{2} \cdot \left(1 + \frac{i}{n_0}\right)$ .

For each constraint  $C$  of the form  $f(y_{i_1} + b_1, \dots, y_{i_k} + b_k)$ , let  $\zeta(C) \in \mathcal{C}_\delta(f)$  be the point used to generate it, and let  $\nu(C)$  denote the corresponding distribution on  $\{0, 1\}^k$ . By [Claim 5.9](#), there exists a distribution  $\nu'(C)$  such that  $\|\nu(C) - \nu'(C)\|_1 = O(k\varepsilon/\delta)$  and such that the biases of the *literals* satisfy  $\mathbb{E}_{\alpha \sim \nu'(C)} [(-1)^{\alpha_i}] = \text{sign}(\zeta_j) \cdot t_{i_j}$ , where  $t_{i_j}$  denotes the bias for the interval to which  $y_{i_j}$  belongs. When  $t_{i_j} \neq 0$ , we negate a variable only when  $\text{sign}(\zeta_j) < 0$ . Thus, we have  $\mathbb{E}_{\alpha \sim \nu'(C)} [(-1)^{\alpha_j + b_j}] = t_{i_j}$ , which is consistent with the bias given by the singleton variables  $x_{(y_{i_j}, 1)}$  and  $x_{(y_{i_j}, -1)}$ . We thus define the local distribution on the set  $S_C$  as  $\mathcal{D}_{S_C}(\alpha) = (\nu'(C))(\alpha + b_C)$ .

For all  $C \in \Phi$ , since  $\zeta(C) \in \mathcal{C}_\delta(f)$ , we have that  $\mathbb{E}_{\alpha \sim \nu(C)} [f(\alpha)] \geq 1 - \delta$ . Also, since  $\|\nu(C) - \nu'(C)\|_1 = O(k\varepsilon/\delta)$ , we get that  $\mathbb{E}_{\alpha \sim \nu'(C)} [f(\alpha)] \geq 1 - \delta - O(k\varepsilon/\delta)$ . Thus, we have for all  $C \in \Phi$ ,  $\mathbb{E}_{\alpha \sim \mathcal{D}_{S_C}} [f(\alpha + b_C)] \geq 1 - \delta - O(k\varepsilon/\delta)$ . Taking  $\delta = \sqrt{\varepsilon}$  proves the claim. ■

### 5.3 Lower bounds for LP extended formulations

A connection between LP integrality gaps for the Sheral-Adams hierarchy, and lower bounds on the size of LP extended formulations, was established by Chan et al. [\[7\]](#). They proved the following:

**Theorem 5.10 ([\[7\]](#))** *Let  $k, q \in \mathbb{N}$  and  $f : [q]^k \rightarrow \{0, 1\}$  be given. Let  $r : \mathbb{N} \rightarrow \mathbb{N}$  be a function such that the relaxation obtained by  $r(n)$  levels of the Sherali-Adams hierarchy cannot achieve a  $(c, s)$  approximation for instances of MAX  $k$ -CSP $_q(f)$  on  $n$  variables. Then, for all large enough  $n$ , no LP extended formulation of size  $n^{(r(n))^2}$  can achieve a  $(c, s)$  approximation on instances of size  $N$ , where  $N \leq n^{10 \cdot r(n)}$*

Combining the above with [Theorem 1.1](#) and taking  $r(n) = \Omega\left(\frac{\log n}{\log \log n}\right)$  yields [Corollary 1.2](#).

**Corollary 1.2** *Let  $f : [q] \rightarrow \{0, 1\}$  be any predicate. Let  $\Phi_0$  be a  $(c, s)$  integrality gap instance for basic LP relaxation of MAX  $k$ -CSP $(f)$ . Then for every  $\varepsilon > 0$ , there exists  $c_\varepsilon > 0$  such that for infinitely  $N \in \mathbb{N}$ , there exist  $(c - \varepsilon, s + \varepsilon)$  integrality gap instances of size  $N$ , for every linear extended formulation of size at most  $\exp\left(c_\varepsilon \cdot \frac{(\log N)^{3/2}}{(\log \log N)^{1/2}}\right)$ .*



## Acknowledgements

We thank Chandra Chekuri, Subhash Khot and Yury Makarychev for helpful discussions. This research was supported by supported by the National Science Foundation under award number CCF-1254044.

## References

- [1] Michael Alekhnovich, Sanjeev Arora, and Iannis Tourlakis. Towards strong nonapproximability results in the Lovasz-Schrijver hierarchy. In *Proceedings of the 37th ACM Symposium on Theory of Computing*, pages 294–303, 2005. [2](#), [11](#)
- [2] Sanjeev Arora, Béla Bollobás, and László Lovász. Proving integrality gaps without knowing the linear program. In *Proceedings of the 43rd IEEE Symposium on Foundations of Computer Science*, pages 313–322. IEEE, 2002. [2](#)
- [3] Sanjeev Arora, Béla Bollobás, László Lovász, and Iannis Tourlakis. Proving integrality gaps without knowing the linear program. *Theory of Computing*, 2(2):19–51, 2006. [2](#), [9](#), [11](#), [12](#)
- [4] Boaz Barak, Siu On Chan, and Pravesh K. Kothari. Sum of squares lower bounds from pairwise independence. In *Proceedings of the 47th ACM Symposium on Theory of Computing*, pages 97–106, New York, NY, USA, 2015. ACM. [2](#), [8](#), [11](#)
- [5] Eli Ben-Sasson. *Expansion in Proof Complexity*. PhD thesis, Hebrew University, 2001. [11](#)
- [6] Siavosh Benabbas, Konstantinos Georgiou, Avner Magen, and Madhur Tulsiani. SDP Gaps from Pairwise Independence. *Theory of Computing*, 8(1):269–289, 2012. [2](#), [11](#)
- [7] Siu On Chan, James Lee, Prasad Raghavendra, and David Steurer. Approximate constraint satisfaction requires large LP relaxations. In *Proceedings of the 54th IEEE Symposium on Foundations of Computer Science*, pages 350–359. IEEE, 2013. [2](#), [3](#), [24](#)
- [8] Moses Charikar, Chandra Chekuri, Ashish Goel, Sudipto Guha, and Serge Plotkin. Approximating a finite metric by a small number of tree metrics. In *Proceedings of the 39th IEEE Symposium on Foundations of Computer Science*, pages 379–388, 1998. [5](#), [15](#)
- [9] Moses Charikar, Konstantin Makarychev, and Yury Makarychev. Local global trade-offs in metric embeddings. In *Proceedings of the 48th IEEE Symposium on Foundations of Computer Science*, 2007. [5](#), [9](#), [11](#), [12](#)
- [10] Moses Charikar, Konstantin Makarychev, and Yury Makarychev. Near-optimal algorithms for maximum constraint satisfaction problems. In *Proceedings of the 18th ACM-SIAM Symposium on Discrete Algorithms*, pages 62–68, 2007. [3](#)
- [11] Moses Charikar, Konstantin Makarychev, and Yury Makarychev. Integrality gaps for Sherali-Adams relaxations. In *Proceedings of the 41st ACM Symposium on Theory of Computing*, 2009. [2](#), [8](#), [9](#), [13](#)

- [12] Wenceslas Fernandez de la Vega and Claire Kenyon-Mathieu. Linear programming relaxations of maxcut. In *Proceedings of the 18th ACM-SIAM Symposium on Discrete Algorithms*, pages 53–61, 2007. 2, 5
- [13] Subhash Khot. On the power of unique 2-prover 1-round games. In *Proceedings of the 34th ACM Symposium on Theory of Computing*, pages 767–775, 2002. 2
- [14] Subhash Khot and Rishi Saket. SDP integrality gaps with local  $\ell_1$ -embeddability. In *Proceedings of the 50th IEEE Symposium on Foundations of Computer Science*, 2009. 2
- [15] Subhash Khot, Madhur Tulsiani, and Pratik Worah. A characterization of strong approximation resistance. In *Proceedings of the 46th ACM Symposium on Theory of Computing*, pages 634–643. ACM, 2014. 3, 4, 5, 16, 22, 23
- [16] Vladimir Kolmogorov, Johan Thapper, and Stanislav Živný. The power of linear programming for general-valued CSPs. *SIAM Journal on Computing*, 44(1):1–36, 2015. 3
- [17] Pravesh Kothari, Raghu Meka, and Prasad Raghavendra. Personal Communication, 2016. 2, 3
- [18] Robert Krauthgamer, James R Lee, Manor Mendel, and Assaf Naor. Measured descent: A new embedding method for finite metrics. *Geometric & Functional Analysis GAFA*, 15(4):839–858, 2005. 5
- [19] L. Lovász and A. Schrijver. Cones of matrices and set-functions and 0-1 optimization. *SIAM J. on Optimization*, 1(12):166–190, 1991. 2
- [20] Prasad Raghavendra. Optimal algorithms and inapproximability results for every CSP? In *STOC*, pages 245–254, 2008. 2, 4
- [21] Prasad Raghavendra and David Steurer. Integrality gaps for strong SDP relaxations of unique games. In *Proceedings of the 50th IEEE Symposium on Foundations of Computer Science*, 2009. 2
- [22] Grant Schoenebeck. Linear level Lasserre lower bounds for certain k-CSPs. In *Proceedings of the 49th IEEE Symposium on Foundations of Computer Science*, 2008. 2
- [23] Hanif D. Sherali and Warren P. Adams. A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM J. Discrete Math.*, 3(3):411–430, 1990. 2
- [24] Johan Håstad. On the Efficient Approximability of Constraint Satisfaction Problems. In *Surveys in Combinatorics*, volume 346, pages 201–222. Cambridge University Press, 2007. 1
- [25] Johan Thapper and Stanislav Živný. The complexity of finite-valued CSPs. In *Proceedings of the forty-fifth annual ACM symposium on Theory of computing*, pages 695–704. ACM, 2013. 3
- [26] Johan Thapper and Stanislav Živný. The power of Sherali-Adams relaxations for general-valued CSPs. *arXiv preprint arXiv:1606.02577*, 2016. 3
- [27] Madhur Tulsiani. CSP gaps and reductions in the Lasserre hierarchy. In *Proceedings of the 41st ACM Symposium on Theory of Computing*, 2009. 2

## A Omitted proofs from Section 3 and Section 5

**Lemma 3.8** *Let  $\eta < 1/4$  and  $m = \gamma \cdot n$  for  $\gamma > 1$ . Then for  $\tau \leq \frac{1}{n_0} \cdot \left(\frac{1}{e \cdot k^{3k} \cdot \gamma}\right)^{1/\eta}$  the following holds:*

$$\mathbb{P}_{H \sim \mathcal{H}_k(m, n, n_0, \Gamma)} [H \text{ is not } (\tau, \eta)\text{-sparse}] \leq 3 \cdot \left(\frac{k^{3k} \cdot \gamma}{n^{\eta/4}}\right)^{1/k}.$$

**Proof:** As in the proof of Lemma 3.5, given a random hypergraph  $H$ , we construct a hypergraph  $H'$  (by contracting all the vertices in  $[n_0] \times \{j\}$  to  $j \in [n]$ ).

Consider a subset of vertices  $S \subseteq V(H)$  and let  $S' \subseteq V(H')$  be the corresponding contracted set in  $H'$ . Since each edge in  $H$  corresponds to an edge in  $H'$  (counting multiplicities), we have

$$|E(S)| \geq \frac{|S|}{k-1-\eta} \Rightarrow |E(S')| \geq \frac{|S|}{k-1-\eta} \geq \frac{|S'|}{k-1-\eta}.$$

Thus, it suffices to show that  $H'$  is  $(\tau', \eta)$ -sparse for  $\tau' = \tau \cdot n_0$ , since  $|S'| \leq \tau \cdot N = (\tau \cdot n_0) \cdot n$ . Given any multiset in  $[n]^k$ , the probability that it corresponds to an edge in  $H'$  is at most  $(k!) \cdot (m/n^k)$ . Thus, the probability that there exists a set  $T$  of size at most  $\tau' \cdot n$ , containing at least  $|T|/(k-1-\eta)$  edges (counting multiplicities) is at most

$$\sum_{h=1}^{\tau' \cdot n} \binom{n}{h} \cdot \binom{h^k}{r} \cdot \left(\frac{k! \cdot m}{n^k}\right)^r,$$

where  $r = \frac{h}{k-1-\eta}$ . Note that we also need to consider  $h = 1$  as edges in  $H'$  correspond to multisets of size  $k$ , and so may not have all distinct vertices. Simplifying the above using  $\binom{a}{b} \leq \left(\frac{a \cdot e}{b}\right)^b$  and  $k! \leq k^k$  gives

$$\begin{aligned} \sum_{h=1}^{\tau' \cdot n} \binom{n}{h} \cdot \binom{h^k}{r} \cdot \left(\frac{k! \cdot m}{n^k}\right)^r &\leq \sum_{h=1}^{\tau' \cdot n} \left(\frac{n \cdot e}{h}\right)^h \cdot \left(\frac{h^k \cdot e}{r}\right)^r \cdot \left(\frac{k^k \cdot m}{n^k}\right)^r \\ &= \sum_{h=1}^{\tau' \cdot n} \left(e^{k-\eta} \cdot (k-1-\eta) \cdot k^k \cdot \gamma \cdot \left(\frac{h}{n}\right)^\eta\right)^{h/(k-1-\eta)} \\ &\leq \sum_{h=1}^{\tau' \cdot n} \left(k^{3k} \cdot \gamma \cdot \left(\frac{h}{n}\right)^\eta\right)^{h/(k-1-\eta)} \end{aligned}$$

Let  $\theta = \eta/(2k)$ . We divide the above summation in two parts and first consider

$$\begin{aligned} \sum_{h=n^\theta}^{\tau' \cdot n} \left(k^{3k} \cdot \gamma \cdot \left(\frac{h}{n}\right)^\eta\right)^{h/(k-1-\eta)} &\leq \sum_{h=n^\theta}^{\tau' \cdot n} \left(k^{3k} \cdot \gamma \cdot (\tau')^\eta\right)^{n^\theta/(k-1-\eta)} \\ &\leq 2 \cdot \exp\left(-\frac{n^\theta}{k}\right) \\ &\leq 2 \cdot \frac{k}{n^\theta}, \end{aligned}$$

for  $\tau' \leq (e \cdot k^{3k} \cdot \gamma)^{-1/\eta}$ . Considering the first half of the summation, we get

$$\begin{aligned} \sum_{h=1}^{n^\theta} \left( k^{3k} \cdot \gamma \cdot \left( \frac{h}{n} \right)^\eta \right)^{h/(k-1-\eta)} &\leq n^\theta \cdot \left( \frac{k^{3k} \cdot \gamma}{n^{(1-\theta)\eta}} \right)^{1/k} \\ &\leq \left( \frac{k^{3k} \cdot \gamma}{n^{\eta/4}} \right)^{1/k} = k^3 \cdot \gamma^{1/k} \cdot n^{-\theta/2}. \end{aligned}$$

Combining the two bounds gives that the probability is at most  $3k^3 \cdot \gamma^{1/k} \cdot n^{-\theta/2}$ , which equals the desired bound.  $\blacksquare$

**Claim 5.9** Let  $\zeta \in \mathcal{C}_\delta(f)$  and let  $\nu$  be the corresponding distribution supported in  $f^{-1}(1)$  such that for all  $i \in [k]$ , we have  $\zeta_i = \mathbb{E}_{\alpha \sim \nu} [(-1)^{\alpha_i}]$ . Let  $t_1, \dots, t_k \in [0, 1]$  be such that for all  $i \in [k]$ ,  $|t_i - |\zeta_i|| \leq \varepsilon$  for  $\varepsilon < \delta/2$ . Then there exists a distribution  $\nu'$  on  $\{0, 1\}^k$  such that

$$\|\nu - \nu'\|_1 = O(k \cdot (\varepsilon/\delta)) \quad \text{and} \quad \forall i \in [k], \quad \mathbb{E}_{\alpha \sim \nu'} [(-1)^{\alpha_i}] = \text{sign}(\zeta_i) \cdot t_i.$$

**Proof:** Let  $r_j = \text{sign}(\zeta_j) \cdot t_j$  be the desired bias of the  $j^{\text{th}}$  coordinate. Then,  $|\zeta(j) - r_j| \leq \varepsilon$  for all  $j \in [k]$ . We construct a sequence of distributions  $\nu_0, \dots, \nu_k$  such that  $\nu_0 = \nu$  and  $\nu_k = \nu'$ . In  $\bar{\nu}_j$ , the biases are  $(r_1, \dots, r_j, \zeta_{j+1}, \dots, \zeta_k)$ .

The biases in  $\nu_0$  satisfy the above by definition. We obtain  $\bar{\nu}_j$  from  $\bar{\nu}_{j-1}$  as,

$$\nu_j = (1 - \tau_j) \cdot \nu_{j-1} + \tau_j \cdot D_j,$$

where  $D_j$  is the distribution in which all bits, except for the  $j^{\text{th}}$  one, are set independently according to their biases in  $\bar{\nu}_{j-1}$ . For the  $j^{\text{th}}$  bit, we set it to  $\text{sign}(r_j - \zeta_j)$  (if  $r_j - \zeta_j = 0$ , we can simply proceed with  $\bar{\nu}_j = \bar{\nu}_{j-1}$ ). The biases for all except for the  $j^{\text{th}}$  bit are unchanged. For the  $j^{\text{th}}$  bit, the bias now becomes  $r_j$  if

$$r_j = (1 - \tau_j) \cdot \zeta_j + \tau_j \cdot \text{sign}(r_j - \zeta_j) \implies \tau_j \cdot (\text{sign}(r_j - \zeta_j) - r_j) = (1 - \tau_j) \cdot (r_j - \zeta_j).$$

Since  $\zeta \in \mathcal{C}_\delta(f)$ , we know that  $|\text{sign}(r_j - \zeta(j)) - r_j| \geq \delta/2$ . Also,  $|r_j - \zeta(j)| \leq \varepsilon$  by assumption. Thus, we can choose  $\tau_j = O(\varepsilon/\delta)$  which gives that  $\|\bar{\nu}_j - \bar{\nu}_{j-1}\|_1 = O(\varepsilon/\delta)$ . The final bound then follows by triangle inequality.  $\blacksquare$