

# On the impossibility of entropy reversal, and its application to zero-knowledge proofs

Shachar Lovett\*

University of California, San Diego

Jiapeng Zhang<sup>†</sup>

University of California, San Diego

## Abstract

Zero knowledge proof systems have been widely studied in cryptography. In the statistical setting, two classes of proof systems studied are Statistical Zero Knowledge (SZK) and Non-Interactive Statistical Zero Knowledge (NISZK), where the difference is that in NISZK only very limited communication is allowed between the verifier and the prover. It is an open problem whether these two classes are in fact equal. In this paper, we rule out efficient black box reductions between SZK and NISZK.

We achieve this by studying algorithms which can reverse the entropy of a function. The problem of estimating the entropy of a circuit is complete for NISZK. Hence, reversing the entropy of a function is equivalent to a black box reduction of NISZK to its complement, which is known to be equivalent to a black box reduction of SZK to NISZK [Goldreich et al, CRYPTO 1999]. We show that any such algorithm incurs an exponential loss of parameters, and hence cannot be implemented efficiently.

---

\*Email: [slovett@cs.ucsd.edu](mailto:slovett@cs.ucsd.edu). Research supported by NSF CAREER award 1350481 and a Sloan fellowship.

<sup>†</sup>Email: [jpeng.zhang@gmail.com](mailto:jpeng.zhang@gmail.com). Research supported by NSF CAREER award 1350481.

# 1 Introduction

The notion of *Zero-Knowledge Proof Systems* was introduced in the seminal paper of Goldwasser, Micali and Rackoff [GMR89]. Informally, an interactive proof system is a protocol that involves a computational unbounded prover  $P$  and a polynomial time verifier  $V$ . The prover attempts to convince the verifier that an assertion is a YES instance  $x$  of some promise problem.

A promise problem  $\Pi$  consists of two disjoint sets  $\Pi_Y$  and  $\Pi_N$ , e.g., yes instances and no instances. A zero-knowledge proof system for the problem  $\Pi$  requires the following three conditions:

- *Completeness*: If  $x \in \Pi_Y$ , then  $\Pr[(P, V)(x) \text{ accepts}] \geq 2/3$ .
- *Soundness*: If  $x \in \Pi_N$ , then for every adversary  $P^*$ ,  $\Pr[(P^*, V)(x) \text{ accepts}] \leq 1/3$ .
- *Zero-knowledge*: There is a polynomial time simulator  $S$  such that  $S(x)$  and  $(P, V)(x)$  are “indistinguishable”, for every  $x \in \Pi_Y$ .

Different zero knowledge proof systems differ in the allowed communication protocol, and in the notion of indistinguishability applied to the simulator. In this paper, we restrict our attention to statistical proof systems (SZK and NISZK), where the corresponding notion is that of statistical indistinguishability.

**Statistical zero knowledge (SZK).** The complexity class SZK consists all of the problems that has a statistical zero-knowledge proof, where any efficient interactive communication is allowed between the verifier and the prover. Surprisingly, there are complete problems for SZK which has nothing to do with interaction. This was first discovered by Sahai and Vadhan [SV03].

A distribution  $D$  over  $\{0, 1\}^m$  is said to be *efficiently sampleable* if there exists a polynomial size boolean circuit  $C : \{0, 1\}^n \rightarrow \{0, 1\}^m$ , such that the distribution  $D$  can be obtained by applying  $C$  on uniformly chosen bits. By an abuse of notation, we identify  $C$  with this distribution. Given two distributions  $C_1, C_2$  over  $\{0, 1\}^m$ , we denote by  $\text{dist}(C_1, C_2)$  their statistical distance. The following problem, called *Statistical Difference*, was shown by Sahai and Vadhan [SV03] to be complete for SZK.

**Definition 1.1 (Statistical Difference [SV03]).** The promise problem *Statistical Difference*, denoted by  $\text{SD} = (\text{SD}_Y, \text{SD}_N)$ , consists of

- $\text{SD}_Y = \{(C_1, C_2) : \text{dist}(C_1, C_2) \leq 1/3\}$
- $\text{SD}_N = \{(C_1, C_2) : \text{dist}(C_1, C_2) \geq 2/3\}$

Here  $C_1, C_2$  denote polynomial size circuits with the same output length.

**Theorem 1.2 ([SV03]).** *SD is SZK-complete.*

In a follow up work, Goldreich and Vadhan [GV99] gave another SZK-complete problem, called *Entropy Difference*. Below,  $H(C)$  denotes the Shannon entropy of the distribution induced by  $C$ .

**Definition 1.3 (Entropy Difference [GV99]).** The promise problem *Entropy Difference*, denoted by  $\text{ED} = (\text{ED}_Y, \text{ED}_N)$ , consists of

- $\text{ED}_Y = \{(C_1, C_2) : H(C_1) \geq H(C_2) + 1\}$
- $\text{ED}_N = \{(C_1, C_2) : H(C_2) \geq H(C_1) + 1\}$

Here  $C_1, C_2$  denote polynomial size circuits with the same output length.

**Theorem 1.4 ([GV99]).** *The problem ED is SZK-complete.*

**Non-interactive statistical zero knowledge (NISZK).** The notion of *Non-Interactive Zero-Knowledge Proof Systems*, or NISZK was introduced by Blum, Feldman and Micali [BFM88], allows for very restricted communication between the verifier and the prover. Both parties share a common uniformly random string (a random challenge), and the prover sends a single message to the verifier based on this random challenge.

Since the model has been introduced, several problems are proved to be in the class of NISZK. Originally these were problems arising in number theory, such as Quadratic Nonresiduosity and its variants [BFM88; BDMP91; DDP94; DDP97; GMR98]. More recently, this was extended to several natural problems in lattices [PV08].

The problem of finding complete problems for NISZK arose naturally. De Santis et al. [DSD-CPY98], introduced a problem called *Image Density*, and proved that is complete for NISZK. Subsequently, Goldreich, Sahai and Vadhan [GSV99] studied the following two problems and showed that they are too complete for NISZK. Below,

**Definition 1.5 (Statistical Difference from Uniform [GSV99]).** The promise problem *Statistical Difference from Uniform*, denoted by  $\text{SDU} = (\text{SDU}_Y, \text{SDU}_N)$ , consists of

- $\text{SDU}_Y = \{C : \text{dist}(C, \mathcal{U}) \leq 1/n\}$
- $\text{SDU}_N = \{C : \text{dist}(C, \mathcal{U}) \geq 1 - 1/n\}$

Here  $C$  denotes a polynomial size circuit and  $\mathcal{U}$  denotes the uniform distribution on the output space of  $C$ .

**Definition 1.6 (Entropy Approximation [GSV99]).** The promise problem *Entropy Approximation*, denoted by  $\text{EA} = (\text{EA}_Y, \text{EA}_N)$ , consists of

- $\text{EA}_Y = \{(C, k) : H(C) \geq k + 1\}$
- $\text{EA}_N = \{(C, k) : H(C) \leq k - 1\}$

Here  $C$  denotes a polynomial size circuit and  $k \geq 1$ .

**Theorem 1.7 ([GSV99]).** *SDU and EA are NISZK-complete.*

The main open problem that motivated the current paper is *what is the relationship between NISZK and SZK*. Goldreich, Sahai and Vadhan [GSV99] made a significant progress towards resolving this problem.

**Theorem 1.8 ([GSV99]).** *The following statements are equivalent:*

- 1).  $\text{SZK} = \text{NISZK}$ .
- 2). *NISZK is closed under complement.*
- 3). *NISZK is closed under  $NC^1$  truth-table reductions.*
- 4). *ED, or SD Karp-reduces to EA, or SDU respectively.*
- 5). *EA or SDU Karp-reduces to its complement.*

The main goal of the current paper is to show that these statements are all false, at least in a limited model of computation. Concretely, our goal is to rule out *black box reductions* between NISZK and its complement. When we consider black box reductions, the notion of efficient computation disappears, and we replace the study of circuits with the study of arbitrary functions (which can be seen as oracle functions).

## 1.1 Black-box reductions

We describe the notion of black box reductions of functions in this section.

Let  $\mathcal{F}_{n,m}$  denote the family of functions  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$ . A promise problem  $\Pi$  over  $\mathcal{F}_{n,m}$  consists of a family yes instances  $\Pi_Y$  and a family of no instances  $\Pi_N$ , where  $\Pi_Y, \Pi_N \subset \mathcal{F}_{n,m}$  and  $\Pi_Y \cap \Pi_N = \emptyset$ .

**Definition 1.9 (Black-Box Reduction).** Let  $\Pi = (\Pi_Y, \Pi_N)$  and  $\Pi' = (\Pi'_Y, \Pi'_N)$  be promise problems over functions  $\mathcal{F}_{n,m}$  and  $\mathcal{F}_{n',m'}$ , respectively. a *black-box reduction* from  $\Pi$  to  $\Pi'$  is an algorithm  $A^{(\cdot)} : \{0, 1\}^{n'} \rightarrow \{0, 1\}^{m'}$  with oracle access to function  $f \in \mathcal{F}_{n,m}$ , such that the following holds:

- If  $f \in \Pi_Y$  then  $A^f \in \Pi'_Y$ .
- If  $f \in \Pi_N$  then  $A^f \in \Pi'_N$ .

Given an input  $w \in \{0, 1\}^{n'}$ , the algorithm makes a number of queries to  $f$  (the query locations may depend on  $w$  and be adaptive), and outputs a value  $z \in \{0, 1\}^{m'}$ . We define  $A^f(w) = z$ . The query complexity of  $A$ , denoted  $\text{QC}(A)$ , is the maximal number of queries to  $f$  performed over an input.

Our definition of black-box reduction does not relate to decidability, and instead relates to functionality. This type of black-box reduction is well studied in cryptography. Many reductions in the literature are in fact black-box reductions. Examples include the the flattening lemma of [GSV99], the polarization lemma of [SV03], the reduction from Statistical Difference to its complement [SV03], constructions of pseudorandom generators from one-way functions [HILL99; HHR06], constructions of pseudorandom functions from pseudorandom generators [GGM86], and many more.

## 1.2 Our results

We define the function version of the Entropy Approximation (EA) problem.

**Definition 1.10 (Function Entropy Approximation).** The promise problem *Function Entropy Approximation*, denoted by  $\text{FEA} = (\text{FEA}_Y, \text{FEA}_N)$ , consists of

- $\text{FEA}_Y = \{(f, k) : H(f) \geq k + 1\}$
- $\text{FEA}_N = \{(f, k) : H(f) \leq k - 1\}$

Here  $n, m, k \geq 1$  and  $f \in \mathcal{F}_{n,m}$ . Note that the interesting regime of parameters (where the problem is not trivial) is when  $1 \leq k \leq n - 1$ .

A black box reduction from NISZK to its complement needs to map  $\text{FEA}$  to its complement. In particular, an efficient reduction would stay efficient even if we fix  $n, m, k$  to favourable values (we will later set  $m = 3n, k = n - 3$ ). We call such a reduction an *Entropy Reverser*.

**Definition 1.11 (Entropy Reverser).** Let  $n, m, k, n', m', k' \geq 1$ . An  $(n, m, k; n', m', k')$  entropy reverser is a black box reduction  $A$  from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n',m'}$  such that

- If  $H(f) \geq k + 1$  then  $H(A^f) \leq k' - 1$ .
- If  $H(f) \leq k - 1$  then  $H(A^f) \geq k' + 1$ .

Our main result is that entropy reversers require either exponential output length  $n', m'$  or exponential query complexity. In particular, when they are applied to a function  $f$  computed by a polynomial size circuit, their output  $A^f$  is computed by an exponential size circuit. We state and prove our result for a concrete setting of parameters  $m = 3n, k = n - 3$ . We note that our work can be extended to a much wider set of parameters. However, we did not see any applications of pursuing this.

**Theorem 1.12 (Main theorem).** *Let  $A$  be an  $(n, m, k; n', m', k')$  Entropy Reverser for  $m = 3n, k = n - 3$ . Then  $QC(A) \geq 2^{n/5} / \text{poly}(n', m')$ .*

### 1.3 Related works

Relations between zero knowledge proofs have been previously studied [HR05; Vad16], where certain black box reductions were ruled out. However, previous works only ruled out restricted forms of black box reductions, where the only access to a function  $f$  is via independent and uniform samples. In particular, these reductions are non adaptive. We note that this is a much weaker notion of black box reductions, and indeed some of the black box reductions we already mentioned (e.g. the reduction from Statistical Difference to its complement [SV03]) require the ability to correlate inputs. As far as we know, ours is the first work in this context which rules out general black box reductions without any restriction on the access pattern or adaptivity.

### 1.4 Proof overview

Let  $n \geq 1$  and fix  $m = 3n, k = n - 3$ .

The first step in our proof is to apply a black box reduction of Goldreich et al. [GSV99], which converts high / low entropy distributions to distributions which are close to uniform, or supported on a small set, respectively (Lemma 3.1). This allows us to assume stronger properties of the functions generated by the supposed Entropy Reverser. Concretely, that we are given a black box reduction  $A$  from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n',m'}$  such that:

- If  $H(f) \geq k + 1$  then  $A^f$  is distributed close to uniform (concretely,  $\text{dist}(A^f, \mathcal{U}) \leq 0.1$ ).
- If  $H(f) \leq k - 1$  then the distribution of  $A^f$  is supported on a small set (concretely, of size  $\leq 0.1 \cdot 2^{m'}$ ).

As this black box reduction is efficient, it incurs a blowup of only  $\text{poly}(n', m')$  in the query complexity. See Section 3.1 for the details. From now on, we focus on this stronger notion of an Entropy Reverser, and show that for it it holds that  $QC(A) \geq \Omega(2^{n/5})$ .

Next, we consider several distributions over functions  $\mathcal{F}_{n,m}$ . Fix  $b = 256$ . We denote by  $\mathbf{B} = (B_1, \dots, B_s)$  a partition of the input space  $\{0, 1\}^n$  into  $s$  blocks, each of size  $b$ . For  $0 \leq j \leq s$  we define a distribution  $\mathcal{D}_j$  over  $\mathcal{F}_{n,m}$  as follows:

- Sample a random partition  $\mathbf{B} = (B_1, \dots, B_s)$  of  $\{0, 1\}^n$ .
- Sample  $y_1, \dots, y_j \in \{0, 1\}^m$  uniformly and independently.
- If  $x \in B_i, i \leq j$  then set  $f_j(x) = y_i$ .
- If  $x \in B_i, i > j$  then sample  $f_j(x) \in \{0, 1\}^m$  uniformly and independently.

It is not hard to show that as  $j$  increases, the entropy of  $f_j \sim D_j$  decreases. Concretely, we show (Claim 3.3) that with very high probability it holds that

$$H(f_{s/4}) = n - 2, \quad H(f_{s/2}) = n - 4.$$

Thus, by the assumptions of our Entropy Reverser (as we set  $k = n - 3$ ), it should hold that  $A^{f_{s/4}}$  is supported on a small set, while  $A^{f_{s/2}}$  is distributed close to uniform. We show that this requires exponential query complexity. From now onwards, let  $q = \text{QC}(A)$  denote this query complexity.

Let  $z \in \{0, 1\}^{m'}$  be chosen uniformly, and let  $p_j$  for  $0 \leq j \leq s$  denote the probability that  $z$  belongs to the support of  $A^{f_j}$ :

$$p_j = \Pr[\exists w \in \{0, 1\}^{n'}, z = A^{f_j}(w)].$$

By our assumptions,  $p_{s/2} \geq 0.9$  while  $p_{s/4} \leq 0.1$ . Our goal is to apply a hybrid argument and show that if  $q$  is small then  $p_{j-1} \approx p_j$  for all  $s/4 \leq j \leq s/2$ .

We can couple the choice of  $f_{j-1}, f_j$ , so that we jointly sample  $\mathbf{B}, y_1, \dots, y_j$ , and the only difference between  $f_{j-1}$  and  $f_j$  is that they differ on the block  $B_j$  ( $f_{j-1}$  maps each point in  $B_j$  to a uniformly chosen point in  $\{0, 1\}^{m'}$ , while  $f_j$  maps all the points in  $B_j$  to a single point). As the partition to blocks is random, the probability that a specific query belongs to the block  $B_j$  is  $1/s$ . As the algorithm makes  $q$  queries, this should “intuitively” give the bound

$$|p_j - p_{j-1}| \leq \frac{q}{s}.$$

(we say “intuitively” as the black box reduction is an adaptive algorithm, while the above analysis works straightforwardly only for non-adaptive algorithms). However, such a bound is useless for us, as we need to apply it  $\Omega(s)$  times. Thus, we need a more refined analysis.

In order to do so, let  $f \in \{f_{j-1}, f_j\}$ . We say that an input  $w$  to  $A^f$  respects the block structure of  $\mathbf{B}$  if any block  $B_i$  in  $\mathbf{B}$  is queried at most once by  $A^f(w)$ . Intuitively, such an input should not be able to “distinguish” between  $f_{j-1}$  and  $f_j$ . On the other hand, the probability over a random partition that any two fixed points belong to  $B_j$  is  $\approx 1/s^2$ , and hence as there are  $q$  queries, this should “intuitively” give an improved bound of

$$|p_j - p_{j-1}| \leq \frac{q^2}{s^2}.$$

If such a bound is indeed true, then applying it  $(s/2) - (s/4) = s/4$  times would give that  $|p_{s/4} - p_{s/2}| \leq O(q^2/s)$ , which would imply that  $q^2 \geq \Omega(s) = \Omega(2^n)$ , and hence we obtain an exponential lower bound on  $q$ .

Formalizing this intuition turns out to be quite delicate, as the algorithm  $A^f$  is an adaptive algorithm, and hence various choices are dependent on each other. Our main technical Lemma (Lemma 3.8) show that, if we restrict our attention to inputs which respect the block structure and define

$$p'_j = \Pr[\exists w \in \{0, 1\}^{n'}, z = A^{f_\ell}(w), w \text{ respects the block structure of } \mathbf{B}]$$

then  $p'_j$  is a good proxy for  $p_j$  (Lemma 3.6), for which a better bound can be obtained:

$$|p'_j - p'_{j-1}| \leq O\left(\frac{q^{5/3}}{s^{4/3}}\right).$$

While this bound is worse than the “intuitive” bound of  $q^2/s^2$ , it still suffices for our purposes, as when we apply it  $\Theta(s)$  times we obtain that  $|p'_{s/2} - p'_{s/4}| \leq O(q^{5/3}/s^{1/3})$  and hence we still get an exponential lower bound on  $q$ , namely  $q \geq \Omega(2^{n/5})$ .

**Paper organization.** We give some preliminary definitions in Section 2. In Section 3 we formalize the above proof overview, and give the proof of our main theorem, Theorem 1.12, assuming our main technical lemma, Lemma 3.8. The proof of Lemma 3.8 is given in Section 4. We conclude with some open problems in Section 5.

**Acknowledgements.** We would like to thank Benny Applebaum, Mika Goos, Iftach Haitner, Amit Sahai and Salil Vadhan for helpful discussion. Jiapeng Zhang also would like to thank Yingcong Li and Yue Gao for encouragement.

## 2 Preliminaries

Let  $\mathcal{F}_{n,m}$  denote the family of functions  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$ . A *black box reduction* from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n',m'}$  with is an algorithm which, given query access to  $f \in \mathcal{F}_{n,m}$ , computes a function  $A^f \in \mathcal{F}_{n',m'}$  as follows. Given an input  $w \in \{0, 1\}^{n'}$ , the algorithm makes a number of queries to  $f$  (the query locations can depend on  $w$  and be adaptive), and outputs a value  $z \in \{0, 1\}^{m'}$ . We define  $A^f(w) = z$ . The query complexity of  $A$  is the maximum number of queries to  $f$  performed over an input, which we denote by  $\text{QC}(A)$ .

Let  $X$  be a random variable taking values in  $\{0, 1\}^m$ . We recall some basic definitions. The support of  $X$  is  $\text{supp}(X) = \{x : \Pr[X = x] > 0\}$ . The Shannon entropy of  $X$  is

$$H(X) = \sum_x \Pr[X = x] \cdot \log_2(1/\Pr[X = x]).$$

The statistical distance of two random variables  $X, Y$  is

$$\text{dist}(X, Y) = \frac{1}{2} \sum_x |\Pr[X = x] - \Pr[Y = x]|.$$

We denote by  $\mathcal{U}_m$  the uniform distribution over  $\{0, 1\}^m$ .

We will identify  $f \in \mathcal{F}_{n,m}$  with the random variable of its output distribution in  $\{0, 1\}^m$ , given a uniformly sampled input in  $\{0, 1\}^n$ . As such, we extend the definition of support, Shannon entropy and statistical distance to functions. In the special case where  $f$  is computable by a circuit  $C : \{0, 1\}^n \rightarrow \{0, 1\}^m$  of size  $\text{poly}(m)$ , we say that this distribution is an *efficiently sampleable distribution*.

## 3 Proof of main theorem: Theorem 1.12

### 3.1 A useful reduction

As a first step towards proving Theorem 1.12, we make use of a black box reduction of Goldreich et al. [GSV99]. It allows us to strengthen the assumptions in Theorem 1.12.

**Lemma 3.1** ([GSV99]). *Let  $n', m', k' \geq 1$ . There exists a black box reduction  $A_{3.1}$  from  $\mathcal{F}_{n',m'}$  to  $\mathcal{F}_{n'',m''}$ , where  $n'', m''$ ,  $\text{QC}(A_{3.1}) \leq \text{poly}(n', m')$ , such that the following holds for any  $f \in \mathcal{F}_{n',m'}$ :*

- If  $H(f) \geq k' + 1$  then  $\text{dist}(A_{3.1}^f, \mathcal{U}_{m''}) \leq 0.1$ .
- If  $H(f) \leq k' - 1$  then  $|\text{supp}(A_{3.1}^f)| \leq 0.1 \cdot 2^{m''}$ .

Let  $A_{1.12}$  denote the black box from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n',m'}$  reduction assumed in Theorem 1.12. Let  $A_{3.1}$  denote the black box from  $\mathcal{F}_{n',m'}$  to  $\mathcal{F}_{n'',m''}$  given in Lemma 3.1. Let  $A$  be their composition. Namely,  $A$  is a black box reduction from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n'',m''}$ , obtained by first applying  $A_{1.12}$  and then  $A_{3.1}$ . Namely,

$$A^f = A_{3.1}^{A_{1.12}^f}.$$

Observe that  $\text{QC}(A) \leq \text{QC}(A_{1.12})\text{QC}(A_{3.1}) \leq \text{QC}(A_{1.12})\text{poly}(n', m')$ , that  $n'', m'' \leq \text{poly}(n', m')$ , and that  $A$  satisfies the following:

- If  $H(f) \geq k + 1$  then  $|\text{supp}(A^f)| \leq 0.1 \cdot 2^{m''}$ .
- If  $H(f) \leq k - 1$  then  $\text{dist}(A^f, \mathcal{U}_{m''}) \leq 0.1$ .

We will prove a lower bound on the query complexity of  $A$ , which would then imply a lower bound on the query complexity of  $A_{1.12}$ .

## 3.2 Preparations

In order to prove Theorem 1.12, we will exhibit two distributions over functions, one of high entropy functions, the other of low entropy functions, and show that black box reductions with low query complexity cannot “reverse” the entropy relation between them.

**Definition 3.2 (Sample distribution).** Let  $n, m \geq 1$  and let  $b \geq 2$  be a parameter (block size) to be determined later, and set  $s = 2^n/b$ . We denote by  $\mathbf{B} = (B_1, \dots, B_s)$  a partition of  $\{0, 1\}^n$  into  $s$  blocks of equal size  $2^n/s = b$ . For any  $0 \leq j \leq s$  we define a distribution over partitions  $\mathbf{B}$  and functions  $f_j \in \mathcal{F}_{n,m}$  as follows:

- Sample a random partition  $\mathbf{B} = (B_1, \dots, B_s)$  of  $\{0, 1\}^n$ .
- Sample  $y_1, \dots, y_j \in \{0, 1\}^m$  uniformly and independently.
- If  $x \in B_i, i \leq j$  then set  $f_j(x) = y_i$ .
- If  $x \in B_i, i > j$  then sample  $f_j(x) \in \{0, 1\}^m$  uniformly and independently.

We denote the joint distribution of  $(\mathbf{B}, f_j)$  as  $\mathcal{D}_j$ . With an abuse of notation, when we write  $f_j \sim \mathcal{D}_j$ , we simply omit the block structure from the sample. Note that  $f_0 \sim \mathcal{D}_0$  is uniformly distributed over  $\mathcal{F}_{n,m}$ . The following simple claim argues that as we increase  $j$ , the entropy of  $f_j \sim \mathcal{D}_j$  decreases. It is specialized to our desired application.

**Claim 3.3.** *Let  $m = 3n$ . Sample  $f_j \sim \mathcal{D}_j$ . Then with probability  $1 - 2^{-n}$  over the choice of  $f_j$ , it holds that*

$$H(f_j) = n - (j/s) \log b.$$

*In particular, if we set  $b = 256$  then*

$$H(f_{s/4}) = n - 2, \quad H(f_{s/2}) = n - 4.$$

*Proof.* Let  $0 \leq j \leq s$  and sample  $f_j \sim \mathcal{D}_j$ . Let  $y_1, \dots, y_j$  be the single value that  $f_j$  obtains on blocks  $B_1, \dots, B_j$ . Consider  $y_1, \dots, y_j, (f_j(x) : x \in B_i, i > j)$ . Lets denote by  $E_0$  the event that no two values in this list collide. The probability that any two of these values are equal is  $2^{-m}$ . As there are  $\leq 2^n$  values, the probability that any two intersect is bounded by  $2^{2n-m} \leq 2^{-n}$ . Thus  $\Pr[E_0] \geq 1 - 2^{-n}$ .



Lets assume that  $E_0$  holds. Then, the distribution of  $f_j$  is as follows: there are  $j$  values (namely,  $y_1, \dots, y_j$ ) that each is obtained with probability  $1/s = b/2^n$ . All other  $2^n - bj$  values are each obtained with probability  $2^{-n}$ . Thus

$$H(f_j|E_0) = j \cdot (b/2^n) \cdot \log(2^n/b) + (2^n - bj) \cdot 2^{-n} \cdot \log(2^n) = n - (j/s) \log b.$$

We treat  $\mathcal{D}_{s/4}$  as a distribution over (mostly) high entropy functions, and  $\mathcal{D}_{s/2}$  as a distribution over (mostly) lower entropy functions.

### 3.3 Block compatible inputs

Given  $w \in \{0, 1\}^{n''}$  and  $f \in \mathcal{F}_{n,m}$ , we denote by  $\text{Query}(A^f(w)) \subset \{0, 1\}^n$  the set of inputs of  $f$  queried by  $A^f$  on input  $w$ . To recall, the functions that we focus attention on are defined together with a block structure  $\mathbf{B} = (B_1, \dots, B_s)$ . Below, we specialize our attention to inputs and their corresponding outputs, for which at most one value in each block is queried.

**Definition 3.4 (Block compatible inputs).** Let  $\mathbf{B} = (B_1, \dots, B_s)$  be a partition of  $\{0, 1\}^n$ ,  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$ . We say that  $w \in \{0, 1\}^{n''}$  is a block compatible input with respect to  $(f, \mathbf{B})$  if, when computing  $A^f(w)$ , each block of  $\mathbf{B}$  is queried at most once. We denote by  $I(f, \mathbf{B})$  the set of all block compatible inputs:

$$I(f, \mathbf{B}) = \{w \in \{0, 1\}^{n''} : |\text{Query}(A^f(w)) \cap B_i| \leq 1 \quad \forall i = 1, \dots, s\}$$

**Definition 3.5 (Block compatible outputs).** Let  $\mathbf{B}$  be a partition of  $\{0, 1\}^n$ ,  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$ . We say that  $z \in \{0, 1\}^{m''}$  is a block compatible output with respect to  $(f, \mathbf{B})$  if  $A^f(w) = z$  for a block compatible input  $w$ . We denote by  $O(f, \mathbf{B})$  the set of all block compatible outputs:

$$O(f, \mathbf{B}) = \{z \in \{0, 1\}^{m''} : \exists w \in I(f, \mathbf{B}), A^f(w) = z\}.$$

Observe that the definition of  $I(f, \mathbf{B}), O(f, \mathbf{B})$  does not depend on the order of the blocks in  $\mathbf{B}$ . This will turn out to be crucial later on in the analysis. Thus, for  $\mathbf{B} = (B_1, \dots, B_s)$  define  $\{\mathbf{B}\} = \{B_1, \dots, B_s\}$  (that is, forgetting the order of the blocks) and note that

$$I(f, \mathbf{B}) = I(f, \{\mathbf{B}\}) \quad O(f, \mathbf{B}) = O(f, \{\mathbf{B}\}).$$

It is obvious that  $O(f, \mathbf{B}) \subset \text{supp}(A^f)$ . Next, we argue that if the distribution of  $A^{f_j}$  is close to uniform, then  $O(f_j, \mathbf{B})$  is large.

**Lemma 3.6.** *Sample  $(\mathbf{B}, f_j) \sim \mathcal{D}_j$ , and assume that  $\text{dist}(A^{f_j}, \mathcal{U}_{m''}) \leq \varepsilon$ . Then*

$$\mathbb{E}[|O(f_j, \mathbf{B})|] \geq \left(1 - \frac{q^2}{s} - \varepsilon\right) 2^{m''}.$$

*Proof.* We first argue that for each fixed  $w$ ,

$$\Pr_{(\mathbf{B}, f_j) \sim \mathcal{D}_j} [w \in I(f_j, \mathbf{B})] \geq 1 - \frac{q^2}{s}.$$

To see that, let  $Q = \{(x_1, y_1, \dots, x_q, y_q)\} \subset \{0, 1\}^{q(n+m)}$  be all possible queries and answers made by  $A^f(w)$ . That is,  $x_1 = x_1(w)$  is the first query made. If  $f(x_1) = y_1$  then  $x_2$  is the second query made, and so on. Note that each  $x_i$  is determined by  $w, x_1, y_1, \dots, x_{i-1}, y_{i-1}$ , while  $y_i$  can take any value in  $\{0, 1\}^m$ . In particular,  $|Q| = 2^{mq}$ .

Next, fix  $x_1, \dots, x_q$  and let  $\mathbf{B}$  be a randomly chosen partition. Then

$$\Pr_{\mathbf{B}}[x_1, \dots, x_q \text{ in distinct blocks}] \geq 1 - \sum_{i \neq j} \Pr_{\mathbf{B}}[x_i, x_j \text{ in the same block}] \geq 1 - \frac{q^2}{s}.$$

Note that if  $x_1, \dots, x_q$  are in distinct blocks, then  $f_j(x_1), \dots, f_j(x_q)$  are independently and uniformly chosen in  $\{0, 1\}^m$ . Thus

$$\begin{aligned} \Pr_{(\mathbf{B}, f_j) \sim \mathcal{D}_j} [w \in I(f_j, \mathbf{B})] &= \sum_{(x_1, y_1, \dots, x_q, y_q) \in Q} \Pr[w \in I(f_j, \mathbf{B}) \wedge f_j(x_1) = y_1 \wedge \dots \wedge f_j(x_q) = y_q] \\ &= \sum_{(x_1, y_1, \dots, x_q, y_q) \in Q} \Pr[x_1, \dots, x_q \text{ in distinct blocks} \wedge f_j(x_1) = y_1 \wedge \dots \wedge f_j(x_q) = y_q] \\ &= \sum_{(x_1, y_1, \dots, x_q, y_q) \in Q} \Pr[x_1, \dots, x_q \text{ in distinct blocks}] \cdot \\ &\quad \Pr[f_j(x_1) = y_1 \wedge \dots \wedge f_j(x_q) = y_q | x_1, \dots, x_q \text{ in distinct blocks}] \\ &\geq \sum_{(x_1, y_1, \dots, x_q, y_q) \in Q} \left(1 - \frac{q^2}{s}\right) 2^{-mq} \\ &= 1 - \frac{q^2}{s}. \end{aligned} \quad \square$$

We next consider  $O(f_j, \mathbf{B})$ . Recall that we assume that the distribution of  $A^{f_j}$  is  $\varepsilon$ -close in statistical distance to the uniform distribution  $\mathcal{U}_{m''}$ . Let  $w \in \{0, 1\}^{n''}$  be chosen uniformly and consider the random variable  $z = A^{f_j}(w)$ . We have

$$\Pr_{(\mathbf{B}, f_j) \sim \mathcal{D}_j, w \in \{0, 1\}^{n''}} [z \in O(f_j, \mathbf{B})] \geq \Pr[w \in I(f_j, \mathbf{B})] \geq 1 - \frac{q^2}{s}.$$

On the other hand, let  $u \in \{0, 1\}^{m''}$  be chosen uniformly and independently of all other random variables. As  $\text{dist}(z, u) = \text{dist}(A^{f_j}, \mathcal{U}_{m''}) \leq \varepsilon$  we have, for every fixing of  $\mathbf{B}, f_j$ , that

$$\Pr[u \in O(f_j, \mathbf{B}) | \mathbf{B}, f_j] \geq \Pr[z \in O(f_j, \mathbf{B}) | \mathbf{B}, f_j] - \varepsilon.$$

Averaging over the choices of  $\mathbf{B}, f_j$  we obtain that

$$\Pr[u \in O(f_j, \mathbf{B})] \geq 1 - \frac{q^2}{s} - \varepsilon.$$

This implies that

$$\mathbb{E}[|O(f_j, \mathbf{B})|] = 2^{m''} \Pr[u \in O(f_j, \mathbf{B})] \geq \left(1 - \frac{q^2}{s} - \varepsilon\right) 2^{m''}.$$

### 3.4 Main technical lemma

The main step in proving Theorem 1.12 is showing that  $O(f_{j-1}, \mathbf{B})$  is not much smaller than  $O(f_j, \mathbf{B})$ .

**Definition 3.7.** Let us jointly sample  $\mathbf{B}, f_{j-1}, f_j$  as follows:

- Sample  $(\mathbf{B}, f_{j-1}) \sim \mathcal{D}_{j-1}$ .
- Sample  $y_j \in \{0, 1\}^m$  independently and uniformly, and set

$$f_j(x) = \begin{cases} f_{j-1}(x) & x \notin B_j \\ y_j & x \in B_j. \end{cases}$$

We denote this joint distribution over  $\mathbf{B}, f_{j-1}, f_j$  by  $\mathcal{D}_{j-1,j}$ . Observe that if we omit  $f_{j-1}$ , then the marginal distribution over  $(\mathbf{B}, f_j)$  is indeed  $\mathcal{D}_j$ .

**Lemma 3.8.** *Assume that  $\delta s \leq j \leq (1 - \delta)s$  and sample  $(\mathbf{B}, f_{j-1}, f_j) \sim \mathcal{D}_j$ . Then for any  $z \in \{0, 1\}^{m''}$  it holds that*

$$\Pr[z \in O(f_{j-1}, \mathbf{B})] \geq \Pr[z \in O(f_j, \mathbf{B})] - \varepsilon,$$

where  $\varepsilon = \frac{4q^{5/3}b^{2/3}}{\delta^{4/3}s^{4/3}}$ . If we set  $b = 256$  and  $\delta = 1/4$  then  $\varepsilon = O(q^{5/3}/s^{4/3})$ . In particular, averaging over a uniform choice of  $z$  gives that

$$\mathbb{E}[|O(f_{j-1}, \mathbf{B})|] \geq \mathbb{E}[|O(f_j, \mathbf{B})|] - \varepsilon 2^{m''}.$$

We note that it is crucial in Lemma 3.8 that  $\varepsilon \ll 1/s$ , as we will apply it to relate  $O(f_{s/4}, \mathbf{B})$  to  $O(f_{s/2}, \mathbf{B})$ , which will incur an additional factor of  $s$ . Most of the technical challenge in proving Lemma 3.8 is achieving that, as achieving weaker bounds of the form  $\varepsilon = \text{poly}(q)/s$  is much easier. We defer the proof of Lemma 3.8 to Section 4, and next show how it implies Theorem 1.12.

### 3.5 Deducing Theorem 1.12

Let  $A_{1.12}$  be the assumed black box reduction given in Theorem 1.12, specialized for  $m = 3n, k = n - 3$ . Let  $n', m'$  denote the input and output size of  $A_{1.12}^f$  in this case and let  $k' = k'(n, m, k)$ . Let  $A$  be the black box reduction obtained by first applying  $A_{1.12}$  to  $f \in \mathcal{F}_{n,m}$ , then applying  $A_{3.1}$  to  $A_{1.12}^f$ . Thus  $A^f \leq \mathcal{F}_{n'', m''}$  where  $n'', m'' \leq \text{poly}(n', m')$ . We have  $\text{QC}(A) \leq \text{QC}(A_{1.12})\text{QC}(A_{3.1}) \leq \text{QC}(A_{1.12})\text{poly}(n', m')$ .

**Definition 3.9 (Hybrid distribution).** Sample  $(f_j : j = 0, \dots, s)$  jointly as follows:

- Sample a random partition  $\mathbf{B} = (B_1, \dots, B_s)$  of  $\{0, 1\}^n$ .
- Sample  $y_1, \dots, y_s \in \{0, 1\}^m$  uniformly and independently.
- Sample a uniform function  $g : \{0, 1\}^n \rightarrow \{0, 1\}^m$ .
- If  $x \in B_i, i \leq j$  then set  $f_j(x) = y_i$ .
- If  $x \in B_i, i > j$  then sample  $f_j(x) = g(x)$ .

Observe that the marginal distribution of  $(\mathbf{B}, f_j)$  is  $\mathcal{D}_j$ , and moreover, the marginal distribution of  $(\mathbf{B}, f_{j-1}, f_j)$  is  $\mathcal{D}_{j-1,j}$ . According to Claim 3.3, we have the following statements by setting  $b = 256$ ,

- $\Pr_{f_{s/4} \sim \mathcal{D}_{s/4}}[\text{H}(f_{s/4}) = n - 2] \geq 1 - 2^{-n}$ .
- $\Pr_{f_{s/2} \sim \mathcal{D}_{s/2}}[\text{H}(f_{s/2}) = n - 4] \geq 1 - 2^{-n}$ .

By the guarantees of  $A$  we have that

- If  $H(f_{s/4}) \geq (n-3) + 1$  then  $|\text{supp}(A^{f_{s/4}})| \leq 0.1 \cdot 2^{m''}$ .
- If  $H(f_{s/2}) \leq (n-3) - 1$  then  $|\text{dist}(A^{f_{s/2}}, U_{m''})| \leq 0.1$ .

Let  $q = \text{QC}(A)$ . Applying Lemma 3.6 to  $f_{s/2}$ , and assuming that  $q^2/s \leq 0.1$  gives that

$$\mathbb{E} \left[ |O(A^{f_{s/2}}, \mathbf{B})| \right] \geq 0.8 \cdot 2^{m''}.$$

On the other hand,

$$\mathbb{E} \left[ |O(A^{f_{s/4}}, \mathbf{B})| \right] \leq |\text{supp}(A^{f_{s/4}})| \leq 0.1 \cdot 2^{m''}.$$

Lemma 3.8, applied for  $s/4 \leq j \leq s/2$ , gives that

$$\mathbb{E} \left[ |O(A^{f_{j-1}}, \mathbf{B})| \right] \geq \left[ |O(A^{f_j}, \mathbf{B})| \right] - \varepsilon 2^{m''},$$

where  $\varepsilon = O(q^{5/3}/s^{4/3})$ . For all these to hold we need to have

$$\varepsilon(s/2 - s/4) \geq 0.7$$

which gives the required bound

$$\text{QC}(A) = q \geq \Omega(s^{1/5}).$$

This then gives us the bound

$$\text{QC}(A_{1.12}^f) \geq \Omega(2^{n/5}/\text{poly}(n', m')).$$

## 4 Proof of main technical lemma: Lemma 3.8

We prove Lemma 3.8 in this section. To recall,  $A$  is a black box reduction from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n'',m''}$ . We fix  $1 \leq j \leq s$  and  $z \in \{0,1\}^{m''}$  from here onwards. We sample  $(\mathbf{B}, f_{j-1}, f_j) \sim \mathcal{D}_{j-1,j}$  and wish to compare  $\Pr[z \in O(f_{j-1}, \mathbf{B})]$  and  $\Pr[z \in O(f_j, \mathbf{B})]$ . To simplify notations define

$$O(f_{j-1}) = O(f_{j-1}, \mathbf{B}) \quad I(f_{j-1}) = I(f_{j-1}, \mathbf{B}).$$

Define the events

$$X := [z \in O(f_{j-1})] \quad Y := [z \in O(f_j)].$$

Our goal is to show that if  $Y$  holds, then with high probability also  $X$  holds. The ‘‘common information’’ between  $X, Y$  is captured by the random variable

$$\mathcal{C} := (\{(B_i, f_j|_{B_i})\}_{1 \leq i \leq j-1}, B_j, \{(B_i, f_j|_{B_i})\}_{j+1 \leq i \leq s}).$$

Observe that  $\{\mathbf{B}\}$  can be computed from  $\mathcal{C}$ , which we denote as  $\{\mathbf{B}\} = \{\mathbf{B}\}(\mathcal{C})$ , and that furthermore

$$f_{j-1} = f_{j-1}(\mathcal{C}, f_{j-1}|_{B_j}) \quad f_j = f_j(\mathcal{C}, f_j|_{B_j}).$$

Thus

$$X = X(\mathcal{C}, f_{j-1}|_{B_j}) \quad Y = Y(\mathcal{C}, f_j|_{B_j}).$$

In particular, given any fixing of  $\mathcal{C}$ , we have that  $f_{j-1}|_{B_j}$  is a uniform function from  $B_j$  to  $\{0,1\}^m$ , that  $f_j|_{B_j}$  is a random constant function, and that the two are independent of each other. We obtain the following claim:

**Claim 4.1.** For any fixing of  $\mathcal{C}$ , the random variables  $X|\mathcal{C}$  and  $Y|\mathcal{C}$  are independent.

Recall that  $f_{j-1}$  and  $f_j$  differ only in their evaluation on the block  $B_j$ . We define a partial function  $\hat{f}$  to be the set of inputs where  $f_{j-1}$  and  $f_j$  agree, namely all inputs outside  $B_j$ , and outputs “?” otherwise. Formally, we define the function  $\hat{f} : \{0, 1\}^n \rightarrow (\{0, 1\}^m \cup \{?\})$  as follows:

$$\hat{f}(x) = \begin{cases} f_{j-1}(x) & \text{if } x \notin B_j \\ ? & \text{if } x \in B_j \end{cases}$$

if  $x \notin B_j$  then  $\hat{f}(x) = f_{j-1}(x) = f_j(x)$ . As we now allow for partial functions, we will also need to allow running the black box reduction  $A$  on partial functions. We do so by outputting a “?” if the black box reduction queries a point where the partial function is not defined. Observe that  $\hat{f}$  can be computed given  $\mathcal{C}$ :

$$\hat{f} = \hat{f}(\mathcal{C}).$$

**Definition 4.2 (Black box reduction of a partial function).** Let  $A$  be a black box reduction from  $\mathcal{F}_{n,m}$  to  $\mathcal{F}_{n',m''}$ . Let  $f : \{0, 1\}^n \rightarrow (\{0, 1\}^m \cup \{?\})$  be a partial function. We define  $A^f : \{0, 1\}^{n'} \rightarrow (\{0, 1\}^{m''} \cup \{?\})$  to be the following partial function. When computing  $A^f(w)$ , follows the queries made by  $A$  as if  $f$  was a total function. However, if at any point we query a point  $x$  where  $f(x) = ?$  then we abort and output ?.

We also extend the definition of block compatible inputs and outputs to partial functions in the obvious manner. Define  $O(\hat{f}) := O(\hat{f}, \{\mathbf{B}\})$  and define the event  $E_1$  as

$$E_1 = E_1(\mathcal{C}) := [z \in O(\hat{f})].$$

**Claim 4.3.** If  $E_1$  holds then both  $X$  and  $Y$  also hold:  $E_1 \Rightarrow X \wedge Y$ .

*Proof.* If  $E_1$  holds then by definition, there exists  $w \in I(\hat{f})$  for which  $A^{\hat{f}}(w) = z$ . This implies that  $A^{f_{j-1}}(w) = A^{f_j}(w) = z$ , as since  $A^{\hat{f}}(w)$  didn't return a “?”, it only queried locations outside  $B_j$ , where  $f_{j-1}, f_j$  agree. Also, as  $w \in I(\hat{f})$  this means that  $A^{\hat{f}}(w)$  queries each block  $B_i$  at most once, while block  $B_j$  is never queried. Thus also  $w \in I(f_{j-1}), w \in I(f_j)$ . This implies that  $z \in O(f_{j-1}), z \in O(f_j)$  which means that  $X, Y$  hold.  $\square$

According to Claim 4.3 we have,

$$\Pr[X \wedge E_1] = \Pr[Y \wedge E_1] = \Pr[E_1]$$

and hence

$$\begin{aligned} & \Pr[X] - \Pr[Y] \\ &= \Pr[X \wedge E_1] + \Pr[X \wedge \neg E_1] - \Pr[Y \wedge E_1] - \Pr[Y \wedge \neg E_1] \\ &= \Pr[X \wedge \neg E_1] - \Pr[Y \wedge \neg E_1] \end{aligned}$$

Thus, from now on we focus on the case that  $\neg E_1$  holds.

#### 4.1 Analyzing the case that $E_1$ doesn't hold

For each  $x \in B_j, y \in \{0, 1\}^m$  we define the following extension of  $\hat{f}$ . Define a partial function  $\hat{f}_{x,y} : \{0, 1\}^n \rightarrow (\{0, 1\}^m \cup \{?\})$  as follows:

$$\hat{f}_{x,y}(x') := \begin{cases} \hat{f}(x') & \text{if } x' \notin B_j \\ y & \text{if } x' = x \\ ? & \text{if } x' \in B_j \text{ and } x' \neq x \end{cases}$$

For each  $x \in B_j$  define

$$R(x) = R_{\hat{f}}(x) := \{y \in \{0, 1\}^m : z \in O(f_{x,y})\}.$$

Namely,  $R(x)$  is the set of values  $y$  for which, if we allow the algorithm to make a single query to  $B_j$  at point  $x$  which returns  $y$ , then  $z$  becomes a block compatible output. Observe that the definition of  $R(x)$  depends only on  $\hat{f}$ , and hence on  $\mathcal{C}$ . Crucially, it does not depend on the values of either  $f_{j-1}$  or  $f_j$  on  $B_j$ . We further define

$$r(x) := \Pr_{y \in \{0,1\}^m} [y \in R(x)] = \frac{|R(x)|}{2^m}.$$

As  $R(x), r(x)$  depend only on  $\mathcal{C}$ , we may consider the following experiment: first sample  $\mathcal{C}$  and then sample  $f_{j-1}, f_j$  conditioned on  $\mathcal{C}$ . That is,  $f_{j-1}, f_j$  are equal to  $\hat{f} = \hat{f}(\mathcal{C})$  outside  $B_j = B_j(\mathcal{C})$ , and on  $B_j$  we sample  $f_{j-1}$  as a random function, while  $f_j(x) = y_j$  for all  $x \in B_j$ , where  $y_j$  is randomly chosen. Recall that  $E_1 = E_1(\mathcal{C})$ .

**Claim 4.4.** *For any fixing of  $\mathcal{C}$  sample  $f_{j-1}, f_j | \mathcal{C}$ . Then*

$$\Pr[X | \mathcal{C}, \neg E_1] = 1 - \prod_{x \in B_j} (1 - r(x))$$

and

$$\Pr[Y | \mathcal{C}, \neg E_1] \leq \sum_{x \in B_j} r(x).$$

*Proof.* Consider any fixing of  $\mathcal{C}$  and sample  $f_{j-1}, f_j$  conditioned on it. Recall that  $r(x)$  is a function of  $\mathcal{C}$ .

Let  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$  be any function which agrees with  $\hat{f}$  on all  $x \notin B_j$ . If  $z \in O(f)$  then there exists  $w \in I(f)$  for which  $A^f(w) = z$ . As we assume that  $\neg E_1$  holds, for any such  $w$ ,  $A^f(w)$  must query the block  $B_j$  at least once, and since  $w \in I(f)$  it is exactly once, say at point  $x_w \in B_j$ . But then also  $z \in O(\hat{f}_{x_w, f(x_w)})$ , which means that  $f(x_w) \in R(x_w)$ . The converse direction also holds: if  $f(x) \in R(x)$  for any  $x \in B_j$  then by definition of  $R(x)$ , there exists  $w_x$  such that  $w_x \in I(f)$  and  $A^f(w_x) = z$  (and moreover  $A^f(w_x)$  queries the block  $B_j$  exactly at  $x$ ) and in particular  $z \in O(f)$ . Thus

$$z \in O(f) \iff \bigvee_{x \in B_j} [f(x) \in R(x)].$$

Next, we apply this logic to both  $f_{j-1}$  and  $f_j$ . For  $f_{j-1}$ , each point  $f_{j-1}(x)$  for  $x \in B_j$  is uniformly and independently chosen, hence

$$\begin{aligned} \Pr[X | \mathcal{C}, \neg E_1] &= \Pr[z \in O(f_{j-1}) | \mathcal{C}, \neg E_1] \\ &= 1 - \Pr[f_{j-1}(x) \notin R(x) \forall x \in B_j] \\ &= 1 - \prod_{x \in B_j} (1 - r(x)). \end{aligned}$$

For  $f_j$ , all the evaluations  $\{f_j(x) : x \in B_j\}$  are equal to a uniformly chosen point  $y_j$ . Hence by the union bound

$$\begin{aligned} \Pr[Y|\mathcal{C}, \neg E_1] &= \Pr[z \in O(f_j)|\mathcal{C}, \neg E_1] \\ &\leq \sum_{x \in B_j} \Pr[y_j \in R(x)] \\ &\leq \sum_{x \in B_j} r(x). \end{aligned} \quad \square$$

The following definition allow us to compare the two bounds appearing in Claim 4.4.

**Definition 4.5.** Let  $\gamma > 0$ . A sequence of numbers  $r_1, \dots, r_b$  is said to be  $\gamma$ -balanced if

$$(1 - \gamma) \sum r_i \leq 1 - \prod (1 - r_i).$$

Let  $\gamma > 0$  to be determined later, and define the event  $E_2$  as

$$E_2 = E_2(\mathcal{C}) := [(r(x) : x \in B_j) \text{ is } \gamma\text{-balanced}].$$

The following is a corollary of Claim 4.4 and the definition of  $X, Y$ .

**Claim 4.6.** For any fixing of  $\mathcal{C}$  for which  $\neg E_1, E_2$  hold, it holds that

$$\Pr[X|\mathcal{C}, \neg E_1, E_2] \geq (1 - \gamma) \Pr[Y|\mathcal{C}, \neg E_1, E_2].$$

*Proof.* Fix  $\mathcal{C}$  such that  $\neg E_1, E_2$  hold. This fixes in particular  $B_j$  and  $(r(x) : x \in B_j)$ . As  $E_2$  holds, we obtain by Claim 4.4 that

$$\Pr[X|\mathcal{C}] = 1 - \prod_{x \in B_j} (1 - r(x)) \geq (1 - \gamma) \sum_{x \in B_j} r(x) \geq (1 - \gamma) \Pr[Y|\mathcal{C}].$$

Following up on (1), we have

$$\begin{aligned} \Pr[X] - \Pr[Y] &= (\Pr[X \wedge \neg E_1 \wedge E_2] - \Pr[Y \wedge \neg E_1 \wedge E_2]) \\ &\quad + (\Pr[X \wedge \neg E_1 \wedge \neg E_2] - \Pr[Y \wedge \neg E_1 \wedge \neg E_2]) \end{aligned}$$

The second term can simply be bounded by  $\Pr[\neg E_1 \wedge \neg E_2]$ , which we bound in the next section. For now, lets focus on the first term. Consider any fixing of  $\mathcal{C}$  for which  $\neg E_1, E_2$  hold. By Claim 4.6 we have that  $\Pr[X|\mathcal{C}] \geq (1 - \gamma) \Pr[Y|\mathcal{C}]$ . By averaging over such  $\mathcal{C}$ , we obtain that

$$\Pr[X \wedge \neg E_1 \wedge E_2] - \Pr[Y \wedge \neg E_1 \wedge E_2] \geq -\gamma \Pr[Y \wedge \neg E_1 \wedge E_2].$$

We can bound the right hand side by

$$\Pr[Y \wedge \neg E_1 \wedge E_2] \leq \Pr[Y \wedge \neg E_1] = \Pr[Y] \Pr[\neg E_1|Y].$$

**Claim 4.7.**  $\Pr[\neg E_1|Y] \leq q/j$ .

*Proof.* Sample  $(\mathbf{B}, f_{j-1}, f_j) \sim \mathcal{D}_{j-1,j}$ . In addition, sample  $t \in \{1, \dots, j\}$  uniformly. We will define a “proxy”  $f_{j-1}$  obtained from  $f_j$  by changing the value on  $B_t$  to a random function. We will then argue that with high probability, this misses any specific set of queries. To that end, define the following random variables:

- $f'_{j-1} : \{0, 1\}^n \rightarrow \{0, 1\}^m$  is a random function defined as follows: if  $x \notin B_t$  then  $f'_{j-1}(x) = f_j(x)$ ; and if  $x \in B_t$  then  $f'_{j-1}(x)$  is a uniformly random element in  $\{0, 1\}^m$ .
- $\hat{f}' : \{0, 1\}^n \rightarrow (\{0, 1\}^m \cup \{?\})$  is a partial function defined as follows:  $\hat{f}'(x) = f'_{j-1}(x) = f_j(x)$  if  $x \notin B_t$ , and  $\hat{f}'(x) = ?$  if  $x \in B_t$ .
- $\mathbf{B}'$  is equal to  $\mathbf{B}$  with blocks  $B_t, B_j$  swapped. Namely,  $\mathbf{B}' = (B_1, \dots, B_{t-1}, B_{t+1}, \dots, B_j, B_t, B_{j+1}, \dots, B_s)$ .

Observe that the joint distributions of  $(\mathbf{B}, f_j, f_{j-1}, \hat{f})$  and  $(\mathbf{B}', f_j, f'_{j-1}, \hat{f}')$  are identical. Next, define the following events:

- $Y' := [z \in O(f_j, \{\mathbf{B}'\})]$ .
- $E'_1 := [z \in O(\hat{f}', \{\mathbf{B}'\})]$ . □

Observe that  $Y' = Y$  since  $\{\mathbf{B}'\} = \{\mathbf{B}\}$ , and that the joint distributions of  $(E_1, Y)$  and  $(E'_1, Y) = (E'_1, Y')$  are identical. Next, fix  $\mathbf{B}, f_j$  such that  $Y$  holds. This means that there exists  $w \in I(f_j, \{\mathbf{B}\}) = I(f_j, \{\mathbf{B}'\})$  such that  $A^{f_j}(w) = z$ . Let  $Q = \text{Query}(A^{f_j}(w))$  be the set of queries made by the algorithm, where  $|Q| \leq q$ . Observe that if  $B_t \cap Q = \emptyset$  then  $E'_1$  holds, and that  $Q, t$  are independent random variables. Let  $T = T(\mathbf{B}, f_j) = \{i \in \{1, \dots, j\} : B_i \cap Q \neq \emptyset\}$ , where  $|T| \leq |Q| \leq q$ . Thus

$$\Pr[\neg E'_1 | \mathbf{B}, f_j, Y] \leq \Pr[B_t \cap Q \neq \emptyset | \mathbf{B}, f_j, Y] = \Pr[t \in T | \mathbf{B}, f_j, Y] \leq \frac{|T|}{j} \leq \frac{q}{j}.$$

By averaging over  $\mathbf{B}, f_j$ , we obtain that  $\Pr[\neg E'_1 | Y] \leq q/j$ . Thus also

$$\Pr[\neg E_1 | Y] = \Pr[\neg E'_1 | Y] \leq \frac{q}{j}.$$

We thus have

$$\Pr[X \wedge \neg E_1 \wedge E_2] - \Pr[Y \wedge \neg E_1 \wedge E_2] \geq -\gamma(q/j) \Pr[Y]$$

which implies that

$$\Pr[X] - \Pr[Y] \geq -\gamma(q/j) \Pr[Y] - \Pr[\neg E_1 \wedge \neg E_2]$$

which in turn gives the bound

$$\Pr[X] \geq (1 - \gamma(q/j)) \Pr[Y] - \Pr[\neg E_1 \wedge \neg E_2]. \tag{1}$$

To conclude, we need to upper bound  $\Pr[\neg E_1 \wedge \neg E_2]$ , which is what we do in the next section.

## 4.2 Bounding the probability that both $E_1, E_2$ don't hold

We first need a simple corollary of the definition of  $\gamma$ -balanced.

**Claim 4.8.** *Let  $r_1, \dots, r_b \in [0, 1]$  be a sequence which is not  $\gamma$ -balanced. Then there exist distinct  $1 \leq i, j \leq b$  such that  $r_i, r_j \geq \gamma/b$ .*

*Proof.* Assume not. Then without loss of generality,  $r_2, \dots, r_b \leq \gamma/b$ . By the inclusion-exclusion principle

$$1 - \prod (1 - r_i) \geq \sum r_i - \sum_{i < j} r_i r_j$$

and by our assumption

$$\sum_{i < j} r_i r_j \leq \left( \sum_{j \geq 2} r_j \right) \left( \sum_{i \geq 1} r_i \right) \leq \gamma \sum r_i.$$



We next define the notion of critical blocks. Informally, a block  $B_j$  is critical if all block compatible input  $w$  for which  $A^f(w) = z$ ,  $A^f(w)$  queries exactly one point in  $B_j$ .

**Definition 4.9 (Critical block).** Given  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$  and a partition  $\mathbf{B} = (B_1, \dots, B_s)$  of  $\{0, 1\}^n$ , we say that the block  $B_j$  is critical for  $f$  if

$$(A^f(w) = z) \wedge (w \in I(f, \mathbf{B})) \quad \Rightarrow \quad |\text{Query}(A^f(w)) \cap B_j| = 1.$$

A double critical block is a critical block where the output  $z$  can be obtained by two block compatible inputs  $w_1, w_2$  which query different points  $x_1, x_2$  in it.

**Definition 4.10 (Double critical block).** Given  $f : \{0, 1\}^n \rightarrow \{0, 1\}^m$  and a partition  $\mathbf{B} = (B_1, \dots, B_s)$  of  $\{0, 1\}^n$ , we say that the block  $B_j$  is double critical for  $f$  if

- (i)  $B_j$  is a critical block for  $f$ .
- (ii) There exist  $w_1, w_2 \in I(f, \mathbf{B})$  and distinct  $x_1, x_2 \in B_j$  such that

$$(A^f(w_i) = z) \wedge (\text{Query}(A^f(w_i)) \cap B_j = \{x_i\}) \quad i = 1, 2.$$

**Lemma 4.11.** *Sample  $(\mathbf{B}, f_{j-1}) \sim \mathcal{D}_{j-1}$ . Then*

$$\Pr[B_j \text{ is double critical for } f_{j-1}] \leq \frac{2q^3}{(s-j+1)^2}$$

*Proof.* We can jointly sample  $\mathbf{B}, f_{j-1}$  as follows:

- (1) Sample disjoint blocks  $B_1, \dots, B_{j-1}$  and  $y_1, \dots, y_{j-1} \in \{0, 1\}^m$ , and set  $f_{j-1}(x) = y_i$  if  $x \in B_i$ ,  $i < j$ .
- (2) Let  $U = \{0, 1\}^m \setminus (B_1 \cup \dots \cup B_{j-1})$ . Sample  $f_{j-1}(x) \in \{0, 1\}^m$  uniformly and independently for all  $x \in U$ .
- (3) Sample  $B_j, \dots, B_s$  a random partition of  $U$  to  $s - j + 1$  blocks of size  $b$ .

From now on, we fix  $f_{j-1}, U$  and consider only the randomness in step (3), namely the random partition of  $U$ . For simplicity of notation we say that  $B_j$  is critical, or double critical, where in both cases we refer with respect to  $f_{j-1}$ .

Define

$$W = \{w \in \{0, 1\}^{n''} : A^{f_{j-1}}(w) = z, |\text{Query}(A^{f_{j-1}}(w)) \cap B_i| \leq 1 \forall i = 1, \dots, j-1\}.$$

The set  $W$  is the set of *potential* elements in  $I(f_{j-1}, \mathbf{B})$ , in the sense that they satisfy the requirement  $|\text{Query}(A^{f_{j-1}}(w)) \cap B_i| \leq 1$  for the blocks defined so far, namely  $B_1, \dots, B_{j-1}$ . If  $W$  is empty then no block can be critical. For simplicity of notation define  $Q(w) := \text{Query}(A^{f_{j-1}}(w)) \cap U$ . Note that so far these definitions do not depend on the choice of the partition of  $U$ .

Next, sample a random partition  $(B_j, \dots, B_s)$  of  $U$ . We say that an input  $w$  is *legal* if  $A^{f_{j-1}}(w)$  queries each block at most once:

$$W_{\text{legal}} := \{w \in W : |Q(w) \cap B_i| \leq 1 \forall i = j, \dots, s\}.$$

Equivalently,  $W_{\text{legal}} = \{w \in I(f, \mathbf{B}) : A^{f_{j-1}}(w) = z\}$ . The definitions of critical and double critical can then be cast as

$$\begin{aligned} B_j \text{ is critical} &\Leftrightarrow |B_j \cap Q(w)| = 1, \forall w \in W_{\text{legal}} \\ B_j \text{ is double critical} &\Leftrightarrow B_j \text{ is critical and } |B_j \cap (\cup_{w \in W_{\text{legal}}} Q(w))| \geq 2. \end{aligned}$$

Fix  $w_1 \in W$  and assume for now that  $w_1 \in W_{\text{legal}}$ . We will handle the case that  $w_1 \notin W_{\text{legal}}$  later.

If  $B_j$  is critical then  $|B_j \cap Q(w_1)| = 1$ . Say  $B_j \cap Q(w_1) = \{x_1\}$ . If  $B_j$  is double critical then there must be another legal  $w_2 \in W_{\text{legal}}$  such that  $B_j \cap Q(w_2) = \{x_2\}$  where  $x_2 \neq x_1$ . In particular,  $x_1 \notin Q(w_2)$ . Thus, for each  $x_1 \in Q(w_1)$  define

$$W_{x_1} = \{w \in W : x_1 \notin Q(w)\}.$$

Note that if  $W_{x_1}$  is empty then it is impossible that  $B_j$  is double critical,  $w_1$  is legal and  $B_j \cap Q(w_1) = \{x_1\}$ . Thus let

$$Q'(w_1) = \{x_1 \in Q(w_1) : |W_{x_1}| \geq 1\}.$$

For each  $x_1 \in Q'(w_1)$  fix an arbitrary  $w_{x_1} \in W_{x_1}$ . By definition,  $x_1 \notin Q(w_{x_1})$ . We can bound the probability that  $B_j$  is double critical and  $w_1$  is legal by requiring that  $B_j \cap Q(w_1) = \{x_1\}$  and  $B_j \cap Q(w_{x_1}) = \{x_2\}$ , where by definition  $x_1 \neq x_2$ , and summing over all choices for  $x_1, x_2$ :

$$\Pr[B_j \text{ is double critical} \wedge w_1 \in W_{\text{legal}}] \leq \sum_{x_1 \in Q'(w_1)} \sum_{x_2 \in Q(w_{x_1})} \Pr[x_1, x_2 \in B_j] \leq \frac{q^2}{(s-j+1)^2},$$

where the bound follows from the union bound and the fact that as  $B_j, \dots, B_s$  is a random partition of  $U$ , for any fixed distinct  $x_1, x_2 \in U$  it holds that  $\Pr[x_1, x_2 \in B_j] \leq 1/(s-j+1)^2$ .

To conclude the proof, we need to handle the event that  $w_1$  is not legal. First, note that

$$\Pr[w_1 \notin W_{\text{legal}}] \leq \sum_{x_1, x_2 \in Q(w_1), x_1 \neq x_2} \Pr[x_1, x_2 \text{ in the same block}] \leq \frac{q^2}{s-j+1}.$$

We will bound  $\Pr[B_j \text{ is double critical} | w_1 \notin W_{\text{legal}}]$ . To do that, let's condition on which block does every element of  $Q(w_1)$  belong to. Let  $H_1$  denote the family of all functions  $h : Q(w_1) \rightarrow \{j, \dots, s\}$ . Let  $F_h$  denote the event

$$F_h = [x \in B_{h(x)} \quad \forall x \in Q(w_1)].$$

Note that the events  $F_h$  are disjoint, and that the event  $w_1 \notin W_{\text{legal}}$  is equivalent to  $F_h$  holding where  $h$  has at least one collision. Thus let

$$H_2 = \{h \in H_1 : \exists x_1, x_2 \in Q(w_1), h(x_1) = h(x_2)\}.$$

We have  $w_1 \notin W_l \iff \cup_{h \in H_2} F_h$ .

For each  $h \in H_2$  let  $W_h$  denote the set of  $w \in W$  for which  $Q(w)$  is not already illegal given  $h$ , namely

$$W_h = \{w \in W : \neg \exists x_1, x_2 \in Q_w \cap Q_{w_1}, h(x_1) = h(x_2)\}.$$

If  $W_h$  is empty then it is impossible that  $B_j$  is double critical and that  $F_h$  holds, as there are no legal inputs. Thus let

$$H_3 = \{h \in H_2 : |W_h| \geq 1\}.$$

For each  $h \in H_3$  fix an arbitrary  $w_h \in W_h$ . By definition, if  $B_j$  is double critical then we must have  $|B_j \cap Q(w_h)| = 1$ . We can thus bound

$$\begin{aligned} \Pr[B_j \text{ is double critical} | w_1 \notin W_{\text{legal}}] &= \sum_{h \in H_3} \Pr[B_j \text{ is double critical} | F_h] \Pr[F_h | w_1 \notin W_{\text{legal}}] \\ &\leq \sum_{h \in H_3} \sum_{x \in Q(w_h)} \Pr[x \in B_j | F_h] \Pr[F_h | w_1 \notin W_{\text{legal}}]. \end{aligned}$$

In order to help bound this expression, note that both  $h \in H_3$  and  $\Pr[F_h | w_1 \notin W_{\text{legal}}]$  are invariant to permutations of the output of  $h$ . That is, if we replace  $h(x)$  with  $\pi(h)(x) = \pi(h(x))$  for any permutation  $\pi$  on  $\{j, \dots, s\}$ , then  $h \in H_3 \iff \pi(h) \in H_3$  and  $\Pr[F_h | w_1 \notin W_{\text{legal}}] = \Pr[F_{\pi(h)} | w_1 \notin W_{\text{legal}}]$ . Thus

$$\begin{aligned} \Pr[B_j \text{ is double critical} | w_1 \notin W_{\text{legal}}] &\leq \mathbb{E}_\pi \sum_{h \in H_3} \sum_{x \in Q(w_h)} \Pr[x \in B_j | F_{\pi(h)}] \Pr[F_{\pi(h)} | w_1 \notin W_{\text{legal}}] \\ &= \mathbb{E}_\pi \sum_{h \in H_3} \sum_{x \in Q(w_h)} \Pr[x \in B_{\pi^{-1}(j)} | F_h] \Pr[F_h | w_1 \notin W_{\text{legal}}]. \end{aligned}$$

When we average  $\pi$  we get that  $\Pr[x \in B_{\pi^{-1}(j)} | F_h] = \frac{1}{s-j+1}$ , and hence

$$\Pr[B_j \text{ is double critical} | w_1 \notin W_{\text{legal}}] \leq \frac{1}{s-j+1} \sum_{h \in H_3} \sum_{x \in Q(w_h)} \Pr[F_h | w_1 \notin W_{\text{legal}}] = \frac{q}{s-j+1}.$$

We obtained the bound

$$\begin{aligned} &\Pr[B_j \text{ is double critical} \wedge w_1 \notin W_{\text{legal}}] \\ &= \Pr[B_j \text{ is double critical} | w_1 \notin W_{\text{legal}}] \Pr[w_1 \notin W_{\text{legal}}] \\ &\leq \frac{q}{s-j+1} \cdot \frac{q^2}{s-j+1} = \frac{q^3}{(s-j+1)^2}. \end{aligned}$$

Combining the two bounds we obtained, we conclude that

$$\begin{aligned} &\Pr[B_j \text{ is double critical}] \\ &= \Pr[B_j \text{ is double critical} \wedge w_1 \in W_{\text{legal}}] + \Pr[B_j \text{ is double critical} \wedge w_1 \notin W_{\text{legal}}] \\ &\leq \frac{q^2}{(s-j+1)^2} + \frac{q^3}{(s-j+1)^2} \leq \frac{2q^3}{(s-j+1)^2}. \end{aligned} \quad \square$$

**Claim 4.12.** *Let  $\mathcal{C}$  be such that  $\neg E_1, \neg E_2$  hold. Sample  $f_{j-1} | \mathcal{C}$ . Then*

$$\Pr[B_j \text{ is double critical for } f_{j-1} | \mathcal{C}] \geq (\gamma/b)^2.$$

*In particular,*

$$\Pr[B_j \text{ is double critical for } f_{j-1} | \neg E_1, \neg E_2] \geq (\gamma/b)^2.$$

*Proof.* Fix any  $\mathcal{C}$  such that  $\neg E_1, \neg E_2$  hold. By Claim 4.8 there are distinct  $x_1, x_2 \in B_j$  such that  $r(x_i) \geq \gamma/b$ . Note that if  $f_{j-1}(x_i) \in R(x_i)$  for both  $i = 1, 2$ , then  $B_j$  is double critical for  $f_{j-1}$ . As  $f_{j-1}(x_i)$  are sampled independently, we have

$$\Pr[B_j \text{ is double critical for } f_{j-1} | \mathcal{C}] \geq \Pr[f_{j-1}(x_i) \in R(x_i), i = 1, 2 | \mathcal{C}] \geq (\gamma/b)^2.$$

Combining Lemma 4.11 and Claim 4.12 gives a bound on the probability that both  $E_1, E_2$  don't hold.

**Corollary 4.13.**  $\Pr[\neg E_1 \wedge \neg E_2] \leq \frac{2q^3b^2}{\gamma^2(s-j+1)^2}$ .

*Proof.* Let  $F := [B_j \text{ is double critical for } f_{j-1}]$ . We have

$$\Pr[\neg E_1 \wedge \neg E_2] = \frac{\Pr[\neg E_1 \wedge \neg E_2 \wedge F]}{\Pr[F|\neg E_1 \wedge \neg E_2]} \leq \frac{\Pr[F]}{\Pr[F|\neg E_1, \neg E_2]} \leq \frac{2q^3b^2}{\gamma^2(s-j+1)^2}.$$

The claim then follows. □

From now on, we can prove Lemma 3.8. Appealing to equation(1) we have that

$$\Pr[X] \geq \left(1 - \frac{\gamma q}{j}\right) \Pr[Y] - \frac{2q^3b^2}{\gamma^2(s-j+1)^2} \geq \Pr[Y] - \left(\frac{\gamma q}{j} + \frac{2q^3b^2}{\gamma^2(s-j+1)^2}\right). \quad (2)$$

Let us denote

$$\varepsilon = \frac{\gamma q}{j} + \frac{2q^3b^2}{\gamma^2(s-j+1)^2}.$$

We now choose  $\gamma$  to minimize  $\varepsilon$ . Let us assume (as we have) that  $\delta s \leq j \leq (1-\delta)s$  for some absolute constant  $\delta > 0$ . Then

$$\varepsilon \leq \frac{\gamma q}{\delta s} + \frac{2q^3b^2}{\gamma^2\delta^2s^2}.$$

We choose  $\gamma = (2q^2b^2/\delta s)^{1/3}$  to equate the two terms, so that

$$\varepsilon \leq \frac{4q^{5/3}b^{2/3}}{\delta^{4/3}s^{4/3}}.$$

In particular, as we choose  $b, \delta$  to be absolute constants, we have  $\varepsilon \leq O(q^{5/3}/s^{4/3})$ .

## 5 Conclusions and open problems

In this paper, we studied impossibility of reversing entropy in black-box constructions. An obvious question that remains open is whether our result can be extended to the computational setting, given some complexity assumptions. Note that if we assume that  $P = NP$  then  $P = \text{NISZK} = \text{SZK} = \text{NP}$ .

Besides considering the relationship between NISZK and SZK, it is also interesting to explore relationships between other non-computational zero-knowledge proof systems. Concretely, what are the relationships between NIPZK and PZK, the perfect statistical analogs of NISZK and SZK, and the statistical versions. In particular, Malka [Mal08] gave a complete problem for NIPZK, which can be a good starting point to apply the techniques developed in this paper and separate NISZK and NIPZK.

## References

- [BDMP91] Manuel Blum, Alfredo De Santis, Silvio Micali, and Giuseppe Persiano. “Noninteractive zero-knowledge”. In: *SIAM Journal on Computing* 20.6 (1991), pp. 1084–1118 (cit. on p. 3).
- [BFM88] Manuel Blum, Paul Feldman, and Silvio Micali. “Non-interactive zero-knowledge and its applications”. In: *Proceedings of the twentieth annual ACM symposium on Theory of computing*. ACM. 1988, pp. 103–112 (cit. on p. 3).
- [DDP94] Alfredo De Santis, Giovanni Di Crescenzo, and Giuseppe Persiano. “The knowledge complexity of quadratic residuosity languages”. In: *Theoretical Computer Science* 132.1-2 (1994), pp. 291–317 (cit. on p. 3).
- [DDP97] Alfredo De Santis, Giovanni Di Crescenzo, and Pino Persiano. “Randomness-efficient non-interactive zero knowledge”. In: *International Colloquium on Automata, Languages, and Programming*. Springer. 1997, pp. 716–726 (cit. on p. 3).
- [DSDCPY98] Alfredo De Santis, Giovanni Di Crescenzo, Giuseppe Persiano, and Moti Yung. “Image density is complete for non-interactive-SZK”. In: *International Colloquium on Automata, Languages, and Programming*. Springer. 1998, pp. 784–795 (cit. on p. 3).
- [GGM86] Oded Goldreich, Shafi Goldwasser, and Silvio Micali. “How to construct random functions”. In: *Journal of the ACM (JACM)* 33.4 (1986), pp. 792–807 (cit. on p. 4).
- [GMR89] Shafi Goldwasser, Silvio Micali, and Charles Rackoff. “The knowledge complexity of interactive proof systems”. In: *SIAM Journal on computing* 18.1 (1989), pp. 186–208 (cit. on p. 2).
- [GMR98] Rosario Gennaro, Daniele Micciancio, and Tal Rabin. “An efficient non-interactive statistical zero-knowledge proof system for quasi-safe prime products”. In: *Proceedings of the 5th ACM conference on Computer and communications security*. ACM. 1998, pp. 67–72 (cit. on p. 3).
- [GSV99] Oded Goldreich, Amit Sahai, and Salil Vadhan. “Can statistical zero knowledge be made non-interactive? or on the relationship of SZK and NISZK”. In: *Advances in Cryptology CRYPTO 99*. Springer. 1999, pp. 467–484 (cit. on pp. 3–5, 7).
- [GV99] Oded Goldreich and Salil Vadhan. “Comparing entropies in statistical zero knowledge with applications to the structure of SZK”. In: *Computational Complexity, 1999. Proceedings. Fourteenth Annual IEEE Conference on*. IEEE. 1999, pp. 54–73 (cit. on p. 2).
- [HHR06] Iftach Haitner, Danny Harnik, and Omer Reingold. “On the power of the randomized iterate”. In: *Annual International Cryptology Conference*. Springer. 2006, pp. 22–40 (cit. on p. 4).
- [HILL99] Johan Håstad, Russell Impagliazzo, Leonid A Levin, and Michael Luby. “A pseudorandom generator from any one-way function”. In: *SIAM Journal on Computing* 28.4 (1999), pp. 1364–1396 (cit. on p. 4).
- [HR05] Thomas Holenstein and Renato Renner. “One-way secret-key agreement and applications to circuit polarization and immunization of public-key encryption”. In: *Annual International Cryptology Conference*. Springer. 2005, pp. 478–493 (cit. on p. 5).

- [Mal08] Lior Malka. “How to achieve perfect simulation and a complete problem for non-interactive perfect zero-knowledge”. In: *Theory of Cryptography Conference*. Springer. 2008, pp. 89–106 (cit. on p. 20).
- [PV08] Chris Peikert and Vinod Vaikuntanathan. “Noninteractive statistical zero-knowledge proofs for lattice problems”. In: *Annual International Cryptology Conference*. Springer. 2008, pp. 536–553 (cit. on p. 3).
- [SV03] Amit Sahai and Salil Vadhan. “A complete problem for statistical zero knowledge”. In: *Journal of the ACM (JACM)* 50.2 (2003), pp. 196–249 (cit. on pp. 2, 4, 5).
- [Vad16] Salil Vadhan. “Personal communication”. In: 2016 (cit. on p. 5).