

# Isolating a Vertex via Lattices: Polytopes with Totally Unimodular Faces

Rohit Gurjar<sup>\*1</sup>, Thomas Thierauf<sup>†2</sup>, and Nisheeth K. Vishnoi<sup>3</sup>

<sup>1</sup>Tel Aviv University, Israel

<sup>2</sup>Aalen University, Germany

<sup>3</sup>École Polytechnique Fédérale de Lausanne (EPFL), Switzerland

## Abstract

We derandomize the famous *Isolation Lemma* by Mulmuley, Vazirani, and Vazirani for polytopes given by totally unimodular constraints. That is, we construct a weight assignment such that one vertex in such a polytope is *isolated*, i.e., there is a unique minimum weight vertex. Our weights are quasi-polynomially bounded and can be constructed in quasi-polynomial time. In fact, our isolation technique works even under the weaker assumption that every face of the polytope lies in an affine space defined by a totally unimodular matrix. This generalizes the recent derandomization results for bipartite perfect matching and matroid intersection.

We prove our result by associating a *lattice* to each face of the polytope and showing that if there is a totally unimodular kernel matrix for this lattice, then the number of near-shortest vectors in it is polynomially bounded. The proof of this latter geometric fact is combinatorial and follows from a polynomial bound on the number of near-shortest circuits in a regular matroid. This is the technical core of the paper and relies on Seymour's decomposition theorem for regular matroids. It generalizes an influential result by Karger on the number of minimum cuts in a graph to regular matroids. Both of our results, on lattices and matroids, should be of independent interest.

---

\*The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement number 257575 and from the Israel Science Foundation (grant number 552/16).

†Supported by DFG grant TH 472/4.

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Our Results</b>	<b>5</b>
2.1	Isolating a Vertex in a Polytope . . . . .	5
2.2	Short vectors in lattices associated to polytopes . . . . .	6
2.3	Short circuits in regular matroids . . . . .	6
<b>3</b>	<b>Isolation via the Lattices Associated to the Polytope: Proof of Theorem 2.4</b>	<b>7</b>
<b>4</b>	<b>Matroids</b>	<b>10</b>
4.1	Matroids preliminaries . . . . .	10
4.2	Seymour's Theorem and its variants . . . . .	12
<b>5</b>	<b>A Bound on the Number of Short Vectors in Lattices: Proof of Theorem 2.5</b>	<b>13</b>
<b>6</b>	<b>A Bound on the Number of Short Circuits in Regular Matroids: Proof of Theorem 2.6</b>	<b>15</b>
6.1	Base Case: Graphic and cographic matroids . . . . .	15
6.2	General regular matroids . . . . .	17
<b>A</b>	<b>Proof of Theorem 4.19</b>	<b>25</b>

# 1 Introduction

The Isolation Lemma by Mulmuley, Vazirani, and Vazirani [MVV87] states that for any given family of subsets of a ground set  $E$ , if we assign random weights (bounded in magnitude by  $\text{poly}(|E|)$ ) to the elements of  $E$  then, with high probability, the minimum weight set in the family is unique. Such a weight assignment is called an *isolating weight assignment*. The lemma was introduced in the context of parallel algorithms for the matching problem. Since then it has found numerous other applications: a reduction from CLIQUE to UNIQUE-CLIQUE [MVV87], NL/poly  $\subseteq \oplus\text{L}/\text{poly}$  [Wig94], NL/poly = UL/poly [RA00], an RNC-algorithm for linear matroid intersection [NSV94], and an RP-algorithm for disjoint paths [BH14]. In all these results, the Isolation Lemma is the only place where they need randomness (or advice string). Thus, if the Isolation Lemma can be derandomized, i.e., a polynomially bounded isolating weight assignment can be deterministically constructed, then so can the aforementioned results that rely on it. Unfortunately, it is easy to see that it is impossible to design a polynomially bounded isolating weight assignment for all possible families of subsets of  $E$ . This is because there are exponentially many subsets and the weights are polynomially bounded. For any polynomially bounded weight assignment, there will be two subsets of  $E$  with the same weights. These two subsets form a family where the weight assignment fails. Even a relaxed task – of constructing a poly-size list of poly-bounded weight functions such that for each family  $\mathcal{B} \subseteq 2^E$ , one of the weight functions in the list is isolating – is impossible. This can be shown via arguments involving the polynomial identity testing (PIT) problem. The PIT problem, that is another important consequence of derandomizing the Isolation Lemma, asks if an implicitly given multivariate polynomial is identically zero. Here, the family of sets one needs to consider comes from the family of monomials that have nonzero coefficients in the polynomial. In essence, construction of such a list would imply that there exists a small set of points in  $\mathbb{R}^E$  such that any  $|E|$ -variate polynomial of small degree is nonzero on at least one of the points in the set. One can rule this out by designing a small degree polynomial that vanishes on all the points in the set. Hence, a natural question is to solve the isolation question for families  $\mathcal{B}$ , that have a succinct representation.

In this work, we derandomize the Isolation Lemma for a large class of families via a geometric approach. For a family of sets  $\mathcal{B} \subseteq 2^E$ , define the polytope  $P(\mathcal{B}) \subseteq \mathbb{R}^E$  to be the convex hull of the characteristic vectors of the sets in  $\mathcal{B}$ . We show that, for  $m := |E|$ , there exists an  $m^{O(\log m)}$ -sized family of weight assignments on  $E$ , with weights bounded by  $m^{O(\log m)}$  that isolates any family  $\mathcal{B}$  whose corresponding polytope  $P(\mathcal{B})$  satisfies the following property: *the affine space spanned by any face of  $P(\mathcal{B})$  is parallel to the null space of some totally unimodular (TU) matrix*; see Theorem 2.3. Our weight construction is black-box in the sense that it does not need the description of the family or the polytope.

A large variety of polytopes satisfy this property and, as a consequence, have been extensively studied in combinatorial optimization. The simplest class of polytopes that satisfy this property is when the polytope  $P(\mathcal{B})$  has a description  $Ax \leq b$  with  $A$  being a TU matrix. Thus, a simple consequence of our main result is a resolution to the problem of derandomizing the isolation lemma for polytopes with TU constraints raised in a recent work [ST17]. Further, our results significantly generalize recent work for the family of perfect matchings in a bipartite graph [FGT16] and for the family of common bases of two matroids [GT17]. In the case of perfect matchings in bipartite graphs, the perfect matching polytope can be described by the incidence matrix of the given graph which is a TUM. In the matroid intersection problem, the constraints of the common base polytope are a rank bound on every subset of the ground set. These constraints, in general, do not form a TUM. However, for every face of the polytope there exist two laminar families of subsets that form a basis for the tight constraints of the face. The incidence matrix for the union of two laminar

families is TU (see [Sch03b, Theorem 41.11]). Other examples of families whose polytopes are defined by TU constraints are vertex covers of a bipartite graph, independent sets of a bipartite graph, edge covers of a bipartite graph. Since the constraint matrix defining the polytope (or any of its face) itself does not have to be TU for it to satisfy the condition above, the condition required in Theorem 2.3 on the polytope  $P(\mathcal{B})$  is quite weak and is well studied. Schrijver [Sch03a, Theorem 5.35] shows that this condition is sufficient to prove that the polytope is *box-totally dual integral*. The second volume of Schrijver’s book [Sch03b] gives an excellent overview on polytopes that satisfy the condition of Theorem 2.3:

- up hull of an  $r$ -arborescence polytope [Sch03b, Section 52.4]
- $R - S$  bibranching polytope [Sch03b, Section 54.6]
- directed cut cover polytope [Sch03b, Section 55.2]
- submodular flow polyhedron [Sch03b, Theorem 60.1]
- lattice polyhedron [Sch03b, Theorem 60.4]
- For a submodular set function  $f$  on a set  $E$ , the polytope defined by

$$\sum_{e \in S} x_e \leq f(S) \text{ for } S \subseteq E \quad [\text{Sch03b, Section 44.3}].$$

Schrijver [Sch03b] also shows that the condition required by Theorem 2.3 holds for other polytopes defined via submodular set functions [Sch03b, (46.1), (48.1), (48.23)], submodular and supermodular set functions [Sch03b, (46.13), (46.28), (46.29)], submodular functions on a lattice family [Sch03b, (49.3), (49.12)], intersecting submodular functions [Sch03b, (49.33), (49.39)], and intersecting supermodular functions [Sch03b, (49.53)].

Our starting point is a reformulation of the approach for bipartite perfect matching and matroid intersection [FGT16, GT17] in terms of certain *lattices* associated to polytopes. For each face  $F$  of  $P(\mathcal{B})$ , we consider the lattice  $L_F$  of all integer vectors parallel to  $F$ . We show that, if for each face  $F$  of  $P(\mathcal{B})$ , the number of near-shortest vectors in  $L_F$  is polynomially bounded then we can construct an isolating weight assignment for  $\mathcal{B}$  with quasi-polynomially bounded weights; see Theorem 2.4. Our main technical contribution is to give a polynomial bound on the number of vectors whose length is less than  $3/2$  times that of the shortest vector in  $L_F$ , when this lattice is the set of integral vectors in the null space of a TUM; see Theorem 2.5. This is in contrast to general lattices where the number of such near-shortest vectors could be exponential in the dimension.

The above result can be reformulated using the language of matroid theory: the number of near-shortest circuits in a regular matroid is polynomially bounded; see Theorem 2.6. In fact we show how Theorem 2.5 can be deduced from Theorem 2.6. One crucial ingredient in the proof of Theorem 2.6 is Seymour’s decomposition theorem for regular matroids [Sey80]. Theorem 2.6 answers a question raised by Subramanian [Sub95] and can be viewed as a generalization of known results in the case of graphic and cographic matroids, that is, the number of near-minimum length cycles in a graph is polynomially bounded (see [TK92, Sub95]) and the number of near-mincuts in a graph is polynomially bounded (see [Kar93]).

Thus, not only do our results make significant progress in derandomizing the isolation lemma for combinatorial polytopes, our structural results about the number of near-shortest vectors in lattices and near-shortest circuits in matroids should be of independent interest and raise the question: to what extent are they generalizable?

## 2 Our Results

In this section we explain and state our main theorems. The proofs are given in the subsequent sections.

### 2.1 Isolating a Vertex in a Polytope

For a set  $E$  and a weight function  $w: E \rightarrow \mathbb{Z}$ , we define the extension of  $w$  to any set  $S \subseteq E$  by

$$w(S) := \sum_{e \in S} w(e).$$

Let  $\mathcal{B} \subseteq 2^E$  be a family of subsets of  $E$ . A weight function  $w: E \rightarrow \mathbb{Z}$  is called *isolating for  $\mathcal{B}$* , if the minimum weight set in  $\mathcal{B}$  is unique. In other words, the set  $\arg \min_{S \in \mathcal{B}} w(S)$  is unique. The Isolation Lemma of Mulmuley, Vazirani, and Vazirani [MVV87] asserts that a random weight function is isolating with a good probability for any  $\mathcal{B}$ .

**Lemma 2.1** (Isolation Lemma). *Let  $E$  be a set,  $|E| = m$ , and let  $w: E \rightarrow \{1, 2, \dots, 2m\}$  be a random weight function, where for each  $e \in E$ , the weight  $w(e)$  is chosen uniformly and independently at random. Then for any family  $\mathcal{B} \subseteq 2^E$ , the weight function  $w$  is isolating with probability at least  $1/2$ .*

The task of derandomizing the Isolation Lemma asks for a deterministic construction of an isolating weight function with weights polynomially bounded in  $m = |E|$ . Here, we view the isolation question for  $\mathcal{B}$  as an isolation over a corresponding polytope  $P(\mathcal{B})$ , which is defined as follows. For a set  $S \subseteq E$ , its characteristic vector  $x^S \in \mathbb{R}^m$  is defined as

$$x_e^S := \begin{cases} 1, & \text{if } e \in S, \\ 0, & \text{otherwise.} \end{cases}$$

For any family of sets  $\mathcal{B} \subseteq 2^E$ , the polytope  $P(\mathcal{B}) \subseteq \mathbb{R}^m$  is defined as the convex hull of the characteristic vectors of the sets in  $\mathcal{B}$ ,

$$P(\mathcal{B}) := \text{conv} \{ x^S \mid S \in \mathcal{B} \}.$$

Note that  $P(\mathcal{B})$  is contained in the  $m$ -dimensional unit hypercube.

The isolation question for a family  $\mathcal{B}$  is equivalent to constructing a weight vector  $w \in \mathbb{Z}^E$  such that  $\langle w, x \rangle$  has a unique minimum over  $P(\mathcal{B})$ . The property we need for our isolation approach is in terms of a totally unimodular matrix.

**Definition 2.2** (Totally unimodular matrix). *A matrix  $A \in \mathbb{R}^{n \times m}$  is said to be totally unimodular (TU), if every square submatrix has determinant 0 or  $\pm 1$ .*

Our main theorem gives an efficient quasi-polynomial isolation for a family  $\mathcal{B}$  when each face of the polytope  $P(\mathcal{B})$  lies in the affine space defined by a TU matrix.

**Theorem 2.3 (Main Result).** *Given a set  $E$  with  $|E| = m$ , we can construct a set  $W$  of  $m^{O(\log m)}$  weight assignments on  $E$  with weights bounded by  $m^{O(\log m)}$  with the following property: Let  $\mathcal{B} \subseteq 2^E$  be a family of sets. Suppose that for any face  $F$  of the polytope  $P(\mathcal{B})$ , there exists a TU matrix  $A_F \in \mathbb{R}^{n \times m}$  such that the affine space spanned by  $F$  is given by  $A_F x = b_F$  for some  $b_F \in \mathbb{R}^n$ . Then for the family  $\mathcal{B}$ , one of the weight assignments in  $W$  is isolating.*

## 2.2 Short vectors in lattices associated to polytopes

Our starting point towards proving Theorem 2.3 is a reformulation of the isolation approach for bipartite perfect matching and matroid intersection [FGT16, GT17]. For a set  $E$  and a family  $\mathcal{B} \subseteq 2^E$ , we define a lattice corresponding to each face of the polytope  $P(\mathcal{B})$ . The isolation approach works when this lattice has a small number of short vectors. For any face  $F$  of  $P(\mathcal{B})$ , consider the lattice of all integral vectors parallel to  $F$ ,

$$L_F := \{ v \in \mathbb{Z}^E \mid v = \alpha(x_1 - x_2) \text{ for some } x_1, x_2 \in F \text{ and } \alpha \in \mathbb{R} \}.$$

The length of the shortest nonzero vector of a lattice  $L$  is denoted by

$$\lambda(L) := \min \{ \|v\| \mid 0 \neq v \in L \},$$

where  $\|\cdot\|$  denotes the  $\ell_1$ -norm. We prove the following theorem that asserts that if, for all faces  $F$  of  $P(\mathcal{B})$ , the number of near-shortest vectors in  $L_F$  is small, then we can efficiently isolate a vertex in  $P(\mathcal{B})$ .

**Theorem 2.4.** *Let  $E$  be a set,  $|E| = m$ , and  $\mathcal{B} \subseteq 2^E$  be a family such that there exists a constant  $c > 1$ , such that for any face  $F$  of polytope  $P(\mathcal{B})$ , we have*

$$|\{ v \in L_F \mid \|v\| < c \lambda(L_F) \}| \leq m^{O(1)}.$$

*Then one can construct a set of  $m^{O(\log m)}$  weight functions with weights bounded by  $m^{O(\log m)}$  such that at least one of them is isolating for  $\mathcal{B}$ .*

The main ingredient of the proof of Theorem 2.3 is to show that the hypothesis of Theorem 2.4 is true when the lattice  $L_F$  is the set of all integral vectors in the nullspace of a TU matrix. For any  $n \times m$  matrix  $A$  we define a lattice  $L(A)$  as follows:

$$L(A) := \{ v \in \mathbb{Z}^m \mid Av = 0 \}.$$

**Theorem 2.5.** *For an  $n \times m$  TU matrix  $A$ , let  $\lambda := \lambda(L(A))$ . Then*

$$|\{ v \in L(A) \mid \|v\| < 3/2 \lambda \}| = O(m^5).$$

Theorem 2.5 together with Theorem 2.4 implies Theorem 2.3.

*Proof of Theorem 2.3.* Let  $F$  be a face of the polytope  $P(\mathcal{B})$  and let  $A_F$  be the TU matrix associated with  $F$ . Thus  $A_F x = b_F$  defines the affine span of  $F$ . In other words, the set of vectors parallel to  $F$  is precisely the solution set of  $A_F x = 0$  and the lattice  $L_F$  is given by  $L(A_F)$ . Theorem 2.5 implies the hypothesis of Theorem 2.4 for any  $L_F = L(A_F)$ , when the matrix  $A_F$  is TU.  $\square$

## 2.3 Short circuits in regular matroids

The proof of Theorem 2.5 is combinatorial and uses the language and results from matroid theory. We refer the reader to Section 4 for preliminaries on matroids; here we just recall a few basic definitions. A matroid is said to be *represented by a matrix*  $A$ , if its ground set is the column set of  $A$  and its independent sets are the sets of linearly independent columns of  $A$ . A matroid represented by a TU matrix is said to be a *regular matroid*. A *circuit* of a matroid is a minimal dependent set. The following is one of our main results which gives a bound on the number of short circuits in a regular matroid, which, in turn, implies Theorem 2.5. Instead of the circuit size, we consider the weight of a circuit and present a more general result.

**Theorem 2.6.** *Let  $M = (E, \mathcal{I})$  be a regular matroid with  $m = |E| \geq 2$  and  $w: E \rightarrow \mathbb{N}$  be a weight function. Suppose  $M$  does not have any circuit  $C$  with  $w(C) < r$ , for some number  $r$ . Then*

$$|\{C \mid C \text{ circuit in } M \text{ and } w(C) < 3r/2\}| \leq 240m^5.$$

**Remark 2.7.** *An extension of this result would be to give a polynomial bound on the number of circuits of weight at most  $\alpha r$  for any constant  $\alpha$ . Our current proof does not extend to this setting. This will be the subject of an upcoming work.*

## Organization of the rest of the paper

In Section 3, we present a proof of Theorem 2.4. In Section 4.1, we present a basic introduction to matroids. In Section 4.2, we describe some well-known properties of regular matroids which will be key to the proof of Theorem 2.6. Section 5 describes how Theorem 2.5 follows from Theorem 2.6. Finally, in Section 6, we prove Theorem 2.6.

## 3 Isolation via the Lattices Associated to the Polytope: Proof of Theorem 2.4

This section is dedicated to a proof of Theorem 2.4. That is, we give a construction of an isolating weight assignment for a family  $\mathcal{B} \subseteq 2^E$  assuming that for each face  $F$  of the corresponding polytope  $P(\mathcal{B})$ , the lattice  $L_F$  has small number of short vectors. First, let us see how the isolation question for a family  $\mathcal{B}$  translates in the polytope setting. For any weight function  $w: E \rightarrow \mathbb{Z}$ , we view  $w$  as a function on  $P(\mathcal{B})$ . That is, we define an extension of the weight function  $w$  to  $\mathbb{R}^E$ . For  $x \in \mathbb{R}^E$ ,

$$w(x) := \langle w, x \rangle = \sum_{e \in E} w(e) x_e.$$

That is, we can consider  $w$  as a vector in  $\mathbb{Z}^E$ , and the weight of a vector  $x$  is the inner product with  $w$ . Note that  $\langle w, x^B \rangle = w(B)$ , for any  $B \subseteq E$ . Thus, we get the following.

**Claim 3.1.** *A weight function  $w: E \rightarrow \mathbb{Z}$  is isolating for a family  $\mathcal{B}$  if and only if  $w(x)$  has a unique minimum over the polytope  $P(\mathcal{B})$ .*

Observe that for any  $w: E \rightarrow \mathbb{Z}$ , the points that minimize  $w(x) = \langle w, x \rangle$  in  $P(\mathcal{B})$  will form a face of the polytope  $P(\mathcal{B})$ . The idea is to build the isolating weight function in rounds. In every round, we slightly modify the current weight function to get a smaller minimizing face. Our goal is to significantly reduce the dimension of the minimizing face in every round. We stop when we reach a zero-dimensional face, i.e., we have a unique minimum weight point in  $P(\mathcal{B})$ .

The following claim asserts that if we modify the current weight function on a small scale, then the new minimizing face will be a subset of the current minimizing face. In the following, we will denote the size of set  $E$  by  $m$ .

**Claim 3.2.** *Let  $w: E \rightarrow \mathbb{Z}$  be a weight function and  $F$  be the face of  $P(\mathcal{B})$  that minimizes  $w$ . Let  $w': E \rightarrow \{0, 1, \dots, N-1\}$  be another weight function and let  $F'$  be the face that minimizes the combined weight function  $mNw + w'$ . Then  $F' \subseteq F$ .*

*Proof.* Consider any vertex  $x \in F'$ . We show that  $x \in F$ . By definition of  $F'$ , for any vertex  $y \in P(\mathcal{B})$  we have

$$\langle mNw + w', x \rangle \leq \langle mNw + w', y \rangle.$$

In other words,

$$\langle mN w + w', x - y \rangle \leq 0. \quad (1)$$

Since  $x$  and  $y$  are vertices of  $\mathcal{P}(\mathcal{B})$ , we have  $x, y \in \{0, 1\}^m$ . Thus,  $|\langle w', x - y \rangle| < mN$ . On the other hand, if  $|\langle mN w, x - y \rangle|$  is nonzero then it is at least  $mN$  and thus dominates  $|\langle w', x - y \rangle|$ . Hence, for (1) to hold, it must be that

$$\langle mN w, x - y \rangle \leq 0.$$

It follows that  $\langle w, x \rangle \leq \langle w, y \rangle$ , and therefore  $x \in F$ .  $\square$

Thus, in each round, we will add a new weight function to the current function using a smaller scale and try to get a subface with significantly smaller dimension. Henceforth,  $N$  will be sufficiently large number bounded by  $\text{poly}(m)$ .

**Definition 3.3.** For a face  $F$  of the polytope  $P(\mathcal{B})$ , a vector  $v \in \mathbb{R}^E$  is parallel to  $F$  if  $v = \alpha(x_1 - x_2)$  for some  $x_1, x_2 \in F$  and  $\alpha \in \mathbb{R}$ .

Next we show that a weight vector  $w$  is orthogonal to the face  $F$  it minimizes.

**Claim 3.4.** Let  $F$  be the face of  $P(\mathcal{B})$  minimizing a weight function  $w$ . Let  $v$  be a vector parallel to  $F$ . Then  $\langle w, v \rangle = 0$ .

*Proof.* Since  $v$  is parallel to  $F$ , we have  $v = \alpha(x_1 - x_2)$ , for some  $x_1, x_2 \in F$  and  $\alpha \in \mathbb{R}$ . Hence,

$$\langle w, v \rangle = \langle w, \alpha(x_1 - x_2) \rangle = 0.$$

The last equality holds because  $x_1, x_2 \in F$  and thus,  $\langle w, x_1 \rangle = \langle w, x_2 \rangle$ .  $\square$

Let  $F_0$  be the face minimizing the current weight function  $w_0$ . Let  $v$  be a vector parallel to  $F_0$ . Now, we choose a new weight function  $w' \in \{0, 1, \dots, N-1\}^E$  such that

$$\langle w', v \rangle \neq 0.$$

Let us define  $w_1 := mN w_0 + w'$  and let  $F_1$  be the face minimizing  $w_1$ . Clearly,  $\langle w_1, v \rangle \neq 0$  and thus, by Claim 3.4,  $v$  is not parallel to  $F_1$ . This implies that  $F_1$  is strictly contained in  $F_0$ . To ensure that  $F_1$  is *significantly* smaller than  $F_0$ , we choose many vectors parallel to  $F_0$ , say  $v_1, v_2, \dots, v_k$ , and construct a weight function  $w'$  such that for all  $i \in [k]$ , we have  $\langle w', v_i \rangle \neq 0$ . The following well-known lemma actually constructs a list of weight vectors such that one of them has the desired property (see [FKS84]).

**Lemma 3.5.** Given  $m, k, t$ , let  $q = mk \log t$ . In time  $\text{poly}(m, k, \log t)$  one can construct a set of weight vectors  $w_1, w_2, \dots, w_q \in \{0, 1, 2, \dots, q\}^m$  such that for any set of nonzero vectors  $v_1, v_2, \dots, v_k \in \{-(t-1), \dots, 0, 1, \dots, t-1\}^m$  there exists a  $j \in [q]$  such that for all  $i \in [k]$  we have  $\langle w_j, v_i \rangle \neq 0$ .

*Proof.* First define  $w := (1, t, t^2, \dots, t^{m-1})$ . Clearly,  $\langle w, v_i \rangle \neq 0$  for each  $i$ , because each coordinate of  $v_i$  is less than  $t$  in absolute value. To get a weight vector with small coordinates, we go modulo small numbers. We consider the following weight vectors  $w_j$  for  $1 \leq j \leq q$ :

$$w_j := w \bmod j.$$

We claim that this set of weight vectors has the desired property. We know that

$$W = \prod_{i=1}^k \langle w, v_i \rangle \neq 0.$$

Note that the product  $W$  is bounded by  $t^{mk}$ . On the other hand, it is known that  $\text{lcm}(2, 3, \dots, q) > 2^q = t^{mk}$  for all  $q \geq 7$  [Nai82]. Thus, there must exist a  $2 \leq j \leq q$  such that  $j$  does not divide  $W$ . In other words, for all  $i \in [k]$

$$\langle w, v_i \rangle \not\equiv 0 \pmod{j}$$

which is the desired property.  $\square$

There are two things to note about this lemma: (i) it is black-box in the sense that we do not need to know the set of vectors  $\{v_1, v_2, \dots, v_k\}$ . (ii) We do not know a priori which function will work in the given set of functions. So, one has to try all possibilities.

The lemma tells us that we can ensure  $\langle w', v \rangle \neq 0$  for polynomially many vectors  $v$  whose coordinates are polynomially bounded. It is the foundation of our strategy that was also used previously [FGT16, GT17]. Below, we formally give the weight construction. Recall that for a face  $F$ , the lattice  $L_F$  is the set of all integral vectors parallel to  $F$ .

To prove Theorem 2.4, let  $c$  be the constant in the assumption of the theorem. Let  $N = m^{O(1)}$  be a large enough number and  $p = \lfloor \log_c(m+1) \rfloor$ . Let  $w_0: E \rightarrow \mathbb{Z}$  be a weight function such that  $\langle w_0, v \rangle \neq 0$  for all nonzero  $v \in \mathbb{Z}^E$  with  $\|v\| < c$ . For  $i = 1, 2, \dots, p$ , define

$F_{i-1}$ : the face of  $P(\mathcal{B})$  minimizing  $w_{i-1}$

$w'_i$ : a weight vector in  $\{0, 1, \dots, N-1\}^E$  such that  $\langle w'_i, v \rangle \neq 0$  for all nonzero  $v \in L_{F_{i-1}}$  with  $\|v\| < c^{i+1}$ .

$w_i$ :  $mNw_{i-1} + w'_i$ .

Observe that  $F_i \subseteq F_{i-1}$ , for each  $i$  by Claim 3.2. Hence, also for the associated lattices we have  $L_{F_i} \subseteq L_{F_{i-1}}$ . We show that the lengths of the shortest vectors in the  $L_{F_i}$ 's grow exponentially in  $i$ .

**Claim 3.6.** *For  $i = 0, 1, 2, \dots, p$ , we have  $\lambda(L_{F_i}) \geq c^{i+1}$ .*

*Proof.* Consider a nonzero vector  $v \in L_{F_i}$ . By Claim 3.4, weight  $w_i$  is orthogonal to  $F_i$ . Therefore we have

$$\langle w_i, v \rangle = mN\langle w_{i-1}, v \rangle + \langle w'_i, v \rangle = 0. \quad (2)$$

Since  $v$  is in  $L_{F_i}$ , it is also in  $L_{F_{i-1}}$  and again by Claim 3.4, we have  $\langle w_{i-1}, v \rangle = 0$ . Together with (2) we conclude that  $\langle w'_i, v \rangle = 0$ . By the definition of  $w'_i$ , this implies that  $\|v\| \geq c^{i+1}$ .  $\square$

Finally we argue that  $w_p$  is isolating.

**Claim 3.7.** *The face  $F_p$  is a point.*

*Proof.* Let  $y_1, y_2 \in F_p$  be vertices. Then  $y_1 - y_2 \in L_{F_p}$  and  $\|y_1 - y_2\| \leq m < c^{p+1}$ . By Claim 3.6, we have that  $y_1 - y_2$  must be zero, i.e.,  $y_1 = y_2$ .  $\square$

**Bound on the weights.** To bound the weights of  $w_p$ , we bound  $w'_i$  for each  $i$ . By Claim 3.6, we have  $\lambda(L_{F_{i-1}}) \geq c^i$ , for each  $1 \leq i \leq p$ . The hypothesis of Theorem 2.4 implies

$$|\{v \in L_{F_{i-1}} \mid \|v\| < c^{i+1}\}| \leq m^{O(1)}.$$

Recall that we have to ensure  $\langle w'_i, v \rangle \neq 0$  for all nonzero vectors  $v$  in the above set. We apply Lemma 3.5 with  $k = m^{O(1)}$ . For parameter  $t$ , note that as  $\|v\| < c^{i+1} \leq c^{p+1} \leq c(m+1)$ , each coordinate of  $v$  is less than  $c(m+1)$  and therefore  $t \leq c(m+1)$ . Thus, we get  $w'_i$  with weights bounded by  $m^{O(1)}$ . Therefore the weights in  $w_p$  are bounded by  $m^{O(p)} = m^{O(\log m)}$ .

Recall that Lemma 3.5 actually gives a set of  $m^{O(1)}$  weight vectors for possible choices of  $w'_i$  and one of them has the desired property. Thus, we try all possible combinations for each  $w'_i$ . This gives us a set of  $m^{O(\log m)}$  possible choices for  $w_p$  such that one of them is isolating for  $\mathcal{B}$ . This proves Theorem 2.4.

## 4 Matroids

In Section 4.1, we recall some basic definitions and well-known facts about matroids (see, for example, [Oxl06,Sch03b]). In Section 4.2, we describe Seymour's decomposition theorem for regular matroids.

### 4.1 Matroids preliminaries

We start with some basic definitions.

**Definition 4.1** (Matroid). *A pair  $M = (E, \mathcal{I})$  is a matroid if  $E$  is a finite set and  $\mathcal{I}$  is a nonempty collection of subsets of  $E$  satisfying*

1. *if  $I \in \mathcal{I}$  and  $J \subseteq I$ , then  $J \in \mathcal{I}$ ,*
2. *if  $I, J \in \mathcal{I}$  and  $|I| < |J|$ , then  $I \cup \{z\} \in \mathcal{I}$ , for some  $z \in J \setminus I$ .*

*A subset  $I$  of  $E$  is said to be independent, if  $I$  belongs to  $\mathcal{I}$  and dependent otherwise. An inclusionwise maximal independent subset of  $E$  is a base of  $M$ . An inclusionwise minimal dependent set is a circuit of  $M$ .*

We define some special classes of matroids.

**Definition 4.2** (Linear, binary, and regular matroid). *A matroid  $M = (E, \mathcal{I})$  with  $m = |E|$  is linear or representable over some field  $\mathbb{F}$ , if there is a matrix  $A \in \mathbb{F}^{n \times m}$ , for some  $n$ , such that the collection of subsets of the columns of  $A$  that are linearly independent over  $\mathbb{F}$  is identical to  $\mathcal{I}$ .*

*A matroid  $M$  is binary, if  $M$  is representable over  $\text{GF}(2)$ . A matroid  $M$  is regular, if  $M$  is representable over every field.*

It is well known that regular matroids can be characterized in terms of TU matrices.

**Theorem 4.3** (See [Oxl06,Sch03b]). *A matroid  $M$  is regular if, and only if,  $M$  can be represented by a TU matrix over  $\mathbb{R}$ .*

Two special classes of regular matroids are graphic matroids and their duals, cographic matroids.

**Definition 4.4** (Graphic and cographic matroid). *A matroid  $M = (E, \mathcal{I})$  is said to be a graphic, if there is an undirected graph  $G = (V, E)$  whose edges correspond to the ground set  $E$  of  $M$ , such that  $I \in \mathcal{I}$  if and only if  $I$  forms a forest in  $G$ . By  $M(G)$  we denote the graphic matroid corresponding to  $G$ .*

*The dual of  $M$  is the matroid  $M^* = (E, \mathcal{I}^*)$  over the same ground set such that a set  $I \subseteq E$  is independent in  $M^*$  if and only if  $E \setminus I$  contains a base set of  $M$ . A cographic matroid is the dual of a graphic matroid.*

For  $G = (V, E)$ , we can represent  $M(G)$  by the vertex-edge incidence matrix  $A_G \in \{0, 1\}^{V \times E}$  (over  $\text{GF}(2)$ ),

$$A_G(v, e) = \begin{cases} 1 & \text{if } e \text{ is incident on } v, \\ 0 & \text{otherwise.} \end{cases}$$

**Definition 4.5** (Graph cut and cut-set). For a graph  $G = (V, E)$ , a cut is a partition  $(V_1, V_2)$  of  $V$  into two disjoint subsets. Any cut  $(V_1, V_2)$  uniquely determines a cut-set, the set of edges that have one endpoint in  $V_1$  and the other in  $V_2$ . The size of a cut is the number of edges in the corresponding cut-set. A minimum cut is one of minimum size.

**Fact 4.6.** Let  $G = (V, E)$  be a graph.

1. The circuits of the graphic matroid  $M(G)$  are exactly the simple cycles of  $G$ .
2. The circuits of the cographic matroid  $M^*(G)$  are exactly the inclusionwise minimal cut-sets of  $G$ .

The symmetric difference of two cycles in a graph is a disjoint union of cycles. The analogous statement is true for binary matroids.

**Fact 4.7.** Let  $M$  be binary. If  $C_1$  and  $C_2$  are circuits of  $M$ , then the symmetric difference  $C_1 \Delta C_2$  is a disjoint union of circuits.

To prove Theorem 2.6, we have to bound the number of short circuits in regular matroids. In Lemma 6.1, we start by providing such a bound for graphic and cographic matroids. The lemma is a variant of the following theorem that bounds the number of short cycles [Sub95] and the number of small cuts [Kar93] in a graph.

**Theorem 4.8.** Let  $G = (V, E)$  be a graph with  $m \geq 1$  edges and  $\alpha \geq 2$ .

1. If  $G$  has no cycles of length at most  $r$ , then the number of cycles in  $G$  of length at most  $\alpha r/2$  is bounded by  $(2m)^\alpha$  [Sub95].
2. If  $G$  has no cuts of size at most  $r$ , then the number of cuts in  $G$  of size at most  $\alpha r/2$  is bounded by  $m^\alpha$  [Kar93].

We define two operations on matroids.

**Definition 4.9** (Deletion, contraction, minor). Let  $M = (E, \mathcal{I})$  be a matroid and  $e \in E$ . The matroid obtained from  $M$  by deleting  $e$  is denoted by  $M \setminus e$ . Its independent sets are given by the collection  $\{I \in \mathcal{I} \mid e \notin I\}$ .

The matroid obtained by contracting  $e$  is denoted by  $M/e$ . Its independent sets are given by the collection  $\{I \subseteq E \setminus \{e\} \mid I \cup \{e\} \in \mathcal{I}\}$ .

A matroid obtained after a series of deletion and contraction operations on  $M$  is called a minor of  $M$ .

**Fact 4.10.** Let  $M = (E, \mathcal{I})$  be a matroid and  $e \in E$ .

1. The circuits of  $M \setminus e$  are those circuits of  $M$  that do not contain  $e$ .
2. The classes of regular matroids, graphic matroids, and cographic matroids are minor closed.

For a characterization of regular matroids, we will need a specific matroid  $R_{10}$ , first introduced by [Bix77]. It is a matroid, with 10 elements in the ground set, represented over  $GF(2)$  by the following matrix.

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

**Fact 4.11** ([Sey80]). Any matroid obtained by deleting some elements from  $R_{10}$  is a graphic matroid.

## 4.2 Seymour's Theorem and its variants

The main ingredient for the proof of Theorem 2.6 is a theorem of Seymour [Sey80, Theorem 14.3] that shows that every regular matroid can be constructed from piecing together three kinds of matroids – graphic matroids, cographic matroids, and the matroid  $R_{10}$ . This piecing together is done via matroid operations called 1-sum, 2-sum and 3-sum. These operations are defined for binary matroids.

**Definition 4.12** (Sum of two matroids [Sey80], see also [Oxl06]). *Let  $M_1 = (E_1, \mathcal{I}_1)$  and  $M_2 = (E_2, \mathcal{I}_2)$  be two binary matroids, and let  $S = E_1 \cap E_2$ . The sum of  $M_1$  and  $M_2$  is a matroid denoted by  $M_1 \triangle M_2$ . It is defined over the ground set  $E_1 \triangle E_2$  such that the circuits of  $M_1 \triangle M_2$  are minimal non-empty subsets of  $E_1 \triangle E_2$  that are of the form  $C_1 \triangle C_2$ , where  $C_i$  is a (possibly empty) disjoint union of circuits of  $M_i$ , for  $i = 1, 2$ .*

From the characterization of the circuits of a matroid [Oxl06, Theorem 1.1.4], it can be verified that the sum  $M_1 \triangle M_2$  is indeed a matroid.

We are only interested in three special sums:

**Definition 4.13** (1, 2, 3-sums). *Let  $M_1 = (E_1, \mathcal{I}_1)$  and  $M_2 = (E_2, \mathcal{I}_2)$  be two binary matroids and  $E_1 \cap E_2 = S$ . Let  $m_1 = |E_1|$ ,  $m_2 = |E_2|$ , and  $s = |S|$ . Let furthermore  $m_1, m_2 < |E_1 \triangle E_2| = m_1 + m_2 - 2s$ . The sum  $M_1 \triangle M_2$  is called a*

- 1-sum, if  $s = 0$ ,
- 2-sum, if  $s = 1$  and  $S$  is not a circuit of  $M_1, M_2, M_1^*$  or  $M_2^*$ ,
- 3-sum, if  $s = 3$  and  $S$  is a circuit of  $M_1$  and  $M_2$  that does not contain a circuit of  $M_1^*$  or  $M_2^*$ .

Note that the condition  $m_1, m_2 < m_1 + m_2 - 2s$  implies that

$$m_1, m_2 \geq 2s + 1 \tag{3}$$

To get an intuition, we describe how the sum operation looks like for graphic matroids. For two graphs  $G_1$  and  $G_2$ , their 2-sum  $G$  is obtained as follows: for an edge  $(u_1, v_1)$  in  $G_1$  and an edge  $(u_2, v_2)$  in  $G_2$ , identify the two edges, that is, identify  $u_1$  with  $u_2$  and  $v_1$  with  $v_2$ . Finally,  $G$  is the union of  $G_1$  and  $G_2$  where the edge  $(u_1, v_1) = (u_2, v_2)$  is removed. Here,  $S$  is the set containing the single edge  $(u_1, v_1) = (u_2, v_2)$ .

It is instructive to see how a cycle  $C$  in  $G$  looks like. Either  $C$  is a cycle in  $G_1$  or  $G_2$  that avoids nodes  $u_1, v_1, u_2, v_2$ , or it is a union of a path  $u_1 \rightsquigarrow v_1$  in  $G_1$  and a path  $v_2 \rightsquigarrow u_2$  in  $G_2$ . Equivalently, it is a symmetric difference  $C = C_1 \triangle C_2$ , for two cycles  $C_1$  in  $G_1$  and  $C_2$  in  $G_2$ , such that  $C_1$  and  $C_2$  both contain the common edge  $(u_1, v_1) = (u_2, v_2)$ . Analogously, the sum operation for two binary matroids is defined such that any circuit  $C$  of the sum  $M_1 \triangle M_2$  is either a circuit in  $M_1$  or  $M_2$  that avoids the elements in  $S$ , or it is  $C = C_1 \triangle C_2$ , for circuits  $C_1$  and  $C_2$  of  $M_1$  and  $M_2$ , respectively, such that both  $C_1$  and  $C_2$  contain a common element from  $S$ .

From the definition of  $M_1 \triangle M_2$  the following fact follows easily.

**Fact 4.14.** *Let  $C_i$  be a disjoint union of circuits of  $M_i$ , for  $i = 1, 2$ . If  $C_1 \triangle C_2$  is a subset of  $E_1 \triangle E_2$  then it is a disjoint union of circuits of  $M_1 \triangle M_2$ .*

In particular, it follows that for  $i = 1, 2$ , any circuit  $C_i$  of  $M_i$  with  $C_i \subseteq E_i \setminus S$  is a circuit of  $M_1 \triangle M_2$ . Further, for 1-sums, circuits are easy to characterize.

**Fact 4.15** (Circuits in a 1-sum). *If  $M$  is a 1-sum of  $M_1$  and  $M_2$  then any circuit of  $M$  is either a circuit of  $M_1$  or a circuit of  $M_2$ .*

Thus, if one is interested in the number of circuits, one can assume that the given matroid is not a 1-sum of two smaller matroids.

**Definition 4.16** (Connected matroid). *A matroid  $M$  is connected if it cannot be written as a 1-sum of two smaller matroids.*

A characterization of circuits in a 2-sum or 3-sum is not as easy. Seymour [Sey80, Lemma 2.7] provides a unique representation of the circuits for these cases.

**Lemma 4.17** (Circuits in a 2- or 3-sum, [Sey80]). *Let  $\mathcal{C}_1$  and  $\mathcal{C}_2$  be the sets of circuits of  $M_1$  and  $M_2$ , respectively. Let  $M$  be a 2- or 3-sum of  $M_1$  and  $M_2$ . For  $S = E_1 \cap E_2$ , we have  $|S| = 1$  or  $|S| = 3$ , respectively. Then for any circuit  $C$  of  $M$ , one of the following holds:*

1.  $C \in \mathcal{C}_1$  and  $S \cap C = \emptyset$ , or
2.  $C \in \mathcal{C}_2$  and  $S \cap C = \emptyset$ , or
3. there exist unique  $e \in S$ ,  $C_1 \in \mathcal{C}_1$  and  $C_2 \in \mathcal{C}_2$  such that

$$S \cap C_1 = S \cap C_2 = \{e\} \text{ and } C = C_1 \Delta C_2.$$

Seymour proved the following decomposition theorem for regular matroids.

**Theorem 4.18** (Seymour's Theorem, [Sey80]). *Every regular matroid can be obtained by means of 1-sums, 2-sums and 3-sums, starting from matroids that are graphic, cographic or  $R_{10}$ .*

However, to prove Theorem 2.6, we need a refined version of Seymour's theorem that was proved by Truemper [Tru98]. Seymour's theorem decomposes a regular matroid into a sum of two smaller regular matroids. Truemper showed that one of the two smaller regular matroids can be chosen to be graphic, cographic, or the  $R_{10}$  matroid. The theorem we write here slightly differs from the one by Truemper [Tru98, Lemma 11.3.18]. A proof of Theorem 4.19 can be found in Appendix A.

**Theorem 4.19** (Truemper's decomposition for regular matroids, [Tru98]). *Let  $M$  be a connected regular matroid, that is not graphic or cographic and is not isomorphic to  $R_{10}$ . Let  $\tilde{e}$  be a fixed element of the ground set of  $M$ . Then  $M$  is a 2-sum or 3-sum of  $M_1$  and  $M_2$ , where  $M_1$  is a graphic or cographic matroid, or a matroid isomorphic to  $R_{10}$  and  $M_2$  is a regular matroid that contains  $\tilde{e}$ .*

## 5 A Bound on the Number of Short Vectors in Lattices: Proof of Theorem 2.5

In this section, we show that Theorem 2.5 follows from Theorem 2.6. We define a circuit of a matrix and show that it is sufficient to upper bound the number of short circuits of a TU matrix. We argue that this, in turn, is implied by a bound on the number of short circuits of a regular matroid. Recall that for an  $n \times m$  matrix  $A$ , the lattice  $L(A)$  is defined as the set of integer vectors in its kernel,  $L(A) := \{v \in \mathbb{Z}^m \mid Av = 0\}$ .

**Definition 5.1** (Circuit). *For an  $n \times m$  matrix  $A$ , a vector  $u \in L(A)$  is a circuit of  $A$  if*

- there is no nonzero  $v \in L(A)$  with  $\text{supp}(v) \subsetneq \text{supp}(u)$ , and

- $\gcd(u_1, u_2, \dots, u_m) = 1$ .

We have the following properties of circuits. The first follows directly from the definition. The second one is well known (see [Onn10, Lemma 3.18]).

**Fact 5.2.** *Let  $A$  be a matrix.*

1. *If  $u$  is a circuit of  $A$ , then also  $-u$ .*
2. *Let  $A$  be TU. Then every circuit of  $A$  has its coordinates in  $\{-1, 0, 1\}$ .*

Now, we define a notion of conformality, which will allow us to show that any sufficiently small vector in  $L(A)$  is a circuit, when  $A$  is TU.

**Definition 5.3** (Conformal [Onn10]). *Let  $u, v \in \mathbb{R}^m$ . We say that  $u$  is conformal to  $v$ , denoted by  $u \sqsubseteq v$ , if  $u_i v_i \geq 0$  and  $|u_i| \leq |v_i|$ , for each  $1 \leq i \leq m$ .*

Observe that for vectors  $u$  and  $v$  with  $u \sqsubseteq v$ , we have

$$\|v - u\| = \|v\| - \|u\|. \quad (4)$$

The following lemma follows from [Onn10, Lemma 3.19].

**Lemma 5.4.** *Let  $A$  be a TU matrix. Then for any nonzero vector  $v \in L(A)$ , there is a circuit  $u$  of  $A$  that is conformal to  $v$ .*

We use the lemma to argue that any small enough vector in  $L(A)$  must be a circuit.

**Lemma 5.5.** *Let  $A$  be a TU matrix and let  $\lambda := \lambda(L(A))$ . Then any nonzero vector  $v \in L(A)$  with  $\|v\| < 2\lambda$  is a circuit of  $A$ .*

*Proof.* Let  $v \in L(A)$  be not a circuit of  $A$ . We show that  $\|v\| \geq 2\lambda$ .

By Lemma 5.4, there is a circuit  $u$  of  $A$  with  $u \sqsubseteq v$ . Since  $v$  is not a circuit,  $v - u \neq 0$ . By Equation (4), we have

$$\|v\| = \|v - u\| + \|u\|. \quad (5)$$

Since both  $u$  and  $v - u$  are nonzero vectors in  $L(A)$ , we have  $\|u\|, \|v - u\| \geq \lambda$ . Together with Equation (5) we get that  $\|v\| \geq 2\lambda$ .  $\square$

Recall that by Theorem 4.3, a matroid represented by a TU matrix is a regular matroid. The following lemma shows that the two definitions of circuits, 1) for TU matrices and 2) for regular matroids, coincide.

**Lemma 5.6.** *Let  $M = (E, \mathcal{I})$  be a regular matroid, represented by a TU matrix  $A$ . Then there is a one to one correspondence between the circuits of  $M$  and the circuits of  $A$  (up to change of sign).*

*Proof.* If  $u \in \mathbb{R}^E$  is a circuit of  $A$ , then the columns in  $A$  corresponding to the set  $\text{supp}(u)$  are minimally dependent. Thus, the set  $\text{supp}(u)$  is a circuit of matroid  $M$ .

In the other direction, a circuit  $C \subseteq E$  of matroid  $M$  is a minimal dependent set. Thus, the set of columns of  $A$  corresponding to  $C$  is minimally linear dependent. Hence, there are precisely two vectors  $u, -u \in L(A)$  with their support being  $C$ .  $\square$

To prove Theorem 2.5, let  $A$  be TU matrix. By Lemma 5.5, it suffices to bound the number of near-shortest circuits of  $A$ . By Lemma 5.6, the circuits of  $A$  and the circuits of the regular matroid  $M$  represented by  $A$ , coincide. Moreover, the size of a circuit of  $M$  is same as the  $\ell_1$ -norm of the corresponding circuit of  $A$ , as a circuit of  $A$  has its coordinates in  $\{-1, 0, 1\}$  by Fact 5.2. Now Theorem 2.5 follows from Theorem 2.6 when we define the weight of each element being 1.

## 6 A Bound on the Number of Short Circuits in Regular Matroids: Proof of Theorem 2.6

In this section, we prove our main technical tool: in a regular matroid, the number of circuits that have size close to a shortest circuit is polynomially bounded (Theorem 2.6). The proof argues along the decomposition provided by Theorem 4.19. First, we need to show a bound on the number of circuits for the two base cases – graphic and cographic matroids.

### 6.1 Base Case: Graphic and cographic matroids

We actually prove a lemma for graphic and cographic matroids that does more – it gives an upper bound on the number of circuits that contain a fixed element of the ground set. For a weight function  $w: E \rightarrow \mathbb{N}$  on the ground set, the weight of any subset  $C \subseteq E$  is defined as  $w(C) := \sum_{e \in C} w(e)$ .

**Lemma 6.1.** *Let  $M = (E, \mathcal{I})$  be a graphic or cographic matroid, where  $|E| = m \geq 2$ , and  $w: E \rightarrow \mathbb{N}$  be a weight function. Let  $R \subseteq E$  with  $|R| \leq 1$  (possibly empty) and  $r$  be a positive integer.*

*If there is no circuit  $C$  in  $M$  such that  $w(C) < r$  and  $C \cap R = \emptyset$ , then, for any integer  $\alpha \geq 2$ , the number of circuits  $C$  such that  $R \subseteq C$  and  $w(C) < \alpha r/2$  is at most  $(2(m - |R|))^\alpha$ .*

*Proof. Part 1:  $M$  graphic.* (See [TK92, Sub95] for a similar argument as in this case.) Let  $G = (V, E)$  be the graph corresponding to the graphic matroid  $M$ . By the assumption of the lemma, any cycle  $C$  in  $G$  such that  $C \cap R = \emptyset$  has weight  $w(C) \geq r$ . Consider a cycle  $C$  in  $G$  with  $R \subseteq C$  and  $w(C) < \alpha r/2$ . Let the edge sequence of the cycle  $C$  be  $(e_1, e_2, e_3, \dots, e_q)$  such that if  $R$  is nonempty then  $R = \{e_1\}$ . We choose  $\alpha$  edges of the cycle  $C$  as follows: Let  $i_1 = 1$  and for  $j = 2, 3, \dots, \alpha$ , define  $i_j$  to be the least index greater than  $i_{j-1}$  (if one exists) such that

$$\sum_{a=i_{j-1}+1}^{i_j} w(e_a) \geq r/2. \quad (6)$$

If such an index does not exist then define  $i_j = q$ . Removing the edges  $e_{i_1}, e_{i_2}, \dots, e_{i_\alpha}$  from  $C$  gives us  $\alpha$  paths: for  $j = 1, 2, \dots, \alpha - 1$

$$p_j := (e_{i_j+1}, e_{i_j+2}, \dots, e_{i_{j+1}-1}),$$

and

$$p_\alpha := (e_{i_\alpha+1}, e_{i_\alpha+2}, \dots, e_q).$$

Note that some of these paths might be empty. By the choice of  $i_j$  we know that  $w(p_j) < r/2$  for  $j = 1, 2, \dots, \alpha - 1$ . Combining (6) with the fact that  $w(C) < \alpha r/2$ , we obtain that  $w(p_\alpha) < r/2$ . We associate the ordered tuple of oriented edges  $(e_{i_1}, e_{i_2}, \dots, e_{i_\alpha})$  with the cycle  $C$ .

**Claim 6.2.** *For two distinct cycles  $C, C'$  in  $G$ , such that both contain  $R$  and  $w(C), w(C') < \alpha r/2$ , the two associated tuples (defined as above) are different.*

*Proof.* For the sake of contradiction, assume that the associated tuples are same for both the cycles. Thus,  $C$  and  $C'$  pass through  $(e_{i_1}, e_{i_2}, \dots, e_{i_\alpha})$  with the same orientation of these edges. Further, there are  $\alpha$  paths connecting them, say  $p_1, p_2, \dots, p_\alpha$  from  $C$  and  $p'_1, p'_2, \dots, p'_\alpha$  from  $C'$ . Since  $C$  and  $C'$  are distinct, for at least one  $j$ , it must be that  $p_j \neq p'_j$ . However, since the starting points and the end points of  $p_j$  and  $p'_j$  are same,  $p_j \cup p'_j$  contains a cycle  $C''$ . Moreover, since  $w(p_j), w(p'_j) < r/2$ , we can deduce that  $w(C'') < r$ . Finally, since neither of  $p_j$  and  $p'_j$  contain  $e_1$ , we get  $C'' \cap R = \emptyset$ . This is a contradiction.  $\square$

Since, each cycle  $C$  with  $w(C) < \alpha r/2$  and  $R \subseteq C$  is associated with a different tuple, the number of such tuples upper bounds the number of such cycles. We bound the number of tuples depending on whether  $R$  is empty or not.

- When  $R$  is empty, the number of tuples of  $\alpha$  oriented edges is at most  $(2m)^\alpha$ .
- When  $R = \{e_1\}$ , the number of choices for the rest of the  $\alpha - 1$  edges and their orientation is at most  $(2(m - 1))^{\alpha-1}$ .

**Part 2:  $M$  cographic.** Let  $G = (V, E)$  be the graph corresponding to the cographic matroid  $M$  and let  $n = |V|$ . Recall from Fact 4.6 that circuits in cographic matroids are inclusionwise minimal cut-sets in  $G$ . By the assumption of the lemma, any cut-set  $C$  in  $G$  with  $R \cap C = \emptyset$  has weight  $w(C) \geq r$ . Note that this implies that  $G$  is connected, and therefore  $m \geq n - 1$ . We want to give a bound on the number of cut-sets  $C \subseteq E$  such that  $w(C) < \alpha r/2$  and  $R \subseteq C$ .

We argue similar to the probabilistic construction of a minimum cut of Karger [Kar93]. The basic idea is to contract randomly chosen edges. *Contraction of an edge  $e = (u, v)$*  means that all edges between  $u$  and  $v$  are deleted and then  $u$  is identified with  $v$ . Note that we get a multi-graph that way: if there were two edges  $(u, w)$  and  $(v, w)$  before the contraction, they become two parallel edges after identifying  $u$  and  $v$ . The contracted graph is denoted by  $G/e$ . The intuition behind contraction is, that randomly chosen edges are likely to avoid the edges of a minimum cut.

The following algorithm implements the idea. It does  $k \leq n$  contractions in the first phase and then chooses a random cut within the remaining nodes of the contracted graph in the second phase that contains the edges of  $R$ . Note that any cut-set of the contracted graph is also a cut-set of the original graph.

SMALL CUT ( $G = (V, E), R, \alpha$ )

*Contraction*

- 1 **Repeat**  $k = n - \alpha - |R|$  times
- 2     **randomly choose**  $e \in E \setminus R$  with probability  $w(e)/w(E \setminus R)$
- 3      $G \leftarrow G/e$
- 4      $R \leftarrow R \cup \{\text{new parallel edges to the edges in } R\}$

*Selection*

- 5 Among all possible cut-sets  $C$  in the obtained graph  $G$  with  $R \subseteq C$ , choose one uniformly at random and return it.

Let  $C \subseteq E$  be a cut-set with  $w(C) < \alpha r/2$  and  $R \subseteq C$ . We want to give a lower bound on the probability that SMALL CUT outputs  $C$ .

Let  $G_0 = G$  and  $G_i = (V_i, E_i)$  be the graph after the  $i$ -th contraction, for  $i = 1, 2, \dots, k$ . Note that  $G_i$  has  $n_i = n - i$  nodes since each contraction decreases the number of nodes by 1. Let  $R_i$  denote the set  $R$  after the  $i$ -th contraction. That is, if  $R = \{e_1\}$ , then  $R_i$  contains all edges parallel to  $e_1$  in  $G_i$ . In case that  $R = \emptyset$ , also  $R_i = \emptyset$ . Note that in either case  $R_i \subseteq C$ , if no edge of  $C$  has been contracted till iteration  $i$ .

Conditioned on the event that no edge in  $C$  has been contracted in iterations 1 to  $i$ , the probability that an edge from  $C$  is contracted in the  $(i + 1)$ -th iteration is at most

$$w(C \setminus R_i)/w(E_i \setminus R_i).$$

We know that  $w(C \setminus R_i) \leq w(C) < \alpha r/2$ . For a lower bound on  $w(E_i \setminus R_i)$ , consider the graph  $G'_i$  obtained from  $G_i$  by contracting the edges in  $R_i$ . The number of nodes in  $G'_i$  will be  $n'_i = n - i - |R|$  and its set of edges will be  $E_i \setminus R_i$ . For any node  $v$  in  $G'_i$ , consider the set  $\delta(v)$  of edges incident

on  $v$  in  $G'_i$ . The set  $\delta(v)$  forms a cut-set in  $G'_i$  and also in  $G$ . Note that  $\delta(v) \cap R = \emptyset$ , as the edge in  $R$  has been contracted in  $G'_i$ . Thus, we can deduce that  $w(\delta(v)) \geq r$ . By summing this up for all nodes in  $G'_i$ , we obtain

$$w(E_i \setminus R_i) \geq r n'_i / 2.$$

Hence,

$$w(E_i \setminus R_i) \geq r(n - i - |R|) / 2.$$

Therefore the probability that an edge from  $C$  is contracted in the  $(i + 1)$ -th iteration is

$$\leq \frac{w(C \setminus R_i)}{w(E_i \setminus R_i)} \leq \frac{\alpha r / 2}{r(n - i - |R|) / 2} = \frac{\alpha}{n - i - |R|}.$$

This bound becomes greater than 1, when  $i > n - \alpha - |R|$ . This is the reason why we stop the contraction process after  $k = n - \alpha - |R|$  iterations.

The probability that no edge from  $C$  is contracted in any of the rounds is

$$\begin{aligned} &\geq \prod_{i=0}^{k-1} \left( 1 - \frac{\alpha}{n - i - |R|} \right) \\ &= \prod_{i=0}^{k-1} \left( 1 - \frac{\alpha}{k + \alpha - i} \right) \\ &= \prod_{i=0}^{k-1} \frac{k - i}{k + \alpha - i} \\ &= \frac{1}{\binom{k+\alpha}{k}} \\ &= \frac{1}{\binom{n-|R|}{\alpha}}. \end{aligned}$$

After  $n - \alpha - |R|$  contractions we are left with  $\alpha + |R|$  nodes. We claim that the number of possible cut-sets on these nodes that contain  $R$  is  $2^{\alpha-1}$ . In case when  $R = \emptyset$ , then the number of partitions of  $\alpha$  nodes into two sets is clearly  $2^{\alpha-1}$ . When  $R = \{e_1\}$ , then the number of partitions of  $\alpha + 1$  nodes, such that the endpoints of  $e_1$  are in different parts, is again  $2^{\alpha-1}$ . We choose one of these cuts randomly. Thus, the probability that  $C$  survives the *contraction* process and is also chosen in the *selection* phase is at least

$$\frac{1}{2^{\alpha-1} \binom{n-|R|}{\alpha}} \geq \frac{1}{(n - |R|)^\alpha}.$$

Note that in the end we get exactly one cut-set. Thus, the number of cut-sets  $C$  of weight  $< \alpha r / 2$  and  $R \subseteq C$  must be at most  $(n - |R|)^\alpha$ , which is bounded by  $(2(m - |R|))^\alpha$  because  $m \geq n - 1$ .  $\square$

## 6.2 General regular matroids

In this section, we prove our main result about regular matroids.

**Theorem** (Theorem 2.6). *Let  $M = (E, \mathcal{I})$  be a regular matroid with  $m = |E| \geq 2$  and  $w: E \rightarrow \mathbb{N}$  be a weight function. Suppose  $M$  does not have any circuit  $C$  such that  $w(C) < r$ , for some number  $r$ . Then*

$$|\{C \mid C \text{ circuit in } M \text{ and } w(C) < 3r/2\}| \leq 240 m^5.$$

*Proof.* The proof is by an induction on  $m$ , the size of the ground set. For the base case, let  $m \leq 10$ . There are at most  $2^m$  circuits in  $M$ . This number is bounded by  $240 m^5$ , for any  $2 \leq m \leq 10$ .

For the inductive step, let  $M = (E, \mathcal{I})$  be a regular matroid with  $|E| = m > 10$  and assume that the theorem holds for all smaller regular matroids. Note that  $M$  cannot be  $R_{10}$  since  $m > 10$ . We can also assume that matroid  $M$  is neither graphic nor cographic, otherwise the bound follows from Lemma 6.1. By Theorem 4.18, matroid  $M$  can be written as a 1-, 2-, or 3-sum of two regular matroids  $M_1 = (E_1, \mathcal{I}_1)$  and  $M_2 = (E_2, \mathcal{I}_2)$ . We define

$$\begin{aligned} S &:= E_1 \cap E_2, \\ s &:= |S|, \\ m_i &:= |E_i|, \text{ for } i = 1, 2, \\ \mathcal{C}_i &:= \{ C \mid C \text{ is a circuit of } M_i \}. \end{aligned}$$

In case that  $M$  is the 1-sum of  $M_1$  and  $M_2$ , we have  $S = \emptyset$ , and therefore  $m = m_1 + m_2$ . By Fact 4.15, the set of circuits of  $M$  is the union of the sets of circuits of  $M_1$  and  $M_2$ . From the induction hypothesis, we have that  $M_i$  has at most  $240 m_i^5$  circuits of weight less than  $3r/2$ , for  $i = 1, 2$ . For the number of such circuits in  $M$  we get the bound of

$$240 m_1^5 + 240 m_2^5 \leq 240 m^5.$$

This proves the theorem in case of a 1-sum. Hence, in the following it remains to consider the case that  $M$  cannot be written as a 1-sum. In other words, we may assume that  $M$  is connected (Definition 4.16).

Now we can apply Theorem 4.19 and assume that  $M$  is a 2- or 3-sum of  $M_1$  and  $M_2$ , where  $M_1$  is a graphic, cographic or the  $R_{10}$  matroid, and  $M_2$  is a regular matroid.

We define for  $i = 1, 2$  and  $e \in S$

$$\begin{aligned} \mathcal{C}_{i,e} &:= \{ C \mid C \text{ is a circuit of } M_i \text{ and } C_i \cap S = \{e\} \}, \\ M'_i &:= M_i \setminus S, \\ \mathcal{C}'_i &:= \{ C \mid C \text{ is a circuit of } M'_i \}. \end{aligned}$$

By Facts 4.10 and 4.11, matroid  $M'_1$  is graphic or cographic, and  $M'_2$  is regular. Recall from Lemma 4.17 that any circuit  $C$  of  $M$  can be uniquely written as  $C_1 \triangle C_2$  such that one of the following holds:

- $C_1 = \emptyset$  and  $C_2 \in \mathcal{C}'_2$ .
- $C_2 = \emptyset$  and  $C_1 \in \mathcal{C}'_1$ .
- $C_1 \in \mathcal{C}_{1,e}$ , and  $C_2 \in \mathcal{C}_{2,e}$ , for some  $e \in S$ .

Thus, we will view each circuit  $C$  of  $M$  as  $C_1 \triangle C_2$  and consider cases based on how the weight of  $C$  is distributed among  $C_1$  and  $C_2$ . Recall that the weight function  $w$  is defined on  $E = E_1 \triangle E_2$ . We extend  $w$  to a function on  $E_1 \cup E_2$  by defining

$$w(e) = 0, \text{ for } e \in S.$$

Now, for the desired upper bound, we will divide the set of circuits of  $M$  of weight less than  $3r/2$  into three cases.

**Case 1.**  $C_1 \in \mathcal{C}'_1$ .

**Case 2.**  $w(C_1) < r/2$ . This includes the case that  $C_1 = \emptyset$ .

**Case 3.**  $w(C_1) \geq r/2$  and  $C_2 \neq \emptyset$ .

In the following, we will derive an upper bound for the number of circuits in each of the three cases. Then the sum of these bounds will be an upper bound on the number of circuits in  $M$ . We will show that the sum is less than  $240m^5$ .

**Case 1:**  $C_1 \in \mathcal{C}'_1$

We have  $C_2 = \emptyset$  and  $C = C_1 \in \mathcal{C}'_1$ . That is, we need to bound the number of circuits of  $M'_1$ . Recall that any circuit of  $M'_1$  is also a circuit of  $M$ . Hence, we know there is no circuit  $C_1$  in  $M'_1$  with  $w(C_1) < r$ . Since  $M'_1$  is graphic or cographic, from Lemma 6.1, the number of circuits  $C_1$  of  $M'_1$  with  $w(C_1) < 3r/2$  is at most  $(2(m_1 - s))^3$ . Recall from (3) that  $m_1 \geq 2s + 1$ . For any  $m_1 \geq 2s + 2$ , one can verify that

$$(2(m_1 - s))^3 \leq 240(m_1 - 2s)^5 =: T_0.$$

On the other hand, when  $m_1 = 2s + 1$ , the number of circuits can be at most  $2^{m_1 - s} \leq 2^4$ , which is again bounded by  $T_0$ .

**Case 2:**  $w(C_1) < r/2$

The main point why we distinguish case 2 is that here  $C_1$  is uniquely determined.

**Claim 6.3.** *For any  $e \in S$ , there is at most one circuit  $C_1 \in \mathcal{C}_{1,e}$  with  $w(C_1) < r/2$ .*

*Proof.* For the sake of contradiction, assume that there are two circuits  $C_1, C'_1 \in \mathcal{C}_{1,e}$ , with  $w(C_1), w(C'_1) < r/2$ . By Fact 4.7, we know that  $C_1 \Delta C'_1$  is a disjoint union of circuits in  $M_1$ . Note that  $C_1 \cap S = C'_1 \cap S = \{e\}$ , and hence  $(C_1 \Delta C'_1) \cap S = \emptyset$ . Thus,  $C_1 \Delta C'_1$  is in fact a disjoint union of circuits in  $M$ . Let  $\tilde{C}$  be a subset of  $C_1 \Delta C'_1$  that is a circuit. For the weight of  $\tilde{C}$  we have

$$w(\tilde{C}) \leq w(C_1 \Delta C'_1) \leq w(C_1) + w(C'_1) < r/2 + r/2 = r.$$

This is a contradiction because  $M$  has no circuit of weight less than  $r$ . □

Thus, as we will see, it suffices to bound the number of circuits  $C_2$  in  $M_2$ . Let  $C_e^*$  be the unique choice of a circuit provided by Claim 6.3 (if one exists) for element  $e \in S$ . For the ease of notation, we assume in the following that there is a  $C_e^*$  for every  $e \in S$ . Otherwise we would delete any element  $e \in S$  from  $M_2$  for which no  $C_e^*$  exists, and then would consider the resulting smaller matroid. It might actually be that we thereby delete all of  $S$  from  $M_2$ .

We define a weight function  $w'$  on  $E_2$  as follows:

$$w'(e) := \begin{cases} w(C_e^*), & \text{if } e \in S, \\ w(e), & \text{otherwise.} \end{cases}$$

We now have that any circuit  $C$  of Case 2 can be written as  $C_e^* \Delta C_2$ , for some  $e \in S$ , or  $C = C_2$  when  $C_1 = \emptyset$ . Because  $C_e^*$  is unique, the mapping  $C \mapsto C_2$  is injective for circuits  $C$  of Case 2. Moreover, we have  $w(C) = w'(C_2)$ . This follows from the definition in case that  $C = C_2$ . In the other case, we have

$$w(C) = w(C_e^* \Delta C_2) = w(C_e^*) + w(C_2) = w'(C_2). \tag{7}$$

For the equalities, recall that  $w(e) = 0$  for  $e \in S$ .

We conclude that the number of circuits  $C_2$  in  $M_2$  with  $w'(C_2) < 3r/2$  is an upper bound on the number of Case 2 circuits  $C$  of  $M$  with  $w(C) < 3r/2$ . Now, to get an upper bound on the number of circuits in  $M_2$ , we want to apply induction hypothesis. We need the following claim.

**Claim 6.4.** *There is no circuit  $C_2$  in  $M_2$  with  $w'(C_2) < r$ .*

*Proof.* For the sake of contradiction let  $C_2$  be such a circuit. We show that there exists a circuit  $C'$  in  $M$  with  $w(C') < r$ . This would contradict the assumption of the lemma.

Case(i):  $C_2 \cap S = \emptyset$ . Then  $C_2 \in \mathcal{C}'_2$  itself yields the contradiction because it is a circuit of  $M$  and  $w(C_2) = w'(C_2) < r$ .

Case(ii):  $C_2 \cap S = \{e\}$ . By Fact 4.14, the set  $C_2 \Delta C_e^*$  is a disjoint union of circuits of  $M$ . Let  $C' \subseteq C_2 \Delta C_e^*$  be a circuit of  $M$ . Then, because  $w(e) = 0$ , we have

$$w(C') \leq w(C_e^* \Delta C_2) = w(C_e^*) + w(C_2) = w'(C_2) < r.$$

Case(iii):  $C_2 \cap S = \{e_1, e_2\}$ . By Fact 4.14, similar as in case (ii), there is a set  $C' \subseteq C_2 \Delta C_{e_1}^* \Delta C_{e_2}^*$  that is a circuit of  $M$ . Then, because  $w(e_1) = w(e_2) = 0$ , we have

$$w(C') \leq w(C_2 \Delta C_{e_1}^* \Delta C_{e_2}^*) \leq w(C_2) + w(C_{e_1}^*) + w(C_{e_2}^*) = w'(C_2) < r.$$

Case(iv):  $C_2 \cap S = \{e_1, e_2, e_3\}$ . Since  $S$  is a circuit, it must be the case that  $C_2 = S$ . Since  $C_{e_1}^*, C_{e_2}^*, C_{e_3}^*$  and  $S$  constitute all the circuits of  $M_1$ , the set  $C_{e_1}^* \Delta C_{e_2}^* \Delta C_{e_3}^* \Delta S$  contains a circuit  $C'$  of  $M_1$ . Since  $\{e_i\} = C_{e_i}^* \cap S$ , for  $i = 1, 2, 3$ , we know that  $S \cap C' = \emptyset$ . Thus,  $C' \in \mathcal{C}'_1$  is a circuit of  $M$ . Since  $w(e_1) = w(e_2) = w(e_3) = 0$ , we obtain that

$$w(C') \leq w(C_{e_1}^*) + w(C_{e_2}^*) + w(C_{e_3}^*) = w'(S) = w'(C_2) < r.$$

This proves the claim. □

By Claim 6.4, we can apply the induction hypothesis for  $M_2$  with the weight function  $w'$ . We get that the number of circuits  $C_2$  in  $M_2$  with  $w'(C_2) < 3r/2$  is bounded by

$$T_1 := 240 m_2^5.$$

As mentioned above, this is an upper bound on the number of circuits  $C$  in  $M$  with  $w(C) < 3r/2$  in Case 2.

**Case 3:**  $w(C_1) \geq r/2$

Since  $w(C) = w(C_1) + w(C_2) < 3r/2$ , we have  $w(C_2) < r$  in this case. We also assume that  $C_2 \neq \emptyset$ . Hence, there is an  $e \in S$  such that  $C_1 \in \mathcal{C}_{1,e}$  and  $C_2 \in \mathcal{C}_{2,e}$ .

Let  $T_2$  be an upper bound on the number of circuits  $C_1 \in \mathcal{C}_{1,e}$  with  $w(C_1) < 3r/2$ , for each  $e \in S$ . Let  $T_3$  be an upper bound on the number of circuits  $C_2 \in \mathcal{C}_{2,e}$  with  $w(C_2) < r$ , for each  $e \in S$ . Because there are  $s$  choices for the element  $e \in S$ , the number of circuits  $C = C_1 \Delta C_2$  with  $w(C) < 3r/2$  in Case 3 will be at most

$$s T_2 T_3. \tag{8}$$

To get an upper bound on the number of circuits in  $\mathcal{C}_{1,e}$  and  $\mathcal{C}_{2,e}$ , consider two matroids  $M_{1,e}$  and  $M_{2,e}$ . These are obtained from  $M_1$  and  $M_2$ , respectively, by deleting the elements in  $S \setminus \{e\}$ . The ground set cardinalities of these two matroids are  $m_1 - s + 1$  and  $m_2 - s + 1$ .

We know that for  $i = 1, 2$ , any circuit  $C_i$  of  $M_{i,e}$  with  $e \notin C_i$  is in  $\mathcal{C}'_i$  and hence, is a circuit of  $M$ . Therefore, there is no circuit  $C_i$  of  $M_{i,e}$  with  $e \notin C_i$  and  $w(C_i) < r$ . Using this fact, we want to bound the number of circuits  $C_i$  of  $M_{i,e}$  with  $e \in C_i$ . We start with  $M_{1,e}$ .

**Claim 6.5.** *An upper bound on the number of circuits  $C_1$  in  $M_{1,e}$  with  $e \in C_1$  and  $w(C_1) < 3r/2$  is*

$$T_2 := \min\{8(m_1 - s)^3, 2^{m_1 - s}\} \quad (9)$$

*Proof.* Recall that the decomposition of  $M$  was such that  $M_1$  is graphic, cographic or the  $R_{10}$  matroid.

Case(i). When  $M_1$  is graphic or cographic, the matroid  $M_{1,e}$  falls into the same class by Fact 4.10. Recall that the ground set of  $M_{1,e}$  has cardinality  $m_1 - s + 1$ . In this case, we apply Lemma 6.1 to  $M_{1,e}$  with  $R = \{e\}$  and  $\alpha = 3$  and get a bound of  $8(m_1 - s)^3$ . The number of circuits containing  $e$  is also trivially bounded by the number of all subsets that contain  $e$ , which is  $2^{m_1 - s}$ . Thus, we get Equation (9).

Case(ii). When  $M_1$  is the  $R_{10}$  matroid, then the cardinality of  $M_{1,e}$ , that is  $m_1 - s + 1$ , is at most 10. In this case again, we use the trivial upper bound of  $2^{m_1 - s}$ . One can verify that when  $m_1 - s + 1 \leq 10$  then  $2^{m_1 - s} \leq 8(m_1 - s)^3$ . Thus, we get Equation (9).  $\square$

Next, we want to bound the number of circuits  $C_2$  in  $M_{2,e}$  with  $e \in C_2$  and  $w(C_2) < r$ . This is done in Lemma 6.7 below, where we get a bound of  $T_3 := 48(m_2 - s)^2$ .

To finish Case 3, we now have

$$\begin{aligned} T_2 &= \min\{8(m_1 - s)^3, 2^{m_1 - s}\}, \\ T_3 &= 48(m_2 - s)^2. \end{aligned}$$

By Equation (8), the number of circuits in Case 3 is bounded by  $sT_2T_3$ .

**Claim 6.6.** *For  $s = 1, 3$  and  $m_1 \geq 2s + 1$ ,*

$$sT_2T_3 \leq 2400(m_1 - 2s)^3(m_2 - s)^2.$$

*Proof.* We consider  $sT_2$ . For  $m_1 - 2s \geq 12$ , we have

$$s \cdot 8(m_1 - s)^3 \leq 50(m_1 - 2s)^3.$$

On the other hand, when  $m_1 - 2s \leq 11$ ,

$$s \cdot 2^{m_1 - s} \leq 50(m_1 - 2s)^3.$$

This proves the claim.  $\square$

### Summing up Cases 1, 2 and 3

Finally we add the bounds on the number of circuits of Case 1, 2 and 3. The total upper bound we get is

$$\begin{aligned} T_0 + T_1 + sT_2T_3 &\leq 240(m_1 - 2s)^5 + 240m_2^5 + 240 \binom{5}{2} (m_1 - 2s)^3 (m_2 - s)^2 \\ &\leq 240(m_2 + m_1 - 2s)^5 \\ &\leq 240m^5 \end{aligned}$$

This completes the proof of Theorem 2.6, except for the bound on  $T_3$  that we show in Lemma 6.7.  $\square$

Now we move on to prove Lemma 6.7, which completes the proof of Theorem 2.6. The lemma is similar to Theorem 2.6, but differs in two aspects: (i) we want to count circuits up to a smaller weight bound, that is,  $r$ , and (ii) we have a weaker assumption that there is no circuit of weight less than  $r$  that does not contain a fixed element  $e$ .

**Lemma 6.7.** *Let  $M = (E, \mathcal{I})$  be a connected, regular matroid with ground set size  $m \geq 2$  and  $w: E \rightarrow \mathbb{N}$  be a weight function on  $E$ . Let  $r$  be a positive integer and let  $\tilde{e} \in E$  be any fixed element of the ground set. Assume that there is no circuit  $C$  in  $M$  such that  $\tilde{e} \notin C$  and  $w(C) < r$ . Then, the number of circuits  $C$  in  $M$  such that  $\tilde{e} \in C$  and  $w(C) < r$  is bounded by  $48(m-1)^2$ .*

*Proof.* We closely follow the proof of Theorem 2.6. We proceed again by an induction on  $m$ , the size of the ground set  $E$ .

For the base case, let  $m \leq 10$ . There are at most  $2^{m-1}$  circuits that contain  $\tilde{e}$ . This number is bounded by  $48(m-1)^2$ , for any  $2 \leq m \leq 10$ .

For the inductive step, let  $M = (E, \mathcal{I})$  be a regular matroid with  $|E| = m > 10$  and assume that the theorem holds for all smaller regular matroids. Since  $m > 10$ , matroid  $M$  cannot be  $R_{10}$ . If  $M$  is graphic or cographic, then the bound of the lemma follows from Lemma 6.1. Thus, we may assume that  $M$  is neither graphic nor cographic.

By Theorem 4.18, matroid  $M$  can be written as a 1-, 2-, or 3-sum of two regular matroids  $M_1 = (E_1, \mathcal{I}_1)$  and  $M_2 = (E_2, \mathcal{I}_2)$ . We use the same notation as Theorem 2.6,

$$\begin{aligned} S &= E_1 \cap E_2, \\ s &= |S|, \\ m_i &= |E_i|, \text{ for } i = 1, 2, \\ \mathcal{C}_i &= \{C \mid C \text{ is a circuit of } M_i\}. \end{aligned}$$

The case that  $M$  is a 1-sum of  $M_1$  and  $M_2$  is again trivial. Hence, we may assume that  $M$  is connected. By Theorem 4.19,  $M$  is a 2-sum or a 3-sum of  $M_1$  and  $M_2$ , where  $M_1$  is a graphic, cographic or the  $R_{10}$  matroid, and  $M_2$  is a regular matroid containing  $\tilde{e}$ . For  $i = 1, 2$  and  $e \in S$ , define

$$\mathcal{C}_{i,e} := \{C \mid C \text{ is a circuit of } M_i \text{ and } C_i \cap S = \{e\}\}.$$

Also the weight function  $w$  is extended on  $S$  by  $w(e) = 0$ , for any  $e \in S$ .

We again view each circuit  $C$  of  $M$  as  $C_1 \Delta C_2$  and consider cases based on how the weight of  $C$  is distributed among  $C_1$  and  $C_2$ . Note that  $\tilde{e}$  is in  $M_2$  and we are only interested in circuits  $C$  that contain  $\tilde{e}$ . Hence, we have  $\tilde{e} \in C_2$ . Therefore we do not have the case where  $C_2 = \emptyset$ . We consider the following two cases.

**Case (i).**  $w(C_1) < r/2$ .

**Case (ii).**  $w(C_1) \geq r/2$ .

We will give an upper bound for the number of circuits in each of the two cases.

**Case (i):**  $w(C_1) < r/2$

Since  $\tilde{e} \notin C_1$ , we can literally follow the proof for Case 2 from Theorem 2.6 for this case. We have again Claim 6.3, that  $C_1$  is uniquely determined as  $C_1 = C_e^*$ , for  $e \in S$ , or  $C_1 = \emptyset$ . Therefore the mapping  $C \mapsto C_2$  is injective. The only point to notice now is that the mapping maintains

that  $\tilde{e} \in C$  if and only if  $\tilde{e} \in C_2$ . With the same definition of  $w'$ , we also have  $w(C) = w'(C_2)$ . Therefore it suffices to get an upper bound on the number of circuits  $C_2$  in  $M_2$  with  $w'(C_2) < r$  and  $\tilde{e} \in C_2$ .

To apply the induction hypothesis, we need the following variant of Claim 6.4. It has a similar proof.

**Claim 6.8.** *There is no circuit  $C_2$  in  $M_2$  such that  $w'(C_2) < r$  and  $\tilde{e} \notin C_2$ .*

By the induction hypothesis applied to  $M_2$ , the number of circuits  $C_2$  in  $M_2$  with  $w'(C_2) < r$  and  $\tilde{e} \in C_2$  is bounded by

$$T_0 := 48(m_2 - 1)^2.$$

**Case (ii):**  $w(C_1) \geq r/2$

Since  $w(C) = w(C_1) + w(C_2) < r$ , we have  $w(C_2) < r/2$  in this case. This is the major difference to Case 3 from Theorem 2.6 where the weight of  $C_2$  was only bounded by  $r$ . Hence, now we have again a uniqueness property similar as in Claim 6.3, but for  $C_2$  this time. A difference comes with  $\tilde{e}$ . But the proof remains the same.

**Claim 6.9.** *For any  $e \in S$ , there is at most one circuit  $C_2 \in \mathcal{C}_{2,e}$  with  $w(C_2) < r/2$  and  $\tilde{e} \in C_2$ .*

We conclude that any circuit  $C$  in case (ii) can be written as  $C = C_1 \Delta C_e^*$ , for a  $e \in S$  and the unique circuit  $C_e^* \in \mathcal{C}_{2,e}$ . Therefore the mapping  $C \mapsto C_1$  is injective for the circuits  $C$  of case (ii). Thus, it suffices to count circuits  $C_1 \in \mathcal{C}_{1,e}$  with  $w(C_1) < r$ , for every  $e \in S$ .

Let  $e \in S$  and consider the matroid  $M_{1,e}$  obtained from  $M_1$  by deleting the elements in  $S \setminus \{e\}$ . It has  $m_1 - s + 1$  elements. Since  $M_1$  is a graphic, cographic or  $R_{10}$ , the matroid  $M_{1,e}$  is graphic or cographic by Facts 4.10 and 4.11. The circuits in  $\mathcal{C}_{1,e}$  are also circuits of  $M_{1,e}$ .

Any circuit  $C_1$  of  $M_{1,e}$  with  $e \notin C_1$  is also a circuit of  $M$ . Thus, there is no circuit  $C_1$  of  $M_{1,e}$  with  $e \notin C_1$  and  $w(C_1) < r$ . Therefore we can apply Lemma 6.1 to  $M_{1,e}$  with  $R = \{e\}$ . We conclude that the number of circuits  $C_1 \in \mathcal{C}_{1,e}$  with  $w(C_1) < r$  is at most

$$T_1 := 4(m_1 - s)^2.$$

Since there are  $s$  choices for  $e \in S$ , we obtain a bound of  $sT_1$ .

There is also a trivial bound of  $s2^{m_1-s}$  on the number of such circuits. We take the minimum of the two bounds. Recall from the definition of 2-sum and 3-sum that  $m_1 \geq 2s + 1$ .

**Claim 6.10.** *For  $s = 1$  or  $3$  and  $m_1 \geq 2s + 1$ ,*

$$\min\{s2^{m_1-s}, 4s(m_1 - s)^2\} \leq 48(m_1 - 2s)^2.$$

*Proof.* One can verify that when  $m_1 - 2s \leq 4$  then

$$s2^{m_1-s} \leq 48(m_1 - 2s)^2.$$

On the other hand, when  $m_1 - 2s \geq 5$  then

$$4s(m_1 - s)^2 \leq 48(m_1 - 2s)^2.$$

This proves the claim. □

Hence, we get a bound of  $48(m_1 - 2s)^2$  on the number circuits in case (ii). Now we add the number of circuits of case (i) and (ii) and get a total upper bound of

$$\begin{aligned} 48(m_2 - 1)^2 + 48(m_1 - 2s)^2 &\leq 48(m_2 - 1 + m_1 - 2s)^2 \\ &\leq 48(m - 1)^2. \end{aligned}$$

This gives us the desired bound and completes the proof of Lemma 6.7.  $\square$

## References

- [BH14] Andreas Björklund and Thore Husfeldt. Shortest two disjoint paths in polynomial time. In *Automata, Languages, and Programming - 41st International Colloquium, ICALP 2014, Copenhagen, Denmark, July 8-11, 2014, Proceedings, Part I*, pages 211–222, 2014.
- [Bix77] Robert E Bixby. Kuratowski’s and Wagner’s theorems for matroids. *Journal of Combinatorial Theory, Series B*, 22(1):31 – 53, 1977.
- [FGT16] Stephen A. Fenner, Rohit Gurjar, and Thomas Thierauf. Bipartite perfect matching is in quasi-NC. In *Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2016, Cambridge, MA, USA, June 18-21, 2016*, pages 754–763, 2016.
- [FKS84] Michael L. Fredman, János Komlós, and Endre Szemerédi. Storing a sparse table with  $O(1)$  worst case access time. *J. ACM*, 31(3):538–544, June 1984.
- [GT17] Rohit Gurjar and Thomas Thierauf. Linear matroid intersection is in quasi-NC. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2017*, pages 821–830, New York, NY, USA, 2017. ACM.
- [Kar93] David R. Karger. Global min-cuts in RNC, and other ramifications of a simple min-cut algorithm. In *Proceedings of the Fourth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA ’93*, pages 21–30, Philadelphia, PA, USA, 1993. Society for Industrial and Applied Mathematics.
- [MVV87] Ketan Mulmuley, Umesh V. Vazirani, and Vijay V. Vazirani. Matching is as easy as matrix inversion. *Combinatorica*, 7:105–113, 1987.
- [Nai82] Mohan Nair. On Chebyshev-type inequalities for primes. *The American Mathematical Monthly*, 89(2):126–129, 1982.
- [NSV94] H. Narayanan, Huzur Saran, and Vijay V. Vazirani. Randomized parallel algorithms for matroid union and intersection, with applications to arborescences and edge-disjoint spanning trees. *SIAM J. Comput.*, 23(2):387–397, 1994.
- [Onn10] S. Onn. *Nonlinear Discrete Optimization: An Algorithmic Theory*. Zurich lectures in advanced mathematics. European Mathematical Society Publishing House, 2010.
- [Oxl06] James G. Oxley. *Matroid Theory (Oxford Graduate Texts in Mathematics)*. Oxford University Press, Inc., New York, NY, USA, 2006.
- [RA00] Klaus Reinhardt and Eric Allender. Making nondeterminism unambiguous. *SIAM Journal on Computing*, 29(4):1118–1131, 2000.

- [Sch03a] Alexander Schrijver. *Combinatorial optimization: polyhedra and efficiency. Vol. A. , Paths, flows, matchings.* Algorithms and combinatorics. Springer-Verlag, Berlin, Heidelberg, New York, 2003.
- [Sch03b] Alexander Schrijver. *Combinatorial optimization: polyhedra and efficiency. Vol. B. , Matroids, trees, stable sets.* Algorithms and combinatorics. Springer-Verlag, Berlin, Heidelberg, New York, N.Y., et al., 2003.
- [Sey80] Paul D. Seymour. Decomposition of regular matroids. *J. Comb. Theory, Ser. B*, 28(3):305–359, 1980.
- [ST17] Ola Svensson and Jakub Tarnawski. The matching problem in general graphs is in quasi-NC. *CoRR*, abs/1704.01929, 2017.
- [Sub95] Ashok Subramanian. A polynomial bound on the number of light cycles in an undirected graph. *Information Processing Letters*, 53(4):173 – 176, 1995.
- [TK92] C. P. Teo and K. M. Koh. The number of shortest cycles and the chromatic uniqueness of a graph. *Journal of Graph Theory*, 16(1):7–15, 1992.
- [Tru98] Klaus Truemper. *Matroid Decomposition.* Leibniz, Plano, Texas (USA), 1998.
- [Wig94] Avi Wigderson.  $NL/poly \subseteq \oplus L/poly$ . In *Proceedings of the Ninth Annual Structure in Complexity Theory Conference, Amsterdam, The Netherlands, June 28 - July 1, 1994*, pages 59–62, 1994.

## A Proof of Theorem 4.19

We show some properties of the sum operation on matroids. First note that the  $k$ -sum operations are commutative because their definition is based on symmetric set differences which is commutative. Further, it is known that the  $k$ -sum operations are also associative in some cases. We give a proof here for completeness. The 2-sum operation is denoted by  $\oplus_2$ .

**Lemma A.1** (Associativity). *Let  $M = M_1 \oplus_2 M_2$  with  $e$  being the common element in  $M_1$  and  $M_2$ . Let  $M_2 = M_3 \triangle M_4$  be a  $k$ -sum for  $k = 2$  or  $3$  with the common set  $S$ . Further, let  $e \in M_3$ . Then*

$$M = M_1 \oplus_2 (M_3 \triangle M_4) = (M_1 \oplus_2 M_3) \triangle M_4, \quad (10)$$

where  $M_1 \oplus_2 M_3$  is defined via the common element  $e$  and  $(M_1 \oplus_2 M_3) \triangle M_4$  is defined via the common set  $S$ .

*Proof.* We show that the matroids in Equation (10) have the same circuits. This implies the equality. Let  $E_i$  denote the ground set of  $M_i$ , for  $i = 1, 2, 3, 4$ .

Let  $C$  be a circuit of  $M = M_1 \oplus_2 M_2$ . We consider the nontrivial case in Lemma 4.17: we have  $C = C_1 \triangle C_2$  and  $e \in C_1 \cap C_2$ , where  $C_1$  and  $C_2$  are circuits in  $M_1$  and  $M_2 = M_3 \triangle M_4$ , respectively. Similarly, we have  $C_2 = C_3 \triangle C_4$ , for circuits  $C_3$  and  $C_4$  of  $M_3$  and  $M_4$ , respectively. By our assumption, we have  $e \in C_3$ . It follows that  $C_1 \triangle C_3 \subseteq E_1 \triangle E_3$  is a circuit of  $M_1 \oplus_2 M_3$ . Since  $C_4$  is a circuit of  $M_4$ , we get from Fact 4.14 that  $(C_1 \triangle C_3) \triangle C_4$  is a disjoint union of circuits in  $(M_1 \oplus_2 M_3) \triangle M_4$ .

For the reverse direction, consider a circuit  $C$  of  $(M_1 \oplus_2 M_3) \triangle M_4$ . Similarly as above by Lemma 4.17, we can write  $C = C' \triangle C_4$ , where  $C'$  and  $C_4$  are circuits of  $M_1 \oplus_2 M_3$  and  $M_4$ ,

respectively, with  $S \cap C' = S \cap C_4$ . Further,  $C' = C_1 \triangle C_3$ , where  $C_1$  and  $C_3$  are circuits in  $M_1$  and  $M_3$ , respectively. Since  $S$  is disjoint from  $E_1$ , it must be that  $S \cap C' = S \cap C_3$ . Thus,  $C_3 \triangle C_4 \subseteq E_3 \triangle E_4$  is a union of disjoint circuits in  $M_3 \triangle M_4$ . Since,  $C_1$  is a circuit in  $M_1$ , it follows that  $C_1 \triangle (C_3 \triangle C_4)$  is a disjoint union of circuits in  $M_1 \oplus_2 (M_3 \triangle M_4)$ .

Thus, we have shown that a circuit of one matroid in Equation (10) is a disjoint union of circuits in the other matroid and vice-versa. Consequently, by the minimality of circuits, it follows that their sets of circuits must be the same.  $\square$

Truemper proves the statement of Theorem 4.19 for 3-connected matroids.

**Definition A.2** (3-connected matroid [Tru98]). *A matroid  $M = (E, \mathcal{I})$  is said to be 3-connected if for  $\ell = 1, 2$ , and for any partition  $E = E_1 \cup E_2$  with  $|E_1|, |E_2| \geq \ell$  we have*

$$\text{rank}(E_1) + \text{rank}(E_2) \geq \text{rank}(E) + \ell.$$

**Lemma A.3** (Decomposition of a matroid [Tru98]). *If a binary matroid is not 3-connected then it can be written as a 2-sum or 1-sum of two smaller binary matroids.*

**Theorem A.4** (Truemper's decomposition for 3-connected matroids, [Tru98]). *Let  $M$  be a 3-connected, regular matroid, that is not graphic or cographic and is not isomorphic to  $R_{10}$ . Let  $\tilde{e}$  be a fixed element of the ground set of  $M$ . Then  $M$  is a 3-sum of  $M_1$  and  $M_2$ , where  $M_1$  is a graphic or a cographic matroid and  $M_2$  is a regular matroid that contains  $\tilde{e}$ .*

Theorem 4.19 can be seen as the extension of Theorem A.4 to connected regular matroids.

*Proof of Theorem 4.19.* The proof is by induction on the ground set size of  $M$ . If  $M$  is 3-connected then the statement is true by Theorem A.4. If  $M$  is not 3-connected, then we invoke Lemma A.3. Since  $M$  is connected, it can be written as 2-sum of two matroids  $M = M_1 \oplus_2 M_2$ . From the definition of a 2-sum, it follows that  $M_1$  and  $M_2$  are minors of  $M$  (see [Sey80, Lemma 2.6]), and thus are regular matroids by Fact 4.10. Without loss of generality, let the fixed element  $\tilde{e}$  be in  $M_2$ . If  $M_1$  is graphic, cographic or  $R_{10}$  then we are done.

Suppose,  $M_1$  is neither of these. Let  $e'$  be the element common in the ground sets of  $M_1$  and  $M_2$ . By induction,  $M_1$  is a 2-sum or a 3-sum  $M_1 = M_{11} \triangle M_{12}$ , where  $M_{12}$  is a regular matroid that contains  $e'$  and  $M_{11}$  is a graphic or cographic matroid, or a matroid isomorphic to  $R_{10}$ . Since  $M_{12}$  and  $M_2$  share  $e'$ , we can take the 2-sum of these two matroids using  $e'$ . By Lemma A.1, the matroid  $M$  is the same as  $M_{11} \triangle (M_{12} \oplus_2 M_2)$ . The matroid  $M_{12} \oplus_2 M_2$  contains  $\tilde{e}$  and is regular because both  $M_{12}$  and  $M_2$  are regular (see [Tru98, Theorem 11.3.14]). Thus, the two matroids  $M_{11}$  and  $M_{12} \oplus_2 M_2$  satisfy the desired properties.  $\square$