

Unique Minimizers for Permanents, Mixed Discriminants, and Log-concave Polynomials

Leonid Gurvits and Jonathan Leake

September 17, 2026

Abstract

The permanent and mixed discriminant of positive matrices are classic problems for which we do not expect an efficient algorithm for exact computation. Thus much work has been done to understand how well we can bound and approximately compute these quantities. One line of research in this area begins with the results of [Gur08] where van der Waerden lower bounds of $\frac{n!}{n^n}$ are proven for doubly stochastic inputs for both problems, using a simple proof via stable polynomials. Along with the bound itself, the same techniques are used to show that the permanent and mixed discriminant are uniquely minimized at a certain natural symmetric input.

In this paper, we generalize those results in two ways. First, we extend the unique minimization results beyond doubly stochastic inputs to other marginals which are near doubly stochastic. This yields the first such unique minimization results for the mixed discriminant beyond the doubly stochastic case. We also discuss why one cannot hope similar results to hold in general for all marginals. Second, we extend the unique minimization result for real stable polynomials to strongly log-concave (aka Lorentzian) polynomials in the doubly stochastic case. This captures an analogous previous result on unique minimization for the mixed volume. Finally, we discuss various open problems related to these results.

1 Introduction

The permanent has received significant attention due to its connections to fundamental problems in combinatorics and computer science. This has led to various questions and answers regarding bounding and approximating the permanent under a wide range of different assumptions. One of the most famous is that of the Van der Waerden conjecture from the 1930s (settled in 1980ish by [Fal81; Ego81]), which says

$$\text{per}(M) \geq \frac{n!}{n^n}$$

for all doubly stochastic $M \in \mathbb{R}_{\geq 0}^{n \times n}$ (i.e., $M\mathbf{1} = M^\top \mathbf{1} = \mathbf{1}$), and that $M = \frac{1}{n}\mathbf{1} \cdot \mathbf{1}^\top$ is the unique minimizer over this set of matrices.

This fact was, for example, used to give the first deterministic e^n -approximation algorithm for the permanent of a matrix with non-negative entries [LSW98]. Note that computing the permanent is #P-complete and thus efficient exact computation is not expected to be possible. This is true even when $M \in \{0, 1\}^{n \times n}$, where $\text{per}(M)$ counts perfect matchings of the associated bipartite graph.

Around 20 years ago, the first author initiated a line of work which gave simple proofs of these facts using real stable polynomials and polynomial capacity [Gur06a; Gur08]. This proof method has the added benefit of also applying to the mixed discriminant of positive semidefinite matrices [Gur06b], and to some extent to the mixed volume of convex bodies [Gur09a]. Later works extended these techniques to obtain approximation algorithms for various other combinatorial and probabilistic quantities as well, including the traveling salesman problem [KKO21; GKL24]. These works generalized the doubly stochastic lower bound and rank-one unique minimization result for the permanent, to matrices and real stable polynomials more generally which are only “close” to being doubly stochastic. This extra robustness in the bounds proved to be a crucial step towards these new applications.

The goal of this paper is then to extend the robust rank-one minimizer uniqueness results to new regimes. Specifically, we (1) generalize recent unique minimization results for the permanent to the mixed discriminant, (2) generalize coefficient minimization uniqueness for real stable polynomials to strongly log-concave polynomials, and (3) provide some counterexamples to possible extensions of these and related results.

In the rest of this section, we state and discuss our main results. See Section 2 for any not-yet-defined notions.

1.1 Main Result: Mixed Discriminant Unique Rank-One Minimizer

We now extend the fact that $\frac{1}{n}\mathbf{1}\mathbf{1}^\top$ uniquely minimizes the permanent over all doubly stochastic matrices in a variety of ways. We first recall the definition of the **permanent**:

$$\text{per}(M) = \sum_{\sigma \in S_n} \prod_{i=1}^n m_{i, \sigma(i)}$$

for a given $n \times n$ matrix M . The following is an extension of the unique minimization result for the permanent proven in [GKL24].

Theorem 1.1 ([GKL24]). *For all n there exists $\epsilon > 0$ such that: for all $\mathbf{c}$ such that $\|\mathbf{c} - \mathbf{1}\|_1 < \epsilon$ and $\sum_{i=1}^n c_i = n$, the permanent is uniquely minimized by $M = \frac{1}{n}\mathbf{1} \cdot \mathbf{c}^\top$ over the set of non-negative entry $n \times n$ matrices with row sums $\mathbf{1}$ and column sums $\mathbf{c}$.*

Our first main result generalizes this to the **mixed discriminant**, and hence also reproves [Theorem 1.1](#). Recall its definition:

$$\text{MD}(X_1, \dots, X_n) = \frac{1}{n!} \partial_{t_1} \cdots \partial_{t_n} \det(t_1 X_1 + \cdots + t_n X_n)$$

for a given tuple of $n \times n$ matrices $(X_1, \dots, X_n)$.

Theorem 1.2. *For all n there exists $\epsilon > 0$ such that: for all C of trace n and eigenvalues $\mathbf{c}$ such that $\|\mathbf{c} - \mathbf{1}\|_1 < \epsilon$, the mixed discriminant is uniquely minimized by $(X_1, \dots, X_n) = \frac{1}{n}(C, \dots, C)$ over n -tuples $(X_1, \dots, X_n)$ of trace-1 PSD matrices such that $\sum_{i=1}^n X_i = C$.*

The proof of this result can be found in [Section 4](#), where we also give more explicit bounds on the value of ϵ required for the result.

We finally note that unique minimization of the mixed discriminant in the doubly stochastic case (i.e. when $\|\mathbf{c} - \mathbf{1}\|_1 = 0$) was proven in [\[Gur06b\]](#) using the known equality conditions of the Alexandrov-Fenchel inequalities for the mixed discriminant. We are not aware of such easily understood and verified conditions in the strongly log-concave case, even for mixed volumes; the proofs in this paper do not use the Alexandrov-Fenchel inequalities whatsoever. In a certain sense, the proof path using Alexandrov-Fenchel is hidden in the inductive proofs of analogues of the Van der Waerden and Schrijver's inequalities for stable and strongly-logconcave polynomials, where the equality conditions are replaced by much more elementary convex optimization arguments.

1.2 Main Result: Strongly Log-concave Unique Rank-one Minimizer

The permanent result and the mixed discriminant result are both obtained by analyzing real stable polynomials. See [Section 2.1](#) for definitions and discussion of classes of log-concave polynomials. One general result in this direction is given as follows.

Theorem 1.3 ([\[Gur06a\]](#)). *For all n , the all-ones coefficient is uniquely minimized by $p(\mathbf{x}) = \left(\frac{x_1 + \cdots + x_n}{n}\right)^n$ over homogeneous real stable polynomials p such that $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{1}$.*

Our second result is to generalize [Theorem 1.3](#) by extending it beyond real stable to strongly log-concave polynomials [\[Gur09b\]](#) (also called Lorentzian [\[BH20\]](#) and completely log-concave [\[Ana+18\]](#)). Previous results utilized specific properties of real stable polynomials, and here we are able to generalize beyond this setting.

Theorem 1.4. *For all n , the all-ones coefficient of p is uniquely minimized by $p(\mathbf{x}) = \left(\frac{x_1 + \cdots + x_n}{n}\right)^n$ over homogeneous strongly log-concave polynomials p such that $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{1}$.*

We actually prove a more general version of this result based on polynomial capacity; see [Section 5](#) for the statement and proof.

As a last comment, [Theorem 1.4](#) was proven for volume polynomials (which are strongly log-concave) in [\[Gur09a\]](#) using a more complicated proof. The main difference in the proofs comes from the fact that we are able to show an inductive property of doubly stochastic minimizers; i.e. that the partial derivatives must also be doubly stochastic. See Step 3 of [Section 5.4](#) for further details. See also the last paragraph of [Section 1.1](#) for discussion on the connection of these strongly log-concave results to the Alexandrov-Fenchel inequalities.

1.3 Main Result: Failure of Rank-One Minimizers

To understand the delicateness of the above results, we now provide some negative results for statements which naturally generalize the above results. The first result shows why one of the marginal vectors needs to be $\mathbf{1}$ in the above results.

Proposition 1.5. *For all n and all $\mathbf{r}, \mathbf{c} \neq \mathbf{1}$, the rank-one matrix $\frac{1}{n} \mathbf{r} \mathbf{c}^\top$ does not minimize the permanent over the set of non-negative entry $n \times n$ matrices with row sums $\mathbf{r}$ and column sums $\mathbf{c}$.*

And finally, even in the case where the column sums are $\mathbf{1}$, the row sums need to be close to $\mathbf{1}$ for rank-one unique minimization results to hold. We note that this result already appears in the conference paper [\[GKL24\]](#), but we include it and its proof here also for completeness.

Proposition 1.6 ([\[GKL24\]](#)). *For all n , there exists $\mathbf{c}$ such that: every non-negative entry $n \times n$ matrix with row sums $\mathbf{1}$ and column sums $\mathbf{c}$ has positive permanent, but the set of such matrices is minimized on its relative boundary.*

The proofs of these results can be found in [Section 6](#).

1.4 Open Questions

A number of open questions remain. First of all, there is another version of the rank-one unique minimizer question for the mixed discriminant.

Question 1.7. For all n , does there exist $\epsilon > 0$ such that: for all $\mathbf{t}$ such that $\|\mathbf{t} - \mathbf{1}\|_1 < \epsilon$ and $\sum_{i=1}^n t_i = n$, the mixed discriminant is uniquely minimized by $(X_1, \dots, X_n) = \frac{1}{n}(t_1 I, \dots, t_n I)$ over n -tuples $(X_1, \dots, X_n)$ of PSD matrices such that $\text{tr}(X_i) = t_i$ and $\sum_{i=1}^n X_i = I$?

This question is natural because it can be generalized to all real stable polynomials, as given in the following more general question.

Question 1.8. For all n , does there exist $\epsilon > 0$ such that: for all $\mathbf{t}$ such that $\|\mathbf{t} - \mathbf{1}\|_1 < \epsilon$ and $\sum_{i=1}^n t_i = n$, the coefficient of $x_1 x_2 \cdots x_n$ is uniquely minimized by $p(\mathbf{x}) = \frac{1}{n^n} (\sum_{i=1}^n t_i x_i)^n$ over n -homogeneous n -variate real stable polynomials p such that $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{t}$?

Another natural open question here is related to the tightness of capacity bounds on a certain inner product applied to determinantal polynomials.

Question 1.9. Fix n, N such that $n \leq N$, let $\mathbf{u}_1, \dots, \mathbf{u}_N, \mathbf{v}_1, \dots, \mathbf{v}_N \in \mathbb{C}^n$ be such that $\sum_{i=1}^N \|\mathbf{u}_i \mathbf{u}_i^* \mathbf{v}_i\|_2^2 = \sum_{i=1}^N \|\mathbf{v}_i \mathbf{v}_i^* \mathbf{u}_i\|_2^2 = I$, and define

$$p(\mathbf{x}) = \det \left(\sum_{i=1}^N x_i \mathbf{u}_i \mathbf{u}_i^* \right), \quad q(\mathbf{x}) = \det \left(\sum_{i=1}^N x_i \mathbf{v}_i \mathbf{v}_i^* \right).$$

Do we always have that $\sum_{\alpha} p_{\alpha} q_{\alpha} \geq \frac{n!}{n^n}$, where p_{α} is the $\mathbf{x}^{\alpha}$ coefficient of $p(\mathbf{x})$?

An affirmative answer to this question is actually equivalent to Conjecture 6.1 from [Gur06b] on the quantum permanent (also known as the Cayley hyperdeterminant, or 4-dimensional Pascal determinant) of doubly stochastic separable PSD block matrices. The left-hand side of the inequality in Question 1.9 is precisely the quantum permanent of the associated matrix. A weaker (something like e^{-n}) version of the bound in Question 1.9 can be derived from the results of [AO17], and a more general version of the conjecture also appears as Remark 3.10 in [Gur09b].

We further note that the quantum permanent of a general doubly stochastic PSD block matrix is known to not satisfy such a bound; in fact, it can even take the value 0. That said, efficiently algorithmically determining precisely when it is 0 is an open problem [Gar+20; Gur04] whose resolution would yield efficient algorithms for a version of the famous PIT (polynomial identity testing) problem.

Finally, there is quite a bit of literature (e.g. [For80; Min84; Bru85; Hwa85; FS86; Son+97; PSW11; SB24] etc.) on minimizers of the permanent over faces of the Birkhoff polytope. For example, it is open for many classes of faces of the Birkhoff polytope whether or not the permanent is minimized in its relative interior (such faces are called **cohesive** in the literature; see the survey [CW05]). We briefly tried to apply techniques from this paper to various classes of such faces, but we could not immediately obtain anything new. A natural question is then whether or not this is possible in some cases, and how such statements would generalize to matrices with column sums $\mathbf{1}$ and row sums $\approx \mathbf{1}$.

1.5 AI Declaration

We note that the literature review for the last paragraph of Section 1.4, along with some testing to see if our results could apply in the context of cohesive faces of the Birkhoff polytope, was completed with the help of AI (ChatGPT 5.6 Sol). The rest of this paper was created by the authors without the use of any AI tool, except perhaps for creating some .bib file entries.

2 Preliminaries

Throughout we use standard vector and set notation. We let $\mathbb{C}, \mathbb{R}, \mathbb{Z}$ denote the complex, real, and integer numbers, respectively. We use a subscript to modify

these sets; for example $\mathbb{Z}_{\geq 0}$ denote the non-negative integers. We also use other sensible notation; e.g. we let $\mathbb{R}^{n \times n}$ denote the set of $n \times n$ matrices with real entries. Also, given $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ we use $\mathbf{x} \leq \mathbf{y}$ to denote $x_i \leq y_i$ for all $i \in [n]$, we denote $\mathbf{x} \odot \mathbf{y} = (x_1 y_1, \dots, x_n y_n)$, we denote $\mathbf{x}^{\mathbf{y}} = x_1^{y_1} \cdots x_n^{y_n}$ (whenever $\mathbf{x} > \mathbf{0}$), we denote $\|\mathbf{x}\|_1 = \sum_{i=1}^n |x_i|$, and we denote $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^n x_i y_i$.

We will make use of a number of shorthands for various spaces throughout. Specifically, we define certain matrix spaces:

- $\text{Mat}_n(\mathbf{c}) = \left\{ M \in \mathbb{R}_{\geq 0}^{n \times n} : M\mathbf{1} = \mathbf{1}, M^\top \mathbf{1} = \mathbf{c} \right\}$,
- PSD_n = the set of $n \times n$ positive semidefinite matrices,
- $\text{MatTup}_n(C) = \{(M_1, \dots, M_n) \in \text{PSD}_n^n : \text{tr}(M_i) = 1 \ \forall i, \sum_{i=1}^n M_i = C\}$.

2.1 Log-concave Polynomials

Here we give some definitions and useful notation, before recalling a few results we will make use of.

Definition 2.1. An n -homogeneous polynomial $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ is **real stable** if $p(z_1, \dots, z_n) \neq 0$ for all $z_1, \dots, z_n$ in the (open) complex upper half-plane.

Definition 2.2. An n -homogeneous polynomial $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ is **strongly log-concave (SLC)** if $\partial_{x_1}^{\alpha_1} \cdots \partial_{x_n}^{\alpha_n} p(\mathbf{x})$ is either log-concave on $\mathbb{R}_{>0}^n$ or identically zero for all $\alpha \in \mathbb{Z}_{\geq 0}^n$.

Example 2.3. For all $A_1, \dots, A_n \in \text{PSD}_d$, the polynomial $p(\mathbf{x}) = \det(\sum_{i=1}^n x_i A_i)$ is real stable. By considering diagonal matrices, this implies products of linear forms with non-negative coefficients are also real stable. Finally, for all compact convex bodies $K_1, \dots, K_n \subseteq \mathbb{R}^d$, the polynomial $p(\mathbf{x}) = \text{vol}(\sum_{i=1}^n x_i K_i)$ is strongly log-concave, but not necessarily real stable.

We will not explicitly use the following, but it is a well-known fact in the literature.

Proposition 2.4. *Every homogeneous real stable polynomial is strongly log-concave.*

We note that strongly log-concave is equivalent to **Lorentzian** [BH20] and **completely log-concave** [Ana+18] for homogeneous polynomials. That said, we also give one other equivalent definition of strongly log-concave which relies upon the following folklore result in convex analysis.

Lemma 2.5. *If $p \in \mathbb{R}[x_1, \dots, x_n]$ is d -homogeneous and $\mathbf{x} \in \mathbb{R}^n$ is such that $p(\mathbf{x}) > 0$, then the Hessian of $\log p$ at $\mathbf{x}$ is negative semidefinite if and only if the Hessian of $p^{1/d}$ at $\mathbf{x}$ is negative semidefinite.*

This leads to the following alternative definition of strongly log-concave polynomials.

Corollary 2.6. *If $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ is d -homogeneous, then p is log-concave on $\mathbb{R}_{> 0}^n$ if and only if $p^{1/d}$ is concave on $\mathbb{R}_{> 0}^n$.*

We now state a few basic properties of real stable and SLC polynomials from the literature, which have straightforward proofs. We first need a definition, which describes the Newton's inequalities that hold for real-rooted polynomials.

Definition 2.7. A polynomial $f \in \mathbb{R}_{\geq 0}[t]$ of degree at most n is **n -Newton** if for $f(t) = \sum_{k=0}^n \binom{n}{k} f_k t^k$ we have $f_k^2 \geq f_{k-1} f_{k+1}$ for all k .

Lemma 2.8. *Both the classes of real stable and strongly log-concave polynomials are preserved under taking partial derivatives and under external fields (i.e. $p(\mathbf{x}) \mapsto \frac{p(\mathbf{a} \odot \mathbf{x})}{p(\mathbf{a})}$ for $\mathbf{a} \in \mathbb{R}_{> 0}^n$).*

Lemma 2.9. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be d -homogeneous, and fix $x_1, \dots, x_{n-1} \in \mathbb{R}_{\geq 0}$. If p is real stable then $f(t) = p(x_1, \dots, x_{n-1}, t)$ is real-rooted, and if p is strongly log-concave then $f(t) = p(x_1, \dots, x_{n-1}, t)$ is d -Newton.*

Finally, we generalize some of the notation use for matrices above to polynomials. Specifically we define the following spaces of polynomials:

- $\text{HStab}_n(\mathbf{c})$ is the set of all n -variate n -homogeneous real stable polynomials p such that $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{c}$, and
- $\text{SLC}_n(\mathbf{c})$ is the set of all n -variate n -homogeneous strongly log-concave polynomials p such that $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{c}$.

We will also call a homogeneous polynomial p **doubly stochastic** if $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{1}$. Note that this implies p is n -variate and n -homogeneous.

2.2 Polynomial Capacity Bounds

In this section, we recall various results from the literature on polynomial capacity. Given $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ and $\alpha \in \mathbb{R}_{\geq 0}^n$, the **α -capacity** of p is defined as

$$\text{Cap}_{\alpha}(p) := \inf_{\mathbf{x} \in \mathbb{R}_{> 0}^n} \frac{p(\mathbf{x})}{\mathbf{x}^{\alpha}}.$$

We often just say **capacity** when α is understood from context. Polynomial capacity has been used in various contexts to lower-bound combinatorial and probabilistic quantities which are hard to compute. We now recall some of these bounds, which quantifies the difference between the capacity of a polynomial and its coefficients.

Theorem 2.10 ([Gur09b]). *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be a d -homogeneous strongly log-concave polynomial. Fix $\alpha \in \mathbb{R}_{\geq 0}^n$ such that $\sum_i \alpha_i = d$. Then*

$$\frac{p_{\alpha}}{\text{Cap}_{\alpha}(p)} \geq \binom{d}{\alpha} \frac{\alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}}{d^d},$$

where p_{α} is the coefficient of $\mathbf{x}^{\alpha}$ in p .

Theorem 2.11 ([Gur15, Thm. 5.1]). Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be a d -homogeneous real stable polynomial. Fix $\alpha \in \mathbb{R}_{\geq 0}^n$ such that $\sum_i \alpha_i = d$, and let d_i be the degree of x_i in

$$\partial_{x_{i+1}}^{\alpha_{i+1}} \cdots \partial_{x_n}^{\alpha_n} p \Big|_{x_{i+1}=\cdots=x_n=0}$$

for $i = 1, \dots, n$. Then we have

$$\frac{p_\alpha}{\text{Cap}_\alpha(p)} \geq \prod_{i=2}^n \binom{d_i}{\alpha_i} \frac{\alpha_i^{\alpha_i} (d_i - \alpha_i)^{d_i - \alpha_i}}{d_i^{d_i}},$$

where p_α is the coefficient of $\mathbf{x}^\alpha$ in p .

It is straightforward to see that the lower bound of Theorem 2.11 is strictly decreasing in d_i for fixed α_i . With this, we obtain the following immediate corollary of Theorem 2.11.

Corollary 2.12. Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be an n -homogeneous real stable polynomial such that $p(\mathbf{e}_i) = 0$ for some $i \in [n]$. Then

$$\frac{\partial_{x_1} \cdots \partial_{x_n} p(\mathbf{x})}{\text{Cap}_1(p)} \geq \frac{(n-2)^{n-2} (n-1)!}{(n-1)^{2n-3}} > \frac{n!}{n^n}.$$

The next result now gives a lower bound on the capacity itself for real stable polynomials under certain conditions on the gradient of the polynomial. This and similar results were originally used to prove Theorem 1.1 and to improve the approximation factor for the traveling salesman problem.

Theorem 2.13 ([GL21]). If $p \in \mathbb{R}[x_1, \dots, x_n]$ is a n -homogeneous real stable polynomial such that $p(\mathbf{1}) = 1$ and $\|\mathbf{1} - \nabla p(\mathbf{1})\|_1 < 2$, then

$$\text{Cap}_1(p) \geq \left(1 - \frac{\|\mathbf{1} - \nabla p(\mathbf{1})\|_1}{2}\right)^n.$$

Finally, we give one final result on the capacity of product of linear forms. This result is known in the literature, but we give a short proof here. We note that this sort of result can also be proven for capacity on completely positive operators.

Lemma 2.14. For any $n \times n$ matrix M with non-negative entries, we have

$$\text{Cap}_1 \left(\prod_{i=1}^n (M\mathbf{x})_i \right) = \frac{1}{n^n} \left(\inf_{\mathbf{x}, \mathbf{y} > \mathbf{0}} \frac{\langle M\mathbf{x}, \mathbf{y} \rangle}{\mathbf{x}^1 \mathbf{y}^1} \right)^n.$$

In particular,

$$\text{Cap}_1 \left(\prod_{i=1}^n (M\mathbf{x})_i \right) = \text{Cap}_1 \left(\prod_{i=1}^n (M^\top \mathbf{x})_i \right).$$

Proof. Let $p_M(\mathbf{x}) = \prod_{i=1}^n (M\mathbf{x})_i$. For all $\mathbf{x}, \mathbf{y} > \mathbf{0}$, by AM-GM we have

$$\left(\frac{\langle \mathbf{x}, \mathbf{y} \rangle}{n} \right)^n \geq \prod_{i=1}^n x_i y_i = \mathbf{x}^1 \mathbf{y}^1,$$

with equality if and only if $y_i = x_i^{-1}$ for all $i \in [n]$. Thus we first have

$$\frac{1}{n^n} \left(\inf_{\mathbf{x}, \mathbf{y} > \mathbf{0}} \frac{\langle M\mathbf{x}, \mathbf{y} \rangle}{\mathbf{x}^1 \mathbf{y}^1} \right)^n \geq \inf_{\mathbf{x} > \mathbf{0}} \frac{\prod_{i=1}^n (M\mathbf{x})_i}{\mathbf{x}^1} = \text{Cap}_1(p_M).$$

By the above equality condition, we also have

$$\frac{1}{n^n} \left(\inf_{\mathbf{x}, \mathbf{y} > \mathbf{0}} \frac{\langle M\mathbf{x}, \mathbf{y} \rangle}{\mathbf{x}^1 \mathbf{y}^1} \right)^n \leq \frac{1}{n^n} \left(\inf_{\mathbf{x} > \mathbf{0}} \frac{\langle M\mathbf{x}, (M\mathbf{x})^{-1} \rangle}{\mathbf{x}^1 (M\mathbf{x})^{-1}} \right)^n = \text{Cap}_1(p_M)$$

which completes the proof. $\square$

3 Optimizing Forms

In this section, we give some general results on optimizing forms. These results are mainly about determining what sort of optima a form can have on a given feasible region. The main result of this section is [Theorem 3.5](#), which proves that minimizers of symmetric multilinear forms over symmetric compact convex sets are symmetric, so long as no minimizers occur on the boundary. We will later use this result to prove our main result on the uniqueness of the rank-one minimizer for the mixed discriminant. That said, we have decided to state the results here more generally, as they seem interesting in their own right.

The first result is on quadratic forms which we will bootstrap to prove the main result.

Lemma 3.1. *Let $q(\mathbf{x})$ be a quadratic form on $\mathbb{R}^n$, and let $S \subset \mathbb{R}^n$ be a closed convex set containing no affine lines. If $\min\{q(\mathbf{x}) : \mathbf{x} \in S\}$ has no minimizers on the relative boundary of S , then it has at most one minimizer in S .*

Proof. Suppose $\min\{q(\mathbf{x}) : \mathbf{x} \in S\}$ has no minima on the relative boundary S , but $\mathbf{u} \neq \mathbf{v}$ are two minima in the relative interior of S . Letting $W \subseteq \mathbb{R}^n$ denote the orthogonal complement to the vector space parallel to the affine span of S , minimality in the relative interior implies $\nabla q(\mathbf{u}), \nabla q(\mathbf{v}) \in W$. Since q is a quadratic form, its gradient is a linear form, and thus $\nabla q(\mathbf{y}) \in W$ for all $\mathbf{y}$ in the line L through $\mathbf{u}$ and $\mathbf{v}$. Therefore every $\mathbf{y} \in S \cap L$ minimizes $q(\mathbf{x})$ over S . Since $S \cap L$ contains a point of the relative boundary of S , this is a contradiction. $\square$

We now prove a few basic convex geometric results.

Lemma 3.2. *Fix a set $S \subseteq (\mathbb{R}^n)^d$ with $d \geq 2$ and any $\mathbf{w}_3, \dots, \mathbf{w}_d \in \mathbb{R}^n$. If $T = \{(\mathbf{u}, \mathbf{v}) \in (\mathbb{R}^n)^2 : (\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d) \in S\}$ contains an affine line, then S contains an affine line.*

Proof. By assumption, $(\mathbf{u}_0, \mathbf{v}_0) + t \cdot (\mathbf{u}_1, \mathbf{v}_1) \in T$ for all $t \in \mathbb{R}$. Therefore $(\mathbf{u}_0, \mathbf{v}_0, \mathbf{w}_3, \dots, \mathbf{w}_d) + t \cdot (\mathbf{u}_1, \mathbf{v}_1, \mathbf{0}, \dots, \mathbf{0}) \in S$ for all $t \in \mathbb{R}$ as desired. $\square$

Lemma 3.3. *Fix a closed convex set $S \subseteq (\mathbb{R}^n)^d$ with $d \geq 2$ and any $\mathbf{w}_3, \dots, \mathbf{w}_d \in \mathbb{R}^n$. If $(\mathbf{u}, \mathbf{v})$ is on the relative boundary of $T = \{(\mathbf{x}, \mathbf{y}) \in (\mathbb{R}^n)^2 : (\mathbf{x}, \mathbf{y}, \mathbf{w}_3, \dots, \mathbf{w}_d) \in S\}$, then $(\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d)$ is on the relative boundary of S .*

Proof. Fix $(\mathbf{u}, \mathbf{v})$ on the relative boundary of T . Let $B \subset \text{aff}(S)$ be some small relatively open ball centered at $(\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d)$, and define

$$B' := \{(\mathbf{x}, \mathbf{y}) : (\mathbf{x}, \mathbf{y}, \mathbf{w}_3, \dots, \mathbf{w}_d) \in B\}.$$

Thus B' is relatively open in $\text{aff}(T)$ and contains $(\mathbf{u}, \mathbf{v})$. Since $(\mathbf{u}, \mathbf{v})$ is on the relative boundary of T , there exists $(\mathbf{u}', \mathbf{v}') \in B' \setminus T$. Therefore $(\mathbf{u}', \mathbf{v}', \mathbf{w}_3, \dots, \mathbf{w}_d) \in B \setminus S$, which implies $B \not\subset S$. Since this holds for any small ball about $(\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d)$, we must have that $(\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d)$ is on the relative boundary of S . $\square$

Finally, we now prove the main result of the section. First we need a definition.

Definition 3.4. We call a set $S \subseteq (\mathbb{R}^n)^d$ S_d -invariant if for all $(\mathbf{w}_1, \dots, \mathbf{w}_d) \in S$ and all $\sigma \in S_d$ we have that $(\mathbf{w}_{\sigma(1)}, \dots, \mathbf{w}_{\sigma(d)}) \in S$.

Theorem 3.5. *Let $P(\mathbf{x}_1, \dots, \mathbf{x}_d)$ be a symmetric multilinear form on $\mathbb{R}^n$ for some $d \geq 2$, and let $S \subset (\mathbb{R}^n)^d$ be an S_d -invariant closed convex set containing no affine lines. If $\min\{P(\mathbf{x}_1, \dots, \mathbf{x}_d) : (\mathbf{x}_1, \dots, \mathbf{x}_d) \in S\}$ has no minimizers on the relative boundary of S , then all its minimizers over S are of the form $(\mathbf{w}, \dots, \mathbf{w})$ for some $\mathbf{w} \in \mathbb{R}^n$.*

Proof. If P has no minimizer in S , then we are done. Otherwise suppose $(\mathbf{w}_1, \dots, \mathbf{w}_d)$ is a minimizer of P over S , which by assumption is contained in the relative interior of S . Define a function

$$f(\mathbf{u}, \mathbf{v}) := P(\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d),$$

which is a symmetric bilinear form on $\mathbb{R}^n$. Further, one can consider $q(\mathbf{u}, \mathbf{v}) = f(\mathbf{u}, \mathbf{v})$ to be a quadratic form on $\mathbb{R}^{2n}$. Thus $(\mathbf{w}_1, \mathbf{w}_2)$ is a minimizer of q over $T = \{(\mathbf{u}, \mathbf{v}) \in (\mathbb{R}^n)^2 : (\mathbf{u}, \mathbf{v}, \mathbf{w}_3, \dots, \mathbf{w}_d) \in S\}$.

Suppose now that $\min\{q(\mathbf{x}, \mathbf{y}) : (\mathbf{x}, \mathbf{y}) \in T\}$ has a minimizer $(\mathbf{u}', \mathbf{v}')$ on the relative boundary of T . Then $(\mathbf{u}', \mathbf{v}', \mathbf{w}_3, \dots, \mathbf{w}_d)$ is a minimizer of P over S , and also $(\mathbf{u}', \mathbf{v}', \mathbf{w}_3, \dots, \mathbf{w}_d)$ is contained in the relative boundary of S by [Lemma 3.3](#). This contradicts the assumptions, and thus $\min\{q(\mathbf{x}, \mathbf{y}) : (\mathbf{x}, \mathbf{y}) \in T\}$ has no minimizers on the relative boundary of T . Thus by [Lemma 3.1](#) and [Lemma 3.2](#), $\min\{q(\mathbf{x}, \mathbf{y}) : (\mathbf{x}, \mathbf{y}) \in T\}$ has at most one minimizer in T , and therefore $(\mathbf{w}_1, \mathbf{w}_2)$ is the unique minimizer of q over T .

But by symmetry of P and S_d -invariance of S , $(\mathbf{w}_2, \mathbf{w}_1)$ is also a minimizer of q over T , and thus $\mathbf{w}_1 = \mathbf{w}_2$. By applying the same argument to $(\mathbf{w}_i, \mathbf{w}_j)$ for all $i \neq j$, we find that there is some $\mathbf{w} \in S$ such that $\mathbf{w}_i = \mathbf{w}$ for all i . Thus all minimizers of P are of the form $(\mathbf{w}, \dots, \mathbf{w})$ for some $\mathbf{w} \in \mathbb{R}^n$. $\square$

4 Unique Minimizer: Mixed Discriminant

In this section we prove our first main result [Theorem 1.2](#) on the unique minimization of the mixed discriminant. Recall that the **mixed discriminant** is defined via

$$\text{MD}(M_1, \dots, M_n) = \frac{1}{n!} \partial_{x_1} \cdots \partial_{x_n} \det(x_1 A_1 + \cdots + x_n A_n)$$

for a given tuple of $n \times n$ matrices $(A_1, \dots, A_n)$. Our main result can then be re-stated as follows.

Theorem 4.1. *For all n there exists $\epsilon > 0$ such that: for all $C \in \text{PSD}_n$ with eigenvalues $\mathbf{c}$ such that $\text{tr}(C) = n$ and $\|\mathbf{c} - \mathbf{1}\|_1 < \epsilon$, the mixed discriminant is uniquely minimized over $\text{MatTup}_n(C)$ at $\frac{1}{n}(C, C, \dots, C)$.*

More concretely, the result holds for any $C \in \text{PSD}_n$ with eigenvalues $\mathbf{c}$ such that $\text{tr}(C) = n$ and

$$\frac{1}{c_1 \cdots c_n} \left(1 - \frac{\|\mathbf{c} - \mathbf{1}\|_1}{2}\right)^n > \frac{(n-1)^{2n-3}}{n^{n-1}(n-2)^{n-2}}.$$

We now prove [Theorem 4.1](#). Fix $(A_1, \dots, A_n) \in \text{MatTup}_n(C)$. We consider

$$p(x_1, \dots, x_n) = \det\left(\sum_{i=1}^n x_i A_i\right),$$

where the mixed discriminant of $A_1, \dots, A_n$ is proportional to the all-ones coefficient of p .

Step 1: Lower bound the capacity of p . Fix $x_1, \dots, x_n \in \mathbb{R}$. Since $\sum_{i=1}^n x_i A_i$ is Hermitian, there exist unitary U and diagonal D (dependent on x) such that

$$UDU^* = \sum_{i=1}^n x_i A_i.$$

This implies

$$D = \sum_{i=1}^n x_i U^* A_i U,$$

which gives

$$p(x_1, \dots, x_n) = \prod_{i=1}^n \sum_{j=1}^n M_{ij} x_j$$

where $M_{ij} = (U^* A_j U)_{ii} \geq 0$ since A_j is PSD for all $j \in [n]$. Note that for each j we have

$$\sum_{i=1}^n M_{ij} = \sum_{i=1}^n (U^* A_j U)_{ii} = \text{tr}(U^* A_j U) = 1,$$

and for each i we have

$$\sum_{j=1}^n M_{ij} = \sum_{j=1}^n (U^* A_j U)_{ii} = (U^* C U)_{ii}.$$

Thus the column sums of M are $\mathbf{1}$ and the row sums $\mathbf{a}$ of M are **majorized** by $\mathbf{c}$ (see condition (1) of [Definition 4.2](#) below). We recall some equivalent definitions of majorization as follows. We will not use all of these, but have decided to include them for the interested reader.

Definition 4.2 (e.g. [\[MOA11\]](#)). Given $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, we say that $\mathbf{y}$ **majorizes** $\mathbf{x}$, equivalently written $\mathbf{x} \preceq \mathbf{y}$ if one of the following equivalent conditions holds:

1. there is a Hermitian matrix with diagonal entries $\mathbf{x}$ and eigenvalues $\mathbf{y}$,
2. $\|\mathbf{x} - \alpha \mathbf{1}\|_1 \leq \|\mathbf{y} - \alpha \mathbf{1}\|_1$ for all $\alpha \in \mathbb{R}$,
3. $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$ and $\sum_{i=1}^k x_i^\downarrow \leq \sum_{i=1}^k y_i^\downarrow$ for all $k \in [n]$, where $\mathbf{x}^\downarrow$ is $\mathbf{x}$ with entries sorted in non-increasing order,
4. there exists $D \in \text{Mat}_n(\mathbf{1})$ such that $D\mathbf{y} = \mathbf{x}$, and
5. $\text{conv}(\sigma \cdot \mathbf{x} : \sigma \in S_n) \subseteq \text{conv}(\sigma \cdot \mathbf{y} : \sigma \in S_n)$.

Note that condition (2) of [Definition 4.2](#) implies $\|\mathbf{a} - \mathbf{1}\|_1 \leq \|\mathbf{c} - \mathbf{1}\|_1$. Therefore by combining the above with [Lemma 2.14](#) and [Theorem 2.13](#), for all $\mathbf{x} \in \mathbb{R}_{>0}^n$ we have

$$\begin{aligned} \frac{p(\mathbf{x})}{x_1 \cdots x_n} &= \frac{\prod_{i=1}^n (M\mathbf{x})_i}{x_1 \cdots x_n} \\ &\geq \text{Cap}_1 \left(\prod_{i=1}^n (M\mathbf{x})_i \right) = \text{Cap}_1 \left(\prod_{i=1}^n (M^\top \mathbf{x})_i \right) \\ &\geq \left(1 - \frac{\|\mathbf{a} - \mathbf{1}\|_1}{2} \right)^n \geq \left(1 - \frac{\|\mathbf{c} - \mathbf{1}\|_1}{2} \right)^n \end{aligned}$$

whenever $\|\mathbf{c} - \mathbf{1}\|_1 < 2$. This implies

$$\text{Cap}_1(p) \geq \left(1 - \frac{\|\mathbf{c} - \mathbf{1}\|_1}{2} \right)^n. \quad (1)$$

Step 2: Apply [Theorem 3.5](#) We note that MD is a symmetric multilinear form, and that $\text{MatTup}_n(C)$ is a compact convex S_n -invariant set. Thus to apply [Theorem 3.5](#), we need to study the relative boundary of $\text{MatTup}_n(C)$. A tuple $(A_1, \dots, A_n)$ is on the relative boundary of $\text{MatTup}_n(C)$ if and only if A_i is singular for some $i \in [n]$. In this case, we have

$$p(\mathbf{e}_i) = \det(A_i) = 0,$$

which by [Corollary 2.12](#) implies

$$\frac{\text{MD}(A_1, \dots, A_n)}{\text{Cap}_1(p)} \geq \frac{(n-2)^{n-2}}{n(n-1)^{2n-3}}. \quad (2)$$

On the other hand, we compute

$$\text{MD}\left(\frac{C}{n}, \dots, \frac{C}{n}\right) = \frac{\det(C)}{n^n} = \frac{c_1 \cdots c_n}{n^n}.$$

Now if $\mathbf{c}$ is such that

$$\frac{(n-2)^{n-2}}{n(n-1)^{2n-3}} \left(1 - \frac{\|\mathbf{c} - \mathbf{1}\|_1}{2}\right)^n > \frac{c_1 \cdots c_n}{n^n},$$

we combine (1) and (2) to obtain

$$\text{MD}(A_1, \dots, A_n) > \frac{c_1 \cdots c_n}{n^n} = \text{MD}\left(\frac{C}{n}, \dots, \frac{C}{n}\right)$$

for all $(A_1, \dots, A_n)$ on the relative boundary of $\text{MatTup}_n(C)$. Therefore we can apply [Theorem 3.5](#) to conclude that all minimizers of MD over $\text{MatTup}_n(C)$ are of the form $(A, \dots, A)$. The only such element of $\text{MatTup}_n(C)$ is $\frac{1}{n}(C, \dots, C)$, and thus this uniquely minimizes MD over $\text{MatTup}_n(C)$.

5 Unique Minimizer: SLC Polynomials

The point of this section is to prove [Theorem 1.4](#); that is, to generalize to strongly log-concave polynomials the fact that the permanent is uniquely minimized over doubly stochastic matrices by the multiple of the all-ones matrix. An analogous fact was proven for doubly stochastic real stable polynomials in [\[Gur06a\]](#). Let us explicitly recall [Theorem 1.4](#).

Theorem 5.1. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous and SLC. If*

$$\partial_{x_1} \cdots \partial_{x_n} p(\mathbf{x}) = \frac{n!}{n^n} \text{Cap}_1(p) > 0,$$

then for $\mathbf{v} = \frac{\nabla p(\mathbf{1})}{n}$ we have $p(\mathbf{x}) = (\sum_i v_i x_i)^n$.

5.1 Univariate n -Newton Polynomials

To prove [Theorem 5.1](#), we first need a number of facts about the class of n -Newton univariate polynomials derived from SLC polynomials. Recall the definition of n -Newton in [Definition 2.7](#). The main result of this section is [Theorem 5.4](#), which gives a refined capacity bound for n -Newton polynomials based on their degree. To prove this, we first need a few basic results on the capacity of n -Newton polynomials.

Lemma 5.2. *For any $a > 0$, we have that*

$$\text{Cap}_1 \left(\left(1 + \frac{a}{n} t \right)^n \right) = a \left(\frac{n}{n-1} \right)^{n-1},$$

with the optimum achieved at $t = \frac{n}{a(n-1)}$.

Proof. We compute

$$\partial_t \left[\frac{1}{t} \left(1 + \frac{a}{n} t \right)^n \right] = \frac{1}{t^2} \left(1 + \frac{a}{n} \right)^{n-1} \left(at \cdot \frac{n-1}{n} - 1 \right),$$

and thus the derivative is 0 exactly at $t = \frac{n}{a(n-1)}$. Therefore $\text{Cap}_1 \left(\left(1 + \frac{a}{n} t \right)^n \right)$ is optimized at $t = \frac{n}{a(n-1)}$ and

$$\text{Cap}_1 \left(\left(1 + \frac{a}{n} t \right)^n \right) = \frac{a(n-1)}{n} \left(\frac{n}{n-1} \right)^n = a \left(\frac{n}{n-1} \right)^{n-1}.$$

□

Lemma 5.3. *Let $f, g \in \mathbb{R}_{\geq 0}[t]$ be such that f is of degree at least 2 and $g(0) > 0$. Then $\text{Cap}_1(f)$ is optimized at some $t_0 > 0$, and if $g(t) < f(t)$ for all $t > 0$ then $\text{Cap}_1(g) < \text{Cap}_1(f)$.*

Proof. The fact that $\lim_{t \rightarrow 0^+} \frac{f(t)}{t} = +\infty = \lim_{t \rightarrow +\infty} \frac{f(t)}{t}$ implies the first claim. Now let $t_0 > 0$ be a value of t which optimizes $\text{Cap}_1(f)$. Then we have

$$\text{Cap}_1(f) = \frac{f(t_0)}{t_0} > \frac{g(t_0)}{t_0} \geq \text{Cap}_1(g),$$

which implies the second claim. □

We now prove the main result of the section. To do this, we make use of the **truncation** operator, which “cuts off” terms of a polynomial above a certain degree. Formally, for $d \leq n$ we define

$$\text{trunc}_d \left(\sum_{k=0}^n c_k t^k \right) = \sum_{k=0}^d c_k t^k.$$

Theorem 5.4. *For all d, n such that $1 \leq d \leq n$ there exist constants $\gamma_{n,d} \geq 1$ such that:*

1. *for all n -Newton polynomials $f(t)$ of degree at most d such that $\partial_t f(0) > 0$ we have*

$$\partial_t f(0) \geq \gamma_{n,d} \left(\frac{n-1}{n} \right)^{n-1} \text{Cap}_1(f),$$

2. $1 = \gamma_{n,n} < \gamma_{n,n-1} < \dots < \gamma_{n,2} < \gamma_{n,1} = \left(\frac{n}{n-1} \right)^{n-1}$, and

3. assuming $d \geq 2$ (and $\partial_t f(0) > 0$), the inequality of (1) is an equality if and only if $f(t) = \text{trunc}_d [(a + bt)^n]$ for some $a, b > 0$.

Proof. We define

$$\gamma_{n,d} = \frac{\text{Cap}_1((1+t)^n)}{\text{Cap}_1(\text{trunc}_d[(1+t)^n])}.$$

Note that $\gamma_{n,n} = 1$ and $\gamma_{n,1} = \left(\frac{n}{n-1}\right)^{n-1}$ by Lemma 5.2, and $\gamma_{n,n} < \gamma_{n,n-1} < \dots < \gamma_{n,2} < \gamma_{n,1}$ by Lemma 5.3. This implies (2) and the ($\Leftarrow$) direction of (3).

Now if $d = 1$ or $f(0) = 0$ then $\partial_t f(0) = \text{Cap}_1(f)$. Thus for the remainder of the proof, we may assume without loss of generality that $f(0) = 1$ and $d \geq 2$. Since f is n -Newton we have

$$\begin{aligned} f(t) &= \binom{n}{0} + \binom{n}{1} \frac{\partial_t f(0)}{n} t + \sum_{k=2}^d \binom{n}{k} c_k t^k \\ &\leq \binom{n}{0} + \binom{n}{1} \frac{\partial_t f(0)}{n} t + \sum_{k=2}^d \binom{n}{k} \left(\frac{\partial_t f(0)}{n}\right)^k t^k \\ &= \text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right]. \end{aligned}$$

Note that if $f(t)$ is not equal to $g(t) := \text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right]$ then the above inequality is strict for all $t > 0$. Thus by Lemma 5.3 we have

$$\text{Cap}_1(f) \leq \text{Cap}_1 \left(\text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right] \right),$$

with strict inequality unless $f(t) = \text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right]$. By Lemma 5.2, we also have $\text{Cap}_1 \left(\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right) = \partial_t f(0) \left(\frac{n}{n-1}\right)^{n-1}$. Since

$$\begin{aligned} \frac{\text{Cap}_1 \left(\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right)}{\text{Cap}_1 \left(\text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right] \right)} &= \frac{\frac{\partial_t f(0)}{n} \text{Cap}_1((1+t)^n)}{\frac{\partial_t f(0)}{n} \text{Cap}_1(\text{trunc}_d[(1+t)^n])} \\ &= \frac{\text{Cap}_1((1+t)^n)}{\text{Cap}_1(\text{trunc}_d[(1+t)^n])} \\ &= \gamma_{n,d}, \end{aligned}$$

we therefore have

$$\begin{aligned} \partial_t f(0) \left(\frac{n}{n-1}\right)^{n-1} &= \text{Cap}_1 \left(\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right) \\ &= \gamma_{n,d} \text{Cap}_1 \left(\text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t\right)^n \right] \right) \\ &\geq \gamma_{n,d} \text{Cap}_1(f), \end{aligned}$$

with strict inequality unless $f(t) = \text{trunc}_d \left[\left(1 + \frac{\partial_t f(0)}{n} t \right)^n \right]$. By rearranging, this implies (1) and also the $(\implies)$ direction of (3). $\square$

5.2 Support and Capacity of SLC Polynomials

In this section we prove various results on the interaction between support, capacity bounds, and capacity minimizers for SLC polynomials. The main result of this section is [Corollary 5.7](#), which gives a capacity-based sufficient condition for an SLC polynomial to have full support. Note that we do not directly use [Corollary 5.7](#) in [Section 5.4](#), but instead give the argument used in the proof. We still give [Corollary 5.7](#) explicitly to unify this section.

First is a more refined version of a derivative capacity bound due to Gurvits, which was originally used to prove [Theorem 2.10](#).

Theorem 5.5. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous and SLC, and let d_i be the degree of x_i in p . If $\partial_{x_1} \cdots \partial_{x_n} p(\mathbf{x}) > 0$ then*

$$\text{Cap}_1(\partial_{x_i}|_{x_i=0} p(\mathbf{x})) \geq \gamma_{n,d_i} \left(\frac{n-1}{n} \right)^{n-1} \text{Cap}_1(p),$$

where $\gamma_{n,d_i} \geq 1$ is as defined in [Theorem 5.4](#).

Proof. Without loss of generality, we assume $i = n$. Thus for all $\mathbf{x} \in \mathbb{R}_{>0}^{n-1}$, the polynomial $f(t) = p(x_1, \dots, x_{n-1}, t)$ is of degree d_n . Further, $\partial_t f(0) > 0$ by assumption, and f is n -Newton by [Lemma 2.9](#).

With this, by [Theorem 5.4](#) we have

$$\partial_t f(0) \geq \gamma_{n,d_n} \left(\frac{n-1}{n} \right)^{n-1} \text{Cap}_1(f)$$

which implies

$$\text{Cap}_1(\partial_{x_n}|_{x_n=0} p(\mathbf{x})) \geq \gamma_{n,d_n} \left(\frac{n-1}{n} \right)^{n-1} \text{Cap}_1(p)$$

by taking inf over $\mathbf{x} \in \mathbb{R}_{>0}^{n-1}$ on both sides. $\square$

Now we prove a basic support lemma for SLC polynomials. This can be proven using the fact that SLC polynomials have M-convex support, but we have decided to give a simpler self-contained proof here.

Lemma 5.6. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous and SLC. If the coefficient of x_i^n in p is positive for all $i \in [n]$ then p has full support.*

Proof. We prove this result by induction on n . Fix p and define

$$H_p = \{\boldsymbol{\alpha} \in \Delta_n \cap \mathbb{Z}^n : \langle \mathbf{x}^{\boldsymbol{\alpha}} \rangle p(\mathbf{x}) = 0\}.$$

So as to obtain a contradiction, suppose H_p is non-empty and let $\boldsymbol{\alpha}$ be the vector which maximizes $\max_{i \in [n]} \alpha_i$ over H_p . By possibly permuting variables, we will assume that $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n$.

Case 1: $\alpha_3 = 0$ or $n \leq 2$. In this case $f(t) = p(t, 1, 0, \dots, 0)$ is n -Newton and has strictly positive t^0 and t^n coefficients by assumption. The n -Newton property then implies the t^k coefficient of f is positive for all $0 \leq k \leq n$, which is a contradiction.

Case 2: $\alpha_3 > 0$. In this case $\alpha_1 \leq n - 2$, which implies the $x_i^{n-1}x_j$ coefficient of p is strictly positive for all i, j by our assumption on α . Thus $\partial_{x_1}|_{x_1=0} p(\mathbf{x})$ is $(n-1)$ -variate and $(n-1)$ -homogeneous, with strictly positive x_i^{n-1} coefficients for all $i \in [n]$. The inductive hypothesis then implies the $\mathbf{x}^\alpha$ coefficient of p is strictly positive, which is a contradiction. $\square$

The above two results combine to give the main result of the section.

Corollary 5.7. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous and SLC. If $\partial_{x_1} \cdots \partial_{x_n} p(\mathbf{x}) > 0$ and*

$$\text{Cap}_1(\partial_{x_i}|_{x_i=0} p(\mathbf{x})) = \left(\frac{n-1}{n}\right)^{n-1} \text{Cap}_1(p)$$

for all $i \in [n]$, then p has full support.

Proof. So as to obtain a contradiction, suppose p does not have full support. Then by Lemma 5.6, there exists $i \in [n]$ such that the coefficient of x_i^n is 0 in p . Thus by Theorem 5.5 we have

$$\text{Cap}_1(\partial_{x_i}|_{x_i=0} p(\mathbf{x})) \geq \gamma_{n,d_i} \left(\frac{n-1}{n}\right)^{n-1} \text{Cap}_1(p),$$

where $d_i < n$. Since $\gamma_{n,d_i} > 1$ by Theorem 5.4, this is a contradiction. $\square$

5.3 General Capacity Minimizers

This section is devoted to some basic results on capacity minimizers and uniqueness, most of which can be found in [GS00]. Note that none of these results require the SLC property. Recall that a homogeneous polynomial $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ is **doubly stochastic** if $p(\mathbf{1}) = 1$ and $\nabla p(\mathbf{1}) = \mathbf{1}$.

Lemma 5.8. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous with full support. Then for all $c > \text{Cap}_1(p)$ we have that $K_c := \{\mathbf{x} > 0 : \prod_i x_i = 1, p(\mathbf{x}) \leq c\}$ is compact. In particular, $\text{Cap}_1(p)$ is minimized at some $\mathbf{y} > \mathbf{0}$.*

Proof. Let $p(\mathbf{x}) = \sum_{\kappa} p_{\kappa} \mathbf{x}^{\kappa}$, and let $p_{\min} > 0$ be the minimum coefficient of p . Fix $c > \text{Cap}_1(p)$. For any $i \neq j$ and any $\mathbf{x} \in K_c$ we have

$$c \geq p(\mathbf{x}) = \sum_{\kappa} p_{\kappa} \mathbf{x}^{\kappa} \geq \mathbf{x}^{\mathbf{1}} \cdot \frac{x_i}{x_j} \cdot p_{\mathbf{1}+\mathbf{e}_i-\mathbf{e}_j} \geq \frac{x_i}{x_j} \cdot p_{\min}.$$

Thus $\max_{i,j} \left(\frac{x_i}{x_j} \right) \leq \frac{c}{p_{\min}}$, which implies $x_i \in [\frac{p_{\min}}{c}, \frac{c}{p_{\min}}]$ since $\prod_i x_i = 1$. Therefore

$$K_c = \left\{ \mathbf{x} \in \left[\frac{p_{\min}}{c}, \frac{c}{p_{\min}} \right]^n : \prod_i x_i = 1, p(\mathbf{x}) \leq c \right\},$$

which implies K_c is compact. $\square$

Lemma 5.9. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be such that $p(\mathbf{1}) = 1$. Then p is doubly stochastic if and only if $\mathbf{1}$ minimizes $\text{Cap}_1(p)$.*

Proof. First, $\mathbf{1}$ minimizes $\text{Cap}_1(p)$ if and only if

$$\inf_{\mathbf{t} \in \mathbb{R}^n} [\log p(e^{\mathbf{t}}) - \langle \mathbf{t}, \mathbf{1} \rangle]$$

is minimized at $\mathbf{t} = \mathbf{0}$. Since $\log p(e^{\mathbf{t}}) - \langle \mathbf{t}, \mathbf{1} \rangle$ is convex and smooth, this is equivalent to $\nabla|_{\mathbf{t}=\mathbf{0}} \log p(e^{\mathbf{t}}) = \mathbf{1}$, where $\nabla|_{\mathbf{t}=\mathbf{0}} \log p(e^{\mathbf{t}}) = \nabla p(\mathbf{1})$. This is the same as saying p is doubly stochastic. $\square$

Lemma 5.10. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous with full support. Then there exists $\mathbf{y} > \mathbf{0}$ such that $\frac{p(\mathbf{y} \odot \mathbf{x})}{p(\mathbf{y})}$ is doubly stochastic.*

Proof. By Lemma 5.8, there exists some $\mathbf{y} > \mathbf{0}$ which minimizes $\text{Cap}_1(p)$. Letting $q(\mathbf{x}) = \frac{p(\mathbf{y} \odot \mathbf{x})}{p(\mathbf{y})}$, we have $q(\mathbf{1}) = 1$. Since $\mathbf{y}$ minimizes $\text{Cap}_1(p)$, we have that $\mathbf{1}$ minimizes $\text{Cap}_1(q)$. Therefore q is doubly stochastic by Lemma 5.9. $\square$

Lemma 5.11. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous with full support. If p is doubly stochastic, then p is uniquely minimized over $\{\mathbf{x} > \mathbf{0} : \prod_i x_i = 1\}$ at $\mathbf{x} = \mathbf{1}$.*

Proof. Let $p(\mathbf{x}) = \sum_{\kappa} p_{\kappa} \mathbf{x}^{\kappa}$. Using Jensen's inequality, $p(\mathbf{1}) = 1$ implies

$$\log p(\mathbf{x}) = \log \left(\sum_{\kappa} p_{\kappa} \mathbf{x}^{\kappa} \right) \geq \sum_{\kappa} p_{\kappa} \log (\mathbf{x}^{\kappa}).$$

Since $\log$ is strictly concave, this inequality is strict unless $\mathbf{x}^{\kappa} = \mathbf{x}^{\kappa'}$ for all κ, κ' . Thus the above inequality is strict unless $\mathbf{x} = \mathbf{1}$. We now further have

$$\sum_{\kappa} p_{\kappa} \log (\mathbf{x}^{\kappa}) = \sum_{i=1}^n \log x_i \sum_{\kappa} \kappa_i p_{\kappa} = \sum_{i=1}^n \log x_i \cdot \partial_{x_i} p(\mathbf{1}) = 0,$$

since $\partial_{x_i} p(\mathbf{1}) = 1$ by double stochasticity and $\sum_{i=1}^n \log x_i = 0$. Therefore, $p(\mathbf{x}) \geq 1$ with strict inequality unless $\mathbf{x} = \mathbf{1}$. $\square$

Corollary 5.12. *Let $p \in \mathbb{R}_{\geq 0}[x_1, \dots, x_n]$ be n -homogeneous with full support. Then p is uniquely minimized over $\{\mathbf{x} > \mathbf{0} : \prod_i x_i = 1\}$.*

Proof. By [Lemma 5.10](#), there exists $\mathbf{y} > \mathbf{0}$ such that $q(\mathbf{x}) = \frac{p(\mathbf{y} \odot \mathbf{x})}{p(\mathbf{y})}$ is doubly stochastic. By [Lemma 5.11](#), q is uniquely minimized over $\{\mathbf{x} > 0 : \prod_i x_i = 1\}$ at $\mathbf{x} = \mathbf{1}$. Therefore p is uniquely minimized over $\{\mathbf{x} > 0 : \prod_i x_i = 1\}$ at $\mathbf{x} = \mathbf{y}$. $\square$

As a final remark, we note that [Corollary 5.12](#) can be strengthened to say that we have unique minimization if and only if p has full support. In fact, by the proof of [Lemma 5.8](#), the support condition can be weakened to just the coefficients “nearby” to the all-ones vector. These facts were already proven in the hyperbolic case in [\[Gur06a\]](#).

5.4 Proof of [Theorem 5.1](#)

Let $p(\mathbf{x})$ be an n -variate n -homogeneous SLC polynomial for which $\partial_{x_1} \cdots \partial_{x_n} p = \frac{n!}{n^n} \text{Cap}_1(p) > 0$. Define $p_i(\mathbf{x}) = \partial_{x_i}|_{x_i=0} p(\mathbf{x})$ for all $i \in [n]$, let d_i denote the degree of x_i in $p(\mathbf{x})$, and let $\mathbf{v} = \frac{\nabla p(\mathbf{1})}{n}$. Our goal is to prove that $p(\mathbf{x}) = (\sum_i v_i x_i)^n$.

Step 1: Prove p has full support. By [Theorem 5.5](#) we have

$$\text{Cap}_1(p_i) \geq \gamma_{n,d_i} \left(\frac{n-1}{n} \right)^{n-1} \text{Cap}_1(p)$$

for all $i \in [n]$, where $\gamma_{n,d_i} \geq 1$ is as defined in [Theorem 5.4](#). Thus for $i = n$, by iteratively using [Theorem 5.5](#) we obtain

$$\begin{aligned} \frac{n!}{n^n} \text{Cap}_1(p) &= \partial_{x_1} \cdots \partial_{x_n} p \\ &= \text{Cap}_1(\partial_{x_2}|_{x_2=0} \cdots \partial_{x_n}|_{x_n=0} p) \\ &\geq \left(\frac{1}{2} \right)^1 \text{Cap}_1(\partial_{x_3}|_{x_3=0} \cdots \partial_{x_n}|_{x_n=0} p) \\ &\geq \left(\frac{1}{2} \right)^1 \left(\frac{2}{3} \right)^2 \text{Cap}_1(\partial_{x_4}|_{x_4=0} \cdots \partial_{x_n}|_{x_n=0} p) \\ &\geq \cdots \\ &\geq \left[\prod_{k=1}^{n-1} \left(\frac{k-1}{k} \right)^{k-1} \right] \text{Cap}_1(\partial_{x_n}|_{x_n=0} p) \\ &\geq \gamma_{n,d_n} \left[\prod_{k=1}^n \left(\frac{k-1}{k} \right)^{k-1} \right] \text{Cap}_1(p) \\ &= \gamma_{n,d_n} \cdot \frac{n!}{n^n} \text{Cap}_1(p) \\ &\geq \frac{n!}{n^n} \text{Cap}_1(p), \end{aligned}$$

which implies all inequalities are actually equalities. The same argument works for any ordering of the variables, and thus $\gamma_{n,d_i} = 1$ and

$$\text{Cap}_1(p_i) = \left(\frac{n-1}{n}\right)^{n-1} \text{Cap}_1(p)$$

for all $i \in [n]$. Since $\gamma_{n,d_i} = 1$ implies $d_i = n$ by [Theorem 5.4](#), we therefore have that p has full support by [Lemma 5.6](#).

Step 2: Reduce to doubly stochastic p . By [Lemma 5.10](#), there exists $\mathbf{y} > 0$ such that $q(\mathbf{x}) := \frac{p(\mathbf{y} \cdot \mathbf{x})}{p(\mathbf{y})}$ is doubly stochastic. Therefore by assumption we have

$$\frac{p(\mathbf{y})}{y_1 \cdots y_n} \cdot \partial_{x_1} \cdots \partial_{x_n} q(\mathbf{x}) = \partial_{x_1} \cdots \partial_{x_n} p(\mathbf{x}) = \frac{n!}{n^n} \text{Cap}_1(p).$$

By [Theorem 2.10](#) and [Lemma 5.9](#), we also have

$$\partial_{x_1} \cdots \partial_{x_n} q(\mathbf{x}) \geq \frac{n!}{n^n}$$

since q is doubly stochastic. Combining these gives

$$\frac{n!}{n^n} \text{Cap}_1(p) = \frac{p(\mathbf{y})}{y_1 \cdots y_n} \cdot \partial_{x_1} \cdots \partial_{x_n} q(\mathbf{x}) \geq \frac{n!}{n^n} \cdot \frac{p(\mathbf{y})}{y_1 \cdots y_n} \geq \frac{n!}{n^n} \text{Cap}_1(p),$$

and thus these inequalities must actually be equalities. Therefore

$$\partial_{x_1} \cdots \partial_{x_n} q(\mathbf{x}) = \frac{n!}{n^n},$$

and if the theorem holds for doubly stochastic q , we would obtain $q(\mathbf{x}) = \left(\frac{1}{n} \sum_i x_i\right)^n$. Thus

$$p(\mathbf{x}) = p(\mathbf{y}) \cdot q(\mathbf{y}^{-1} \odot \mathbf{x}) = p(\mathbf{y}) \cdot \left(\frac{1}{n} \sum_{i=1}^n \frac{x_i}{y_i}\right)^n,$$

which implies p is a power of a linear form. Therefore we must have $p(\mathbf{x}) = (\sum_i v_i x_i)^n$.

Step 3: Prove $p(e_i) = \frac{1}{n^n}$ for all i . From now on we assume that p is doubly stochastic. By [Lemma 5.11](#), $\text{Cap}_1(p)$ is uniquely minimized over $\{\mathbf{x} > 0 : \prod_i x_i = 1\}$ at $\mathbf{1}$. Similarly, since p_n also has full support and is $(n-1)$ -variate and $(n-1)$ -homogeneous, we have that $\text{Cap}_1(p_n)$ is also uniquely minimized over $\{\mathbf{x} > 0 : \prod_i x_i = 1\}$ at some $\mathbf{y}$ by [Corollary 5.12](#).

Now $f(t) = p(y_1, \dots, y_{n-1}, t)$ is n -Newton by [Lemma 2.9](#). Thus by [Theorem 5.4](#) we have

$$\begin{aligned} \text{Cap}_{\mathbf{1}}(p_n) &= \partial_t f(0) \\ &\geq \left(\frac{n-1}{n}\right)^{n-1} \text{Cap}_{\mathbf{1}}(f) \\ &\geq \left(\frac{n-1}{n}\right)^{n-1} \text{Cap}_{\mathbf{1}}(p) \\ &= \text{Cap}_{\mathbf{1}}(p_n), \end{aligned}$$

which implies all inequalities are actually equalities. In particular,

$$\text{Cap}_{\mathbf{1}}(p) = \text{Cap}_{\mathbf{1}}(f) = \text{Cap}_{\mathbf{1}}(p(y_1, \dots, y_{n-1}, t)),$$

and since $\text{Cap}_{\mathbf{1}}(p)$ is optimized uniquely at $\mathbf{1}$ over vectors with product equal to 1, we have $\mathbf{y} = \mathbf{1}$ as well. By the above equalities, we also have

$$\partial_t f(0) = \left(\frac{n-1}{n}\right)^{n-1} \text{Cap}_{\mathbf{1}}(f)$$

which implies $f(t) = (a + bt)^n$ for some $a, b > 0$ by [Theorem 5.4](#). Since $f(1) = p(\mathbf{1}) = 1$ and $\partial_t f(0) = \partial_{x_n}|_{x_n=0} p(\mathbf{1}) = 1$ by double stochasticity, we have that $a = \frac{n-1}{n}$ and $b = \frac{1}{n}$. Therefore $p(\mathbf{e}_n) = \langle t^n \rangle f(t) = \frac{1}{n^n}$. Applying a similar argument to p_i implies $p(\mathbf{e}_i) = \frac{1}{n^n}$ for all $i \in [n]$.

Step 4: Prove $p(\mathbf{x}) = \left(\frac{x_1 + \dots + x_n}{n}\right)^n$. Since p is a homogeneous polynomial, it is sufficient to prove the equality on the standard simplex Δ_n . By concavity of $p^{1/n}$ (see [Corollary 2.6](#)), for all $\mathbf{x} \in \Delta_n$ we have

$$p^{1/n}(\mathbf{x}) \geq \sum_{i=1}^n x_i p^{1/n}(\mathbf{e}_i) = \frac{1}{n} \sum_{i=1}^n x_i.$$

Concavity of $p^{1/n}$ also implies

$$p^{1/n}(\mathbf{x}) \leq p^{1/n}(\mathbf{1}) + \nabla p^{1/n}(\mathbf{1})^\top (\mathbf{x} - \mathbf{1}) = 1 + \frac{1}{n} \mathbf{1}^\top (\mathbf{x} - \mathbf{1}) = \frac{1}{n} \sum_{i=1}^n x_i.$$

Therefore we have

$$\left(\frac{x_1 + \dots + x_n}{n}\right)^n \leq p(\mathbf{x}) \leq \left(\frac{x_1 + \dots + x_n}{n}\right)^n.$$

Since this holds for all $\mathbf{x} \in \Delta_n$, we thus have that $p(\mathbf{x}) = \left(\frac{x_1 + \dots + x_n}{n}\right)^n$ for all $\mathbf{x}$.

6 Failure of Rank-One Minimizers

In this section, we prove our main counter results to simple minimizer uniqueness.

6.1 Permanent, with both marginals non-uniform

Fix positive $\mathbf{r}, \mathbf{c} \neq \mathbf{1}$ such that $\|\mathbf{r}\|_1 = \|\mathbf{c}\|_1 = n$, and consider $\text{Mat}_n(\mathbf{r}, \mathbf{c})$ which is the set of all $n \times n$ matrices with non-negative entries and row sums $\mathbf{r}$ and column sums $\mathbf{c}$. Note that this set contains the matrix $\frac{1}{n}\mathbf{r}\mathbf{c}^\top$. We will now prove [Proposition 1.5](#), which says that $\frac{1}{n}\mathbf{r}\mathbf{c}^\top$ never minimizes the permanent over $\text{Mat}_n(\mathbf{r}, \mathbf{c})$ in this case.

Consider the polynomial $\text{per}(X)$ on a matrix of variables. The gradient of this polynomial can be expressed as a matrix, with entries

$$\nabla \text{per}(X)_{ij} = \text{per}(X_{ij}),$$

where X_{ij} is the matrix X with row i and column j removed. In particular, we have

$$\nabla \text{per}\left(\frac{1}{n}\mathbf{r}\mathbf{c}^\top\right) = \frac{1}{n^n} \prod_{k \neq i} r_k \prod_{k \neq j} c_k = \text{per}\left(\frac{1}{n}\mathbf{r}\mathbf{c}^\top\right) \cdot \frac{1}{r_i c_j}.$$

Now, any minimizer of the permanent which lies in the relative interior of $\text{Mat}_n(\mathbf{r}, \mathbf{c})$ must have gradient in $W^\perp$ where W is the linear space parallel to the affine span of $\text{Mat}_n(\mathbf{r}, \mathbf{c})$. The space W consists of all matrices with row and column sums all equal to 0, and is of dimension $n^2 - (2n - 1)$. Thus $W^\perp$ consists of all matrices M with entries of the form

$$m_{ij} = a_i + b_j$$

for some $a_i, b_j \in \mathbb{R}$. (Note that the dimension of this space of matrices is $2n - 1$.)

Now we prove [Proposition 1.5](#). Since $\frac{1}{n}\mathbf{r}\mathbf{c}^\top$ lies in the relative interior of $\text{Mat}_n(\mathbf{r}, \mathbf{c})$, if it is a minimizer then we need $\nabla \text{per}\left(\frac{1}{n}\mathbf{r}\mathbf{c}^\top\right) \in W^\perp$ which implies

$$\frac{1}{r_i c_j} = a_i + b_j \quad \text{for all } i, j,$$

for some $a_i, b_j \in \mathbb{R}$. This cannot happen if $\mathbf{r}, \mathbf{c} \neq \mathbf{1}$, by the following result applied to $\mathbf{x} = \mathbf{a}$, $\mathbf{y} = \mathbf{1}$, and $\mathbf{z} = \mathbf{b}$.

Lemma 6.1. *If for non-zero vectors $\mathbf{x}, \mathbf{y}, \mathbf{z}$ the matrix $\mathbf{x}\mathbf{y}^\top + \mathbf{y}\mathbf{z}^\top$ is rank one, then one of $\mathbf{x}, \mathbf{z}$ is collinear to $\mathbf{y}$.*

Proof. Suppose $\mathbf{y}, \mathbf{z}$ are linearly independent. Since $\mathbf{x}\mathbf{y}^\top + \mathbf{y}\mathbf{z}^\top$ is rank one, there is some t, s not both zero such that $(\mathbf{x}\mathbf{y}^\top + \mathbf{y}\mathbf{z}^\top)(t\mathbf{y} + s\mathbf{z}) = \mathbf{0}$. This implies

$$\mathbf{0} = (t\mathbf{y}^\top \mathbf{y} + s\mathbf{y}^\top \mathbf{z})\mathbf{x} + (t\mathbf{z}^\top \mathbf{y} + s\mathbf{z}^\top \mathbf{z})\mathbf{y}.$$

Thus if $\mathbf{x}, \mathbf{y}$ are linearly independent, then we have

$$t\mathbf{y}^\top \mathbf{y} + s\mathbf{y}^\top \mathbf{z} = 0 = t\mathbf{z}^\top \mathbf{y} + s\mathbf{z}^\top \mathbf{z}.$$

Therefore

$$0 = t(t\mathbf{y}^\top \mathbf{y} + s\mathbf{y}^\top \mathbf{z}) + s(t\mathbf{z}^\top \mathbf{y} + s\mathbf{z}^\top \mathbf{z}) = (t\mathbf{y} + s\mathbf{z})^\top (t\mathbf{y} + s\mathbf{z}) > 0,$$

a contradiction. Thus either $\mathbf{x}, \mathbf{y}$ are linearly dependent, or else $\mathbf{y}, \mathbf{z}$ are linearly dependent. $\square$

6.2 Permanent, with marginals far from all-ones

We now study an example which proves [Proposition 1.6](#). Fix $t > 0$ and $n \in \mathbb{N}$, and define $\epsilon := \frac{1}{n^{1+t}}$. Further define $\alpha \in \mathbb{R}_{>0}^n$ and $\mathbf{c} \in \mathbb{R}_{>0}^{n+1}$ via

$$\alpha := (1-\epsilon, 1-\epsilon, \dots, 1-\epsilon) \in \mathbb{R}_{>0}^n \quad \text{and} \quad \mathbf{c} := (1+\alpha_1, \alpha_2, \alpha_3, \dots, \alpha_n, \sum_{j=1}^n (1-\alpha_j)) \in \mathbb{R}_{>0}^{n+1}.$$

Note that $\|\mathbf{c} - \mathbf{1}\|_1 = 1 + (n-2)\epsilon + 1 - n\epsilon = 2(1-\epsilon) < 2$, which by [Theorem 2.13](#) implies $\text{per}(X) > 0$ for all $X \in \text{Mat}_{n+1}(\mathbf{c})$. We first have

$$\text{per}\left(\frac{1}{n+1} \mathbf{1} \cdot \mathbf{c}^\top\right) = \frac{(n+1)!}{(n+1)^{n+1}} (2-\epsilon)(1-\epsilon)^{n-1} n\epsilon = \frac{n!}{n^{(1+t)n}} \cdot \frac{2 - n^{-1-t}}{1 - n^{-1-t}} \cdot \frac{(n^{1+t} - 1)^n}{n^t (n+1)^n}.$$

Now consider the matrix

$$M = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 1 - \sum_{j=1}^1 (1 - \alpha_j) & \sum_{j=1}^1 (1 - \alpha_j) & 0 & \cdots & 0 & 0 \\ 0 & 1 - \sum_{j=1}^2 (1 - \alpha_j) & \sum_{j=1}^2 (1 - \alpha_j) & \cdots & 0 & 0 \\ 0 & 0 & 1 - \sum_{j=1}^3 (1 - \alpha_j) & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \sum_{j=1}^{n-1} (1 - \alpha_j) & 0 \\ 0 & 0 & 0 & \cdots & 1 - \sum_{j=1}^n (1 - \alpha_j) & \sum_{j=1}^n (1 - \alpha_j) \end{bmatrix}.$$

Note that $M \in \text{Mat}_{n+1}(\mathbf{c})$. The matrix M is upper-triangular, and thus we have

$$\text{per}(M) = \prod_{k=1}^n \sum_{j=1}^k (1 - \alpha_j) = \prod_{k=1}^n (k\epsilon) = n! \cdot \epsilon^n = \frac{n!}{n^{(1+t)n}}.$$

We then further have

$$\frac{2 - n^{-1-t}}{1 - n^{-1-t}} \cdot \frac{(n^{1+t} - 1)^n}{n^t (n+1)^n} \approx 2n^{(n-1)t} \cdot \frac{(n - n^{-t})^n}{(n+1)^n} \geq 2n^{(n-1)t} \cdot \frac{(n-1)^n}{(n+1)^n} \approx \frac{2n^{(n-1)t}}{e^2} > 1,$$

which implies $\text{per}(M) < \text{per}\left(\frac{1}{n+1} \mathbf{1} \cdot \mathbf{c}^\top\right)$ for large enough n .

References

- [Ana+18] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant. “Log-concave polynomials III: Mason’s ultra-log-concavity conjecture for independent sets of matroids”. In: *arXiv preprint arXiv:1811.01600* (2018) (cit. on pp. [3](#), [6](#)).
- [AO17] Nima Anari and Shayan Oveis Gharan. “A generalization of permanent inequalities and applications in counting and optimization”. In: *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*. 2017, pp. 384–396 (cit. on p. [5](#)).

- [BH20] Petter Brändén and June Huh. “Lorentzian polynomials”. In: *Annals of Mathematics* 192.3 (2020), pp. 821–891 (cit. on pp. 3, 6).
- [Bru85] Richard A. Brualdi. “An interesting face of the polytope of doubly stochastic matrices”. In: *Linear and Multilinear Algebra* 17 (1985), pp. 5–18. DOI: [10.1080/03081088508817637](https://doi.org/10.1080/03081088508817637) (cit. on p. 5).
- [CW05] Gi-Sang Cheon and Ian M. Wanless. “An update on Minc’s survey of open problems involving permanents”. In: *Linear Algebra and its Applications* 403.1–3 (2005), pp. 314–342. DOI: [10.1016/j.laa.2005.02.030](https://doi.org/10.1016/j.laa.2005.02.030) (cit. on p. 5).
- [Ego81] Gregory P Egorychev. “The solution of van der Waerden’s problem for permanents”. In: *Advances in Mathematics* 42.3 (1981), pp. 299–305 (cit. on p. 1).
- [Fal81] Dmitry I Falikman. “Proof of the van der Waerden conjecture regarding the permanent of a doubly stochastic matrix”. In: *Mathematical notes of the Academy of Sciences of the USSR* 29 (1981), pp. 475–479 (cit. on p. 1).
- [For80] Thomas H. Foregger. “On the minimum value of the permanent of a nearly decomposable doubly stochastic matrix”. In: *Linear Algebra and its Applications* 32 (1980), pp. 75–85. DOI: [10.1016/0024-3795\(80\)90008-7](https://doi.org/10.1016/0024-3795(80)90008-7) (cit. on p. 5).
- [FS86] Thomas H. Foregger and Richard Sinkhorn. “On matrices minimizing the permanent on faces of the polyhedron of the doubly stochastic matrices”. In: *Linear and Multilinear Algebra* 19.4 (1986), pp. 395–397. DOI: [10.1080/03081088608817734](https://doi.org/10.1080/03081088608817734) (cit. on p. 5).
- [Gar+20] Ankit Garg, Leonid Gurvits, Rafael Oliveira, and Avi Wigderson. “Operator scaling: Theory and applications”. In: *Foundations of Computational Mathematics* 20.2 (2020), pp. 223–290. DOI: [10.1007/s10208-019-09417-z](https://doi.org/10.1007/s10208-019-09417-z) (cit. on p. 5).
- [GKL24] Leonid Gurvits, Nathan Klein, and Jonathan Leake. “From Trees to Polynomials and Back Again: New Capacity Bounds with Applications to TSP”. In: *51st International Colloquium on Automata, Languages, and Programming (ICALP 2024)*. Schloss Dagstuhl–Leibniz-Zentrum für Informatik. 2024, pp. 79–1 (cit. on pp. 2, 4).
- [GL21] Leonid Gurvits and Jonathan Leake. “Capacity lower bounds via productization”. In: *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*. 2021, pp. 847–858 (cit. on p. 8).
- [GS00] Leonid Gurvits and Alex Samorodnitsky. “A deterministic polynomial-time algorithm for approximating mixed discriminant and mixed volume”. In: *Proceedings of the thirty-second annual ACM symposium on Theory of computing*. 2000, pp. 48–57 (cit. on p. 17).

- [Gur04] Leonid Gurvits. “Classical complexity and quantum entanglement”. In: *Journal of Computer and System Sciences* 69.3 (2004), pp. 448–484. DOI: [10.1016/j.jcss.2004.06.003](https://doi.org/10.1016/j.jcss.2004.06.003) (cit. on p. 5).
- [Gur06a] Leonid Gurvits. “Hyperbolic polynomials approach to Van der Waerden/Schrijver-Valiant like conjectures: sharper bounds, simpler proofs and algorithmic applications”. In: *Proceedings of the thirty-eighth annual ACM symposium on Theory of computing*. 2006, pp. 417–426 (cit. on pp. 2, 3, 13, 19).
- [Gur06b] Leonid Gurvits. “The van der Waerden conjecture for mixed discriminants”. In: *Advances in Mathematics* 200.2 (2006), pp. 435–454 (cit. on pp. 2, 3, 5).
- [Gur08] Leonid Gurvits. “Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all”. In: *The Electronic Journal of Combinatorics* 15.1 (2008), R66. DOI: [10.37236/790](https://doi.org/10.37236/790) (cit. on pp. 1, 2).
- [Gur09a] Leonid Gurvits. “A polynomial-time algorithm to approximate the mixed volume within a simply exponential factor”. In: *Discrete & Computational Geometry* 41 (2009), pp. 533–555 (cit. on pp. 2, 4).
- [Gur09b] Leonid Gurvits. “On multivariate Newton-like inequalities”. In: *Advances in Combinatorial Mathematics: Proceedings of the Waterloo Workshop in Computer Algebra 2008*. Springer. 2009, pp. 61–78 (cit. on pp. 3, 5, 7).
- [Gur15] Leonid Gurvits. “Boolean matrices with prescribed row/column sums and stable homogeneous polynomials: Combinatorial and algorithmic applications”. In: *Information and Computation* 240 (2015), pp. 42–55 (cit. on p. 8).
- [Hwa85] Suk-Geun Hwang. “Minimum permanent on faces of staircase type of the polytope of doubly stochastic matrices”. In: *Linear and Multilinear Algebra* 18.4 (1985), pp. 271–306. DOI: [10.1080/03081088508817694](https://doi.org/10.1080/03081088508817694) (cit. on p. 5).
- [KKO21] Anna R. Karlin, Nathan Klein, and Shayan Oveis Gharan. “A (Slightly) Improved Approximation Algorithm for Metric TSP”. In: *STOC*. ACM, 2021 (cit. on p. 2).
- [LSW98] Nathan Linial, Alex Samorodnitsky, and Avi Wigderson. “A deterministic strongly polynomial algorithm for matrix scaling and approximate permanents”. In: *Proceedings of the thirtieth annual ACM symposium on Theory of computing*. 1998, pp. 644–652 (cit. on p. 2).
- [Min84] Henryk Minc. “Minimum permanents of doubly stochastic matrices with prescribed zero entries”. In: *Linear and Multilinear Algebra* 15 (1984), pp. 225–243. DOI: [10.1080/03081088408817592](https://doi.org/10.1080/03081088408817592) (cit. on p. 5).

- [MOA11] Albert W. Marshall, Ingram Olkin, and Barry C. Arnold. *Inequalities: Theory of Majorization and Its Applications*. 2nd ed. Springer Series in Statistics. New York, NY: Springer, 2011. ISBN: 978-0-387-40087-7. DOI: [10.1007/978-0-387-68276-1](https://doi.org/10.1007/978-0-387-68276-1) (cit. on p. 12).
- [PSW11] Kyle Pula, Seok-Zun Song, and Ian M. Wanless. “Minimum permanents on two faces of the polytope of doubly stochastic matrices”. In: *Linear Algebra and its Applications* 434 (2011), pp. 232–238. DOI: [10.1016/j.laa.2010.08.015](https://doi.org/10.1016/j.laa.2010.08.015) (cit. on p. 5).
- [SB24] Seok-Zun Song and LeRoy B. Beasley. “A face of the polytope of doubly stochastic matrices”. In: *Linear and Multilinear Algebra* 72.10 (2024), pp. 1612–1624. DOI: [10.1080/03081087.2023.2190073](https://doi.org/10.1080/03081087.2023.2190073) (cit. on p. 5).
- [Son+97] Seok-Zun Song, Sung-Min Hong, Young-Bae Jun, and Seon-Jeong Kim. “Minimum permanents on a face of the polytope of doubly stochastic matrices”. In: *Linear Algebra and its Applications* 259 (1997), pp. 169–182. DOI: [10.1016/S0024-3795\(96\)00248-0](https://doi.org/10.1016/S0024-3795(96)00248-0) (cit. on p. 5).